

## Research Article

# Notes on the Propagators of Evolution Equations

Yu Lin,<sup>1</sup> Ti-Jun Xiao,<sup>2</sup> and Jin Liang<sup>3</sup>

<sup>1</sup> Department of Mathematics, University of Science and Technology of China, Hefei, Anhui 230026, China

<sup>2</sup> Shanghai Key Laboratory for Contemporary Applied Mathematics, School of Mathematical Sciences, Fudan University, Shanghai 200433, China

<sup>3</sup> Department of Mathematics, Shanghai Jiao Tong University, Shanghai 200240, China

Correspondence should be addressed to Ti-Jun Xiao, xiaotj@ustc.edu.cn

Received 8 January 2010; Accepted 1 February 2010

Academic Editor: Gaston Mandata N'Guerekata

Copyright © 2010 Yu Lin et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We consider the propagator of an evolution equation, which is a semigroup of linear operators. Questions related to its operator norm function and its behavior at the *critical point* for norm continuity or compactness or differentiability are studied.

## 1. Introduction

As it is well known, each well-posed Cauchy problem for first-order evolution equation in Banach spaces

$$\begin{aligned}u'(t) &= Au(t), \quad t \geq 0, \\u(0) &= u_0\end{aligned}\tag{1.1}$$

gives rise to a well-defined propagator, which is a semigroup of linear operators, and the theory of semigroups of linear operators on Banach spaces has developed quite rapidly since the discovery of the generation theorem by Hille and Yosida in 1948. By now, it is a rich theory with substantial applications to many fields (cf., e.g., [1–6]).

In this paper, we pay attention to some basic problems on the semigroups of linear operators and reveal some essential properties of theirs.

Let  $X$  be a Banach space.

*Definition 1.1* (see [1–6]). *A one-parameter family  $(T(t))_{t \geq 0}$  of bounded linear operators on  $X$  is called a strongly continuous semigroup (or simply  $C_0$ -semigroup) if it satisfies the following conditions:*

- (i)  $T(0) = I$ , with ( $I$  being the identity operator on  $X$ ),
- (ii)  $T(t + s) = T(t)T(s)$  for  $t, s \geq 0$ ,
- (iii) the map  $t \rightarrow T(t)x$  is continuous on  $[0, \infty)$  for every  $x \in X$ .

The *infinitesimal generator*  $A$  of  $(T(t))_{t \geq 0}$  is defined as

$$Ax = \lim_{t \rightarrow 0^+} \frac{T(t)x - x}{t} \quad (1.2)$$

with domain

$$D(A) = \left\{ x \in X; \lim_{t \rightarrow 0^+} \frac{T(t)x - x}{t} \text{ exists} \right\}. \quad (1.3)$$

For a comprehensive theory of  $C_0$ -semigroups we refer to [2].

## 2. Properties of the Function $t \mapsto \|T(t)\|$

Let  $(T(t))_{t \geq 0}$  be a  $C_0$ -semigroup on  $X$  and define  $g(t) := \|T(t)\|$  for  $t \geq 0$ . Clearly, from Definition 1.1 we see that

- (I)  $g(0) = 1$ ,  $g(t) \geq 0$  for  $t \geq 0$ ;
- (II)  $g(t + s) \leq g(t)g(s)$  for  $t, s \geq 0$ .

Furthermore, we can infer from the strong continuity of  $(T(t))_{t \geq 0}$  that

- (III)  $g(t)$  is lower-semicontinuous, that is,

$$g(t) \leq \liminf_{s \rightarrow t} g(s). \quad (2.1)$$

In fact,

$$\|T(t)x\| = \liminf_{s \rightarrow t} \|T(s)x\| \leq \liminf_{s \rightarrow t} \sup_{\|x\|=1} \|T(s)x\| = \liminf_{s \rightarrow t} g(s) \quad (2.2)$$

holds for all  $x \in X$  with  $\|x\| = 1$ . Thus, taking the supremum for all  $x \in X$  with  $\|x\| = 1$  on the left-hand side leads to (2.1).

We ask the following question

*For every function  $g(\cdot)$  satisfying (I), (II), and (III), does there exist a  $C_0$  semigroup  $(T(t))_{t \geq 0}$  on some Banach space  $X$  such that  $\|T(t)\| = g(t)$  for all  $t \geq 0$ ?*

We show that this is not true even if  $X$  is a finite-dimensional space.

**Theorem 2.1.** *Let  $X$  be an  $n$ -dimensional Banach space with  $2 \leq n < \infty$ . Let*

$$g(t) = \begin{cases} 1 + t & \text{if } 0 \leq t \leq 2^n - 1, \\ \left(\frac{2}{2^n - 1}\right)^n t^n & \text{if } t > 2^n - 1. \end{cases} \quad (2.3)$$

Then  $g(\cdot)$  satisfies (I), (II), and (III), and there exists no  $C_0$  semigroup  $(T(t))_{t \geq 0}$  on  $X$  such that  $\|T(t)\| = g(t)$  for all  $t \geq 0$ .

*Proof.* First, we show that  $g(\cdot)$  satisfies (I), (II), and (III).

(I) is clearly satisfied.

To show (III) and (II), we write

$$\gamma := 2^n - 1. \quad (2.4)$$

Then

$$\lim_{t \rightarrow \gamma^-} g(t) = g(\gamma) = 1 + \gamma = 2^n = \lim_{t \rightarrow \gamma^+} g(t), \quad (2.5)$$

hence  $g(\cdot)$  satisfies (III).

For (II), suppose  $t, s \geq 0$ , and consider the following four cases.

*Case 1* ( $t \geq \gamma$  and  $s \geq \gamma$ ). In this case

$$\frac{g(t+s)}{g(t)g(s)} = \left(\frac{\gamma}{2}\right)^n \left(\frac{1}{t} + \frac{1}{s}\right)^n \leq \left(\frac{\gamma}{2}\right)^n \left(\frac{2}{\gamma}\right)^n = 1, \quad (2.6)$$

that is,

$$g(t+s) \leq g(t)g(s). \quad (2.7)$$

*Case 2* ( $t \geq \gamma$  and  $s \leq \gamma$ ). Let

$$h(s) = \left(1 + \frac{s}{\gamma}\right)^n - 1 - s. \quad (2.8)$$

Then

$$h''(s) = \frac{n(n-1)}{\gamma^2} \left(1 + \frac{s}{\gamma}\right)^{n-2} \geq 0, \quad (2.9)$$

and  $h(\cdot)$  is a convex function on  $[0, \gamma]$ . So by Jensen's inequality, we have

$$h(s) \leq \frac{s}{\gamma} h(0) + \frac{\gamma-s}{\gamma} h(\gamma) = \frac{\gamma-s}{\gamma} (2^n - 1 - \gamma) = 0, \quad (2.10)$$

that is,

$$\left(1 + \frac{s}{\gamma}\right)^n \leq 1 + s \quad \text{for } 0 \leq s \leq \gamma. \quad (2.11)$$

Therefore

$$\frac{g(t+s)}{g(t)} = \left(1 + \frac{s}{t}\right)^n \leq \left(1 + \frac{s}{\gamma}\right)^n \leq 1 + s = g(s), \quad (2.12)$$

that is,  $g(t+s) \leq g(t)g(s)$ .

*Case 3* ( $t+s > \gamma$ , but  $t \leq \gamma$  and  $s \leq \gamma$ ). It follows from Case 2 that

$$\begin{aligned} g(t+s) &\leq g(\gamma)g(t+s-\gamma) \\ &= (1+\gamma)(1+t+s-\gamma) \\ &= (1+t)(1+s) - (\gamma-s)(\gamma-t) \\ &\leq (1+t)(1+s) \\ &= g(t)g(s). \end{aligned} \quad (2.13)$$

*Case 4* ( $t+s \leq \gamma$ ). Again we have

$$g(t+s) = 1+t+s \leq 1+t+s+st = (1+t)(1+s) = g(t)g(s). \quad (2.14)$$

Next, we prove that there does not exist any  $C_0$  semigroup  $(T(t))_{t \geq 0}$  on  $X$  such that  $\|T(t)\| = g(t)$ . Suppose  $g(t) = \|T(t)\|$  ( $t \geq 0$ ) for some  $C_0$  semigroup  $(T(t))_{t \geq 0}$  on  $X$  and let  $A$  be its infinitesimal generator.

First we note from (2.3) that

$$\lim_{t \rightarrow +\infty} \|e^{-\varepsilon t} T(t)\| = \lim_{t \rightarrow +\infty} e^{-\varepsilon t} g(t) = 0 \quad (2.15)$$

for every  $\varepsilon > 0$ , while

$$\|T(t)\| = O(t^n) \quad (t \rightarrow +\infty). \quad (2.16)$$

By the well-known Lyapunov theorem [2, Chapter I, Theorem 2.10], all eigenvalues of  $A - \varepsilon I$  (the infinitesimal generator of  $e^{-\varepsilon t} T(t)$ ) have negative parts for every  $\varepsilon > 0$ . Letting  $\lambda_1, \dots, \lambda_r$  be the eigenvalues of  $A$ , we then have

$$\operatorname{Re} \lambda_j - \varepsilon < 0 \quad \text{for every } \varepsilon > 0, j = 1, \dots, r, \quad (2.17)$$

and this implies that

$$\operatorname{Re} \lambda_j \leq 0, \quad j = 1, \dots, r. \tag{2.18}$$

It is known that there is an isomorphism  $P$  of  $\mathbb{C}^n$  onto  $X$  such that

$$A = P \begin{pmatrix} J_1 & \cdots & 0 \\ \vdots & \vdots & \vdots \\ 0 & \cdots & J_r \end{pmatrix} P^{-1}, \tag{2.19}$$

where  $J_j$  is the Jordan block corresponding to  $\lambda_j$ . Therefore

$$T(t) = P \begin{pmatrix} e^{tJ_1} & \cdots & 0 \\ \vdots & \vdots & \vdots \\ 0 & \cdots & e^{tJ_r} \end{pmatrix} P^{-1}. \tag{2.20}$$

Set

$$N_j = J_j - \lambda_j I_{(k_j)}, \quad 1 \leq j \leq r, \tag{2.21}$$

where  $k_j$  is the order of  $J_j$ . Then  $N_j$  is a  $k_j$ th nilpotent matrix with  $k_j \leq n$  for each  $1 \leq j \leq r$ . According to (2.20) and (2.18), we have

$$\begin{aligned} \|T(t)\| &\leq \|P\| \cdot \left\| P^{-1} \right\| \max_{1 \leq j \leq r} \|e^{tJ_j}\| \\ &= \|P\| \cdot \left\| P^{-1} \right\| \max_{1 \leq j \leq r} \|e^{t\lambda_j} e^{tN_j}\| \\ &\leq \|P\| \cdot \left\| P^{-1} \right\| \max_{1 \leq j \leq r} \|e^{tN_j}\|. \end{aligned} \tag{2.22}$$

Observing

$$\begin{aligned} \|e^{tN_i}\| &= \left\| I_{(k_i)} + tN_i + \cdots + \frac{t^{k_i-1}}{(k_i-1)!} N_i^{k_i-1} \right\| \\ &= O\left(t^{k_i-1}\right) \quad (t \rightarrow +\infty), \end{aligned} \tag{2.23}$$

we see that

$$\|e^{tN_i}\| = O(t^{n-1}) \quad (t \rightarrow +\infty). \quad (2.24)$$

Thus,

$$g(t) = \|T(t)\| = O(t^{n-1}) \quad (t \rightarrow +\infty), \quad (2.25)$$

which is a contradiction to (2.16).  $\square$

*Open Problem 1.* Is it possible that there exists an  $X$  with  $\dim X = \infty$  and a  $C_0$  semigroup  $(T(t))_{t \geq 0}$  on  $X$  such that  $\|T(t)\| = g(t)$  for all  $t \geq 0$ ?

### 3. The Critical Point of Norm-Continuous (Compact, Differentiable) Semigroups

The following definitions are basic [1–6].

*Definition 3.1.* A  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  is called norm-continuous for  $t > t_0$  if  $t \mapsto T(t)$  is continuous in the uniform operator topology for  $t > t_0$ .

*Definition 3.2.* A  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  is called compact for  $t > t_0$  if  $(T(t))_{t \geq 0}$  is a compact operator for  $t > t_0$ .

*Definition 3.3.* A  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  is called differentiable for  $t > t_0$  if for every  $x \in X$ ,  $t \mapsto T(t)x$  is differentiable for  $t > t_0$ .

It is known that if a  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  is norm continuous (compact, differentiable) at  $t = t_0$ , then it remains so for all  $t > t_0$ . For instance, the following holds.

**Proposition 3.4.** *If the map  $t \rightarrow T(t)x$  is right differentiable at  $t = t_0$ , then it is also differentiable for  $t > t_0$ .*

Therefore, if we write

$$\begin{aligned} J_n &:= \{t_0 \geq 0; T(t) \text{ is norm-continuous for } t > t_0\}, \\ J_c &:= \{t_0 \geq 0; T(t) \text{ is compact for } t > t_0\}, \\ J_d &:= \{t_0 \geq 0; T(t) \text{ is differentiable for } t > t_0\}, \end{aligned} \quad (3.1)$$

and suppose  $J_n \neq \emptyset$  ( $J_c \neq \emptyset$ ,  $J_d \neq \emptyset$ ), then  $J_n (J_c, J_d)$  takes the form of  $[\tau_0, \infty)$  for a nonnegative real number  $\tau_0$ . In other words, if  $J_n \neq \emptyset$  ( $J_c \neq \emptyset$ ,  $J_d \neq \emptyset$ ), then  $(T(t))_{t \geq 0}$  is norm continuous (compact, differentiable) on the interval  $(\tau_0, \infty)$  but not at any point in  $[0, \tau_0)$ . We call  $\tau_0$  the *critical point* of the norm continuity (compactness, differentiability) of operator semigroup  $(T(t))_{t \geq 0}$ .

A natural question is the following

*Suppose that  $\tau_0$  is the critical point of the norm continuity (compactness, differentiability) of the operator semigroup  $(T(t))_{t \geq 0}$ . Is  $(T(t))_{t \geq 0}$  also norm continuous (compact, differentiable) at  $\tau_0$ ? Of course, concerning norm continuity or differentiability at  $\tau_0$  we only mean right continuity or right differentiability.*

We show that the answer is “yes” in some cases and “no” for other cases.

*Example 3.5. Let  $X = L^2[0, 1]$  and*

$$T(t)f(s) = \begin{cases} f(s+t), & s+t \leq 1, \\ 0, & s+t > 1, \end{cases} \quad t \geq 0. \quad (3.2)$$

Then clearly  $T(t) = 0$  for  $t > 1$ . Moreover,  $(T(t))_{t \geq 0}$  is not norm continuous (not compact, not differentiable) for any  $0 \leq t < 1$  since

$$\|T(t+h) - T(t)\| \geq \|(T(t+h) - T(t))f_h\| = 2 \quad (3.3)$$

for sufficiently small  $h > 0$ , where

$$f_h(s) = \begin{cases} h^{-1/2}, & t+h \leq s \leq t+2h, \\ 0, & \text{otherwise.} \end{cases} \quad (3.4)$$

Therefore, in this case we have  $\tau_0 = 1$ . Since  $T(1) = 0$ , we see that  $T(1)$  is compact and  $(T(t))_{t \geq 0}$  is differentiable at  $t = 1$  from the right.

*Example 3.6. Let*

$$X = c_0 = \left\{ (x_1, \dots, x_n, \dots); x_n \in \mathbb{C}, \lim_{n \rightarrow \infty} x_n \text{ exists} \right\} \quad (3.5)$$

with supremum norm. For any  $x = (x_1, \dots, x_n, \dots) \in X$  set

$$T(t)x = (e^{-t}(\cos(te) + i \sin(te))x_1, \dots, e^{-nt}(\cos(te^n) + i \sin(te^n))x_n, \dots), \quad t \geq 0. \quad (3.6)$$

Then,  $(T(t))_{t \geq 0}$  is compact (hence norm-continuous) for  $t > 0$  since  $T(t)$  is the operator-norm limit of a sequence  $(T_k(t))_{k \in \mathbb{N}}$  of finite-rank operators:

$$T_k(t)x = (y_1(t), \dots, y_n(t), \dots), \quad \text{for each } x = (x_1, \dots, x_n, \dots) \in X, k \in \mathbb{N}, \quad (3.7)$$

where

$$y_n(t) = \begin{cases} e^{-nt}(\cos(te^n) + i \sin(te^n))x_n, & n \leq k, \\ 0, & n > k. \end{cases} \quad (3.8)$$

So the critical point for compactness and norm continuity is  $\tau_0 = 0$ . However, the infinitesimal generator of  $(T(t))_{t \geq 0}$  is given by

$$Ax = ((-1 + ie)x_1, \dots, (-n + ie^n)x_n, \dots) \quad \text{for } x = (x_1, \dots, x_n, \dots) \in D(A) \quad (3.9)$$

with

$$D(A) = \{x \in X; Ax \in X\}. \quad (3.10)$$

In view of that  $A$  is unbounded, we know that  $(T(t))_{t \geq 0}$  is not norm continuous at  $t = 0$ .

For differentiability, we note that  $T(t)x$  is differentiable at  $t = t_0$  if and only if  $T(t_0)x \in D(A)$  for each  $x \in X$ . From

$$\begin{aligned} AT(t)x = & \left( \left( e^{(1-t)}(-\sin(te) + i \cos(te)) - e^{-t}(\cos(te) + i \sin(te)) \right) x_1, \dots, \right. \\ & \left. \left( e^{n(1-t)}(-\sin(te^n) + i \cos(te^n)) - ne^{-nt}(\cos(te^n) + i \sin(te^n)) \right) x_n, \dots \right), \end{aligned} \quad (3.11)$$

$$\text{for each } x = (x_1, \dots, x_n, \dots) \in X, \quad t \geq 0,$$

it follows that when  $t > 1$ ,  $T(t)x \in D(A)$  for every  $x \in X$ . On the other hand, when  $0 \leq t \leq 1$  and  $x$  is any nonzero constant sequence,  $T(t)x \notin D(A)$ . Therefore the critical point for differentiability is  $\tau_0 = 1$ . But  $(T(t))_{t \geq 0}$  is not differentiable at  $t = 1$ .

## Acknowledgments

The authors acknowledge the support from the NSF of China (10771202), the Research Fund for Shanghai Key Laboratory of Modern Applied Mathematics (08DZ2271900), and the Specialized Research Fund for the Doctoral Program of Higher Education of China (2007035805).

## References

- [1] E. B. Davies, *One-Parameter Semigroups*, vol. 15 of *London Mathematical Society Monographs*, Academic Press, London, UK, 1980.
- [2] K.-J. Engel and R. Nagel, *One-Parameter Semigroups for Linear Evolution Equations*, vol. 194 of *Graduate Texts in Mathematics*, Springer, New York, NY, USA, 2000.
- [3] H. O. Fattorini, *The Cauchy Problem*, vol. 18 of *Encyclopedia of Mathematics and Its Applications*, Addison-Wesley, Reading, Mass, USA, 1983.
- [4] J. A. Goldstein, *Semigroups of Linear Operators and Applications*, Oxford Mathematical Monographs, The Clarendon Press/Oxford University Press, New York, NY, USA, 1985.
- [5] E. Hille and R. S. Phillips, *Functional Analysis and Semi-Groups*, American Mathematical Society Colloquium Publications, vol. 31, American Mathematical Society, Providence, RI, USA, 1957.
- [6] A. Pazy, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, vol. 44 of *Applied Mathematical Sciences*, Springer, Berlin, Germany, 1983.