

RESEARCH

Open Access

Existence results for nonlocal boundary value problems of nonlinear fractional q -difference equations

Bashir Ahmad¹, Sotiris K Ntouyas^{2*} and Ioannis K Purnaras²

*Correspondence: sntouyas@uoi.gr

²Department of Mathematics,
University of Ioannina, Ioannina,
451 10, Greece

Full list of author information is
available at the end of the article

Abstract

In this paper, we study a nonlinear fractional q -difference equation with nonlocal boundary conditions. The existence of solutions for the problem is shown by applying some well-known tools of fixed-point theory such as Banach's contraction principle, Krasnoselskii's fixed-point theorem, and the Leray-Schauder nonlinear alternative. Some illustrating examples are also discussed.

MSC: 34A08; 39A05; 39A12; 39A13

Keywords: fractional q -difference equations; nonlocal boundary conditions; existence; Leray-Schauder nonlinear alternative; fixed point

1 Introduction

In recent years, the topic of fractional differential equations has gained considerable attention and has evolved as an interesting and popular field of research. It is mainly due to the fact that several times the tools of fractional calculus are found to be more practical and effective than the corresponding ones of classical calculus in the mathematical modeling of dynamical systems associated with phenomena such as fractals and chaos. In fact, fractional calculus has numerous applications in various disciplines of science and engineering such as mechanics, electricity, chemistry, biology, economics, control theory, signal and image processing, polymer rheology, regular variation in thermodynamics, biophysics, blood flow phenomena, aerodynamics, electrodynamics of complex medium, viscoelasticity and damping, control theory, wave propagation, percolation, identification, fitting of experimental data, *etc.* [1–4]. The development of fractional calculus ranges from the theoretical aspects of existence and uniqueness of solutions to the analytic and numerical methods for finding solutions. For some recent work on fractional differential equations, we refer to [5–13] and the references therein.

The pioneer work on q -difference calculus or *quantum calculus* dates back to Jackson's papers [14, 15], while a systematic treatment of the subject can be found in [16, 17]. For some recent existence results on q -difference equations, see [18–20] and the references therein.

There has also been a growing interest on the subject of discrete fractional equations on the time scale \mathbb{Z} . Some interesting results on the topic can be found in a series of papers [21–29].

Fractional q -difference equations have recently attracted the attention of several researchers. For some earlier work on the topic, we refer to [30] and [31], whereas some recent work on the existence theory of fractional q -difference equations can be found in [32–36]. However, the study of boundary value problems of fractional q -difference equations is at its infancy and much of the work on the topic is yet to be done.

In this paper, we discuss the existence and uniqueness of solutions for the nonlocal boundary value problem of fractional q -difference equations given by

$${}^c D_q^\alpha x(t) = f(t, x(t)), \quad 0 \leq t \leq 1, 1 < \alpha \leq 2, \tag{1.1}$$

$$\alpha_1 x(0) - \beta_1 D_q x(0) = \gamma_1 x(\eta_1), \quad \alpha_2 x(1) + \beta_2 D_q x(1) = \gamma_2 x(\eta_2), \tag{1.2}$$

where $f \in C([0, 1] \times \mathbb{R}, \mathbb{R})$, ${}^c D_q^\alpha$ is the fractional q -derivative of the Caputo type, and $\alpha_i, \beta_i, \gamma_i, \eta_i \in \mathbb{R}$ ($i = 1, 2$).

2 Preliminaries on fractional q -calculus

In this section, we cite some definitions and fundamental results of the q -calculus as well as of the fractional q -calculus [37, 38]. We also give a lemma that will be used in obtaining the main results of the paper.

Let $q \in (0, 1)$ and define

$$[a]_q = \frac{q^a - 1}{q - 1}, \quad a \in \mathbb{R}.$$

The q analogue of the power $(a - b)^n$ is

$$(a - b)^{(0)} = 1, \quad (a - b)^{(n)} = \prod_{k=0}^{n-1} (a - bq^k), \quad a, b \in \mathbb{R}, n \in \mathbb{N}.$$

The q -gamma function is defined by

$$\Gamma_q(x) = \frac{(1 - q)^{(x-1)}}{(1 - q)^{x-1}}, \quad x \in \mathbb{R} \setminus \{0, -1, -2, \dots\}, 0 < q < 1$$

and satisfies $\Gamma_q(x + 1) = [x]_q \Gamma_q(x)$ (see, [38]).

For $0 < q < 1$, we define the q -derivative of a real valued function f as

$$D_q f(t) = \frac{f(t) - f(qt)}{(1 - q)t}, \quad t \in I_q - \{0\}, \quad D_q f(0) = \lim_{t \rightarrow 0} D_q f(t).$$

The higher order q -derivatives are given by

$$D_q^0 f(t) = f(t), \quad D_q^n f(t) = D_q D_q^{n-1} f(t), \quad n \in \mathbb{N}.$$

For $x \geq 0$, we set $J_x = \{xq^n : n \in \mathbb{N} \cup \{0\}\} \cup \{0\}$ and define the definite q -integral of a function $f : J_x \rightarrow \mathbb{R}$ by

$$I_q f(x) = \int_0^x f(s) d_q s = \sum_{n=0}^{\infty} x(1 - q)q^n f(xq^n)$$

provided that the series converges.

For $a, b \in J_x$, we set

$$\int_a^b f(s) d_q s = I_q f(b) - I_q f(a) = (1 - q) \sum_{n=0}^{\infty} q^n [bf(bq^n) - af(aq^n)],$$

provided that the series exist. Throughout the paper, we will assume that the series in the q -integrals converge.

Note that for $a, b \in J_x$, we have $a = xq^{n_1}$, $b = xq^{n_2}$ for some $n_1, n_2 \in \mathbb{N}$, thus the definite integral $\int_a^b f(s) d_q s$ is just a finite sum, so no question about convergence is raised.

We note that

$$D_q I_q f(x) = f(x),$$

while if f is continuous at $x = 0$, then

$$I_q D_q f(x) = f(x) - f(0).$$

For more details of the basic material on q -calculus, see the book [38].

Definition 2.1 ([31]) Let $\alpha \geq 0$ and f be a function defined on $[0, 1]$. The fractional q -integral of the Riemann-Liouville type is $(I_q^\alpha f)(t) = f(t)$ and

$$(I_q^\alpha f)(t) = \int_0^t \frac{(t - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} f(s) d_q(s), \quad \alpha > 0, t \in [0, 1].$$

Definition 2.2 ([39]) The fractional q -derivative of the Riemann-Liouville type of order $\alpha \geq 0$ is defined by $(D_q^\alpha f)(t) = f(t)$ and

$$(D_q^\alpha f)(t) = (D_q^{[\alpha]} I_q^{[\alpha]-\alpha} f)(t), \quad \alpha > 0,$$

where $[\alpha]$ is the smallest integer greater than or equal to α .

Definition 2.3 ([39]) The fractional q -derivative of the Caputo type of order $\alpha \geq 0$ is defined by

$$({}^c D_q^\alpha f)(t) = (I_q^{[\alpha]-\alpha} D_q^{[\alpha]} f)(t), \quad \alpha > 0,$$

where $[\alpha]$ is the smallest integer greater than or equal to α .

Lemma 2.4 Let $\alpha, \beta \geq 0$ and let f be a function defined on $[0, 1]$. Then the next formulas hold:

- (i) $(I_q^\beta I_q^\alpha f)(t) = (I_q^{\alpha+\beta} f)(t),$
- (ii) $(D_q^\alpha I_q^\alpha f)(t) = f(t).$

Lemma 2.5 ([33]) Let $\alpha \geq 0$ and $n \in \mathbb{N}$. Then the following equality holds:

$$(I_q^\alpha D_q^n f)(t) = D_q^n I_q^\alpha f(t) - \sum_{k=0}^{[\alpha]-1} \frac{t^{\alpha-n+k}}{\Gamma_q(\alpha - n + k)} (D_q^k f)(0).$$

Lemma 2.6 ([39]) *Let $\alpha > 0$. Then the following equality holds:*

$$(I_q^{\alpha c} D_q^\alpha f)(t) = f(t) - \sum_{k=0}^{[\alpha]-1} \frac{t^k}{\Gamma_q(k+1)} (D_q^k f)(0).$$

Lemma 2.7 ([40]) *For $\alpha \in \mathbb{R}^+$, $\lambda \in (-1, \infty)$, the following is valid:*

$$I_q^\alpha ((x-a)^{(\lambda)}) = \frac{\Gamma_q(\lambda+1)}{\Gamma(\alpha+\lambda+1)} (x-a)^{(\alpha+\lambda)}, \quad 0 < a < x < b.$$

In particular, for $\lambda = 0$, $a = 0$, using q -integration by parts, we have

$$\begin{aligned} (I_q^\alpha 1)(x) &= \frac{1}{\Gamma_q(\alpha)} \int_0^x (x-qt)^{(\alpha-1)} d_q t = \frac{1}{\Gamma_q(\alpha)} \int_0^x \frac{D_q((x-t)^{(\alpha)})}{-[\alpha]_q} d_q t \\ &= \frac{-1}{\Gamma_q(\alpha+1)} \int_0^x D_q((x-t)^{(\alpha)}) d_q t = \frac{1}{\Gamma_q(\alpha+1)} x^{(\alpha)}. \end{aligned}$$

In order to define the solution for the problem (1.1)-(1.2), we need the following lemma.

Lemma 2.8 *For a given $g \in C([0, 1], \mathbb{R})$ the unique solution of the boundary value problem*

$$\begin{cases} {}^c D_q^\alpha x(t) = g(t), & t \in [0, 1], \\ \alpha_1 x(0) - \beta_1 D_q x(0) = \gamma_1 x(\eta_1), \\ \alpha_2 x(1) + \beta_2 D_q x(1) = \gamma_2 x(\eta_2), \end{cases} \quad (2.1)$$

is given by

$$\begin{aligned} x(t) &= \int_0^t \frac{(t-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} g(s) d_q s \\ &+ \frac{\gamma_1}{\Delta} [(\alpha_2 - \gamma_2)t - (\alpha_2 + \beta_2 - \gamma_2 \eta_2)] \int_0^{\eta_1} \frac{(\eta_1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} g(s) d_q s \\ &- \frac{\gamma_2}{\Delta} [(\alpha_1 - \gamma_1)t + (\beta_1 + \gamma_1 \eta_1)] \int_0^{\eta_2} \frac{(\eta_2 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} g(s) d_q s \\ &+ \frac{\alpha_2}{\Delta} [(\alpha_1 - \gamma_1)t + \beta_1 + \gamma_1 \eta_1] \int_0^1 \frac{(1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} g(s) d_q s \\ &+ \frac{\beta_2}{\Delta} [(\alpha_1 - \gamma_1)t + \beta_1 + \gamma_1 \eta_1] \int_0^1 \frac{(1-qs)^{(\alpha-2)}}{\Gamma_q(\alpha-1)} g(s) d_q s, \end{aligned} \quad (2.2)$$

where

$$\Delta = (\gamma_2 - \alpha_2)(\beta_1 + \gamma_1 \eta_1) + (\gamma_1 - \alpha_1)(\alpha_2 + \beta_2 - \gamma_2 \eta_2) \neq 0. \quad (2.3)$$

Proof In view of Lemmas 2.4 and 2.6, integrating equation in (2.1), we have

$$x(t) = \int_0^t \frac{(t-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} g(s) d_q s + c_0 t + c_1, \quad t \in [0, 1]. \quad (2.4)$$

Using the boundary conditions of (2.1) in (2.4), we have

$$\begin{aligned}
 & -(\beta_1 + \gamma_1 \eta_1)c_0 + (\alpha_1 - \gamma_1)c_1 = \gamma_1 \int_0^{\eta_1} \frac{(\eta_1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} g(s) d_qs, \\
 & (\alpha_2 + \beta_2 - \gamma_2 \eta_2)c_0 + (\alpha_2 - \gamma_2)c_1 \\
 & = \gamma_2 \int_0^{\eta_2} \frac{(\eta_2 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} g(s) d_qs - \alpha_2 \int_0^1 \frac{(1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} g(s) d_qs \\
 & \quad - \beta_2 \int_0^1 \frac{(1 - qs)^{(\alpha-2)}}{\Gamma_q(\alpha - 1)} g(s) d_qs.
 \end{aligned}$$

Solving the above system of equations for c_0, c_1 , we get

$$\begin{aligned}
 c_0 & = \frac{1}{\Delta} \left\{ \gamma_1(\alpha_2 - \gamma_2) \int_0^{\eta_1} \frac{(\eta_1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} g(s) d_qs \right. \\
 & \quad - (\alpha_1 - \gamma_1)\gamma_2 \int_0^{\eta_2} \frac{(\eta_2 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} g(s) d_qs \\
 & \quad \left. + \alpha_2(\alpha_1 - \gamma_1) \int_0^1 \frac{(1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} g(s) d_qs + \beta_2(\alpha_1 - \gamma_1) \int_0^1 \frac{(1 - qs)^{(\alpha-2)}}{\Gamma_q(\alpha - 1)} g(s) d_qs \right\}, \\
 c_1 & = \frac{1}{\Delta} \left\{ -\gamma_1(\alpha_2 + \beta_2 - \gamma_2 \eta_2) \int_0^{\eta_1} \frac{(\eta_1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} g(s) d_qs \right. \\
 & \quad - \gamma_2(\beta_1 + \gamma_1 \eta_1) \int_0^{\eta_2} \frac{(\eta_2 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} g(s) d_qs \\
 & \quad + \beta_2(\beta_1 + \gamma_1 \eta_1) \int_0^1 \frac{(1 - qs)^{(\alpha-2)}}{\Gamma_q(\alpha - 1)} g(s) d_qs \\
 & \quad \left. + \alpha_2(\beta_1 + \gamma_1 \eta_1) \int_0^1 \frac{(1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} g(s) d_qs \right\}.
 \end{aligned}$$

Substituting the values of c_0, c_1 in (2.4), we obtain (2.2). □

In view of Lemma 2.8, we define an operator $F : C([0, 1], \mathbb{R}) \rightarrow C([0, 1], \mathbb{R})$ as

$$\begin{aligned}
 (Fx)(t) & = \int_0^t \frac{(t - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} f(s, x(s)) d_qs \\
 & \quad + \frac{\gamma_1}{\Delta} [(\alpha_2 - \gamma_2)t - (\alpha_2 + \beta_2 - \gamma_2 \eta_2)] \int_0^{\eta_1} \frac{(\eta_1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} f(s, x(s)) d_qs \\
 & \quad - \frac{\gamma_2}{\Delta} [(\alpha_1 - \gamma_1)t + (\beta_1 + \gamma_1 \eta_1)] \int_0^{\eta_2} \frac{(\eta_2 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} f(s, x(s)) d_qs \\
 & \quad + \frac{\alpha_2}{\Delta} [(\alpha_1 - \gamma_1)t + (\beta_1 + \gamma_1 \eta_1)] \int_0^1 \frac{(1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} f(s, x(s)) d_qs \\
 & \quad + \frac{\beta_2}{\Delta} [(\alpha_1 - \gamma_1)t + (\beta_1 + \gamma_1 \eta_1)] \int_0^1 \frac{(1 - qs)^{(\alpha-2)}}{\Gamma_q(\alpha - 1)} f(s, x(s)) d_qs. \tag{2.5}
 \end{aligned}$$

Observe that problem (1.1)-(1.2) has a solution if the operator equation $Fx = x$ has a fixed point, where F is given by (2.5).

3 Main results

Let $\mathcal{C} := C([0, 1], \mathbb{R})$ denote the Banach space of all continuous functions from $[0, 1] \rightarrow \mathbb{R}$ endowed with the norm defined by $\|x\| = \sup\{|x(t)| : t \in [0, 1]\}$.

For the sake of convenience, we set

$$k := (1 + |\alpha_2|\delta_2)(I_q^\alpha L)(1) + |\gamma_1|\delta_1(I_q^\alpha L)(\eta_1) + |\gamma_2|\delta_2(I_q^\alpha L)(\eta_2) + |\beta_2|\delta_2(I_q^{\alpha-1}L)(1), \quad (3.1)$$

where

$$\delta_1 := \frac{|\alpha_2 - \gamma_2| + |\alpha_2 + \beta_2 - \gamma_2\eta_2|}{|\Delta|}, \quad (3.2)$$

and

$$\delta_2 := \frac{|\alpha_1 - \gamma_1| + |\beta_1 + \gamma_1\eta_1|}{|\Delta|}. \quad (3.3)$$

Theorem 3.1 *Assume that $f : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous and that there exists a q -integrable function $L : [0, 1] \rightarrow \mathbb{R}$ such that*

$$(A_1) \quad |f(t, x) - f(t, y)| \leq L(t)|x - y|, \quad t \in [0, 1], \quad x, y \in \mathbb{R}.$$

Then the boundary value problem (1.1)-(1.2) has a unique solution provided

$$k < 1, \quad (3.4)$$

where k is given by (3.1).

Proof Let us fix $\sup_{t \in [0, 1]} |f(t, 0)| = M$ and choose

$$\rho \geq \frac{MA}{1 - k},$$

where

$$A = \frac{1}{\Gamma_q(\alpha + 1)}(1 + |\gamma_1|\delta_1\eta_1^{(\alpha-1)} + |\gamma_2|\delta_2\eta_2^{(\alpha-1)} + |\alpha_2|\delta_2) + \frac{|\beta_2|\delta_2}{\Gamma_q(\alpha)}. \quad (3.5)$$

We define $B_\rho = \{x \in \mathcal{C} : \|x\| \leq \rho\}$ and show that $FB_\rho \subset B_\rho$, where F is defined by (2.5). For $x \in B_\rho$, observe that

$$|f(t, x(t))| \leq |f(t, x(t)) - f(t, 0)| + |f(t, 0)| \leq L(t)|x(t)| + |f(t, 0)| \leq L(t)\rho + M.$$

Then $x \in B_\rho$, $t \in [0, 1]$, we have

$$\begin{aligned} |(Fx)(t)| &\leq \int_0^t \frac{(t - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} [L(s)\rho + M] d_qs \\ &\quad + |\gamma_1|\delta_1 \int_0^{\eta_1} \frac{(\eta_1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} [L(s)\rho + M] d_qs \\ &\quad + |\gamma_2|\delta_2 \int_0^{\eta_2} \frac{(\eta_2 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} [L(s)\rho + M] d_qs \end{aligned}$$

$$\begin{aligned}
 & + |\alpha_2| \delta_2 \int_0^1 \frac{(1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} [L(s)\rho + M] d_qs \\
 & + |\beta_2| \delta_2 \int_0^1 \frac{(1-qs)^{(\alpha-2)}}{\Gamma_q(\alpha-1)} [L(s)\rho + M] d_qs \\
 \leq & M \left\{ \int_0^1 \frac{(1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} d_qs + |\gamma_1| \delta_1 \int_0^{\eta_1} \frac{(\eta_1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} d_qs \right. \\
 & + |\gamma_2| \delta_2 \int_0^{\eta_2} \frac{(\eta_2-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} d_qs \\
 & + |\alpha_2| \delta_2 \int_0^1 \frac{(1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} d_qs + |\beta_2| \delta_2 \int_0^1 \frac{(1-qs)^{(\alpha-2)}}{\Gamma_q(\alpha-1)} d_qs \left. \right\} \\
 & + \rho \left\{ \int_0^1 \frac{(1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} L(s) d_qs + |\gamma_1| \delta_1 \int_0^{\eta_1} \frac{(\eta_1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} L(s) d_qs \right. \\
 & + |\gamma_2| \delta_2 \int_0^{\eta_2} \frac{(\eta_2-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} L(s) d_qs \\
 & + |\alpha_2| \delta_2 \int_0^1 \frac{(1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} L(s) d_qs + |\beta_2| \delta_2 \int_0^1 \frac{(1-qs)^{(\alpha-2)}}{\Gamma_q(\alpha-1)} L(s) d_qs \left. \right\} \\
 \leq & M \left\{ \frac{1}{\Gamma_q(\alpha+1)} (1 + |\gamma_1| \delta_1 \eta_1^{(\alpha-1)} + |\gamma_2| \delta_2 \eta_2^{(\alpha-1)} + |\alpha_2| \delta_2) + \frac{|\beta_2| \delta_2}{\Gamma_q(\alpha)} \right\} \\
 & + \rho \left\{ \int_0^1 \frac{(1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} L(s) d_qs + |\gamma_1| \delta_1 \int_0^{\eta_1} \frac{(\eta_1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} L(s) d_qs \right. \\
 & + |\gamma_2| \delta_2 \int_0^{\eta_2} \frac{(\eta_2-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} L(s) d_qs + |\alpha_2| \delta_2 \int_0^1 \frac{(1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} L(s) d_qs \\
 & \left. + |\beta_2| \delta_2 \int_0^1 \frac{(1-qs)^{(\alpha-2)}}{\Gamma_q(\alpha-1)} L(s) d_qs \right\},
 \end{aligned}$$

which, in view of (3.1) and (3.5), implies that

$$\|Fx\| \leq MA + \rho k \leq \rho.$$

This shows that $FB_\rho \subset B_\rho$.

Now, for $x, y \in \mathcal{C}$, we obtain

$$\begin{aligned}
 & \|(Fx) - (Fy)\| \\
 \leq & \sup_{t \in [0,1]} \left\{ \int_0^t \frac{(t-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} |f(s, x(s)) - f(s, y(s))| d_qs \right. \\
 & + |\gamma_1| \delta_1 \int_0^{\eta_1} \frac{(\eta_1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} |f(s, x(s)) - f(s, y(s))| d_qs \\
 & + |\gamma_2| \delta_2 \int_0^{\eta_2} \frac{(\eta_2-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} |f(s, x(s)) - f(s, y(s))| d_qs \\
 & + |\alpha_2| \delta_2 \int_0^1 \frac{(1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} |f(s, x(s)) - f(s, y(s))| d_qs \\
 & \left. + |\beta_2| \delta_2 \int_0^1 \frac{(1-qs)^{(\alpha-2)}}{\Gamma_q(\alpha-1)} |f(s, x(s)) - f(s, y(s))| d_qs \right\}
 \end{aligned}$$

$$\begin{aligned} &\leq \|x - y\| \sup_{t \in [0,1]} \left\{ \int_0^t \frac{(t - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} L(s) ds \right. \\ &\quad + |\gamma_1| \delta_1 \int_0^{\eta_1} \frac{(\eta_1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} L(s) d_qs + |\gamma_2| \delta_2 \int_0^{\eta_2} \frac{(\eta_2 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} L(s) d_qs \\ &\quad \left. + |\alpha_2| \delta_2 \int_0^1 \frac{(1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} L(s) d_qs + |\beta_2| \delta_2 \int_0^1 \frac{(1 - qs)^{(\alpha-2)}}{\Gamma_q(\alpha - 1)} L(s) d_qs \right\}, \end{aligned}$$

which, in view of (3.1), yields

$$\|(Fx)(t) - (Fy)(t)\| \leq k \|x - y\|.$$

Since $k \in (0, 1)$ by assumption (3.4), therefore, F is a contraction. Hence, it follows by Banach's contraction principle that the problem (1.1)-(1.2) has a unique solution. \square

In case $L(t) = L$ (L is a constant), the condition (3.4) becomes $LA < 1$ and Theorem 3.1 takes the form of the following result.

Corollary 3.2 *Assume that $f : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function and that*

there exists a constant $L \in (0, 1/A)$ with $|f(t, x) - f(t, y)| \leq L|x - y|$, $t \in [0, 1]$, $x, y \in \mathbb{R}$, where A is given by (3.5).

Then the boundary value problem (1.1)-(1.2) has a unique solution.

Our next existence results is based on Krasnoselskii's fixed-point theorem [41].

Lemma 3.3 (Krasnoselskii) *Let M be a closed, bounded, convex, and nonempty subset of a Banach space X . Let A, B be two operators such that:*

- (i) $Ax + By \in M$ whenever $x, y \in M$;
- (ii) A is compact and continuous;
- (iii) B is a contraction mapping.

Then there exists $z \in M$ such that $z = Az + Bz$.

Theorem 3.4 *Let $f : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function satisfying (A_1) . In addition, we assume that*

(A₂) there exists a function $\mu \in C([0, 1], \mathbb{R}^+)$ and a nondecreasing function $\phi \in C([0, 1], \mathbb{R}^+)$ with

$$|f(t, x)| \leq \mu(t)\phi(|x|), \quad (t, x) \in [0, 1] \times \mathbb{R};$$

(A₃) there exists a constant \bar{r} with

$$\begin{aligned} \bar{r} \geq &\left[\frac{1}{\Gamma_q(\alpha + 1)} + |\gamma_1| \delta_1 \frac{\eta_1^{(\alpha-1)}}{\Gamma_q(\alpha + 1)} + |\gamma_2| \delta_2 \frac{\eta_2^{(\alpha-1)}}{\Gamma_q(\alpha + 1)} \right. \\ &\left. + \frac{|\alpha_2| \delta_2}{\Gamma_q(\alpha + 1)} + \frac{|\beta_2| \delta_2}{\Gamma_q(\alpha)} \right] \phi(\bar{r}) \|\mu\|, \end{aligned} \tag{3.6}$$

where $\|\mu\| = \sup_{t \in [0, T]} |\mu(t)|$.

If

$$\begin{aligned}
 & |\gamma_1| \delta_1 \int_0^{\eta_1} \frac{(\eta_1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} L(s) d_qs + |\gamma_2| \delta_2 \int_0^{\eta_2} \frac{(\eta_2 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} L(s) d_qs \\
 & + |\alpha_2| \delta_2 \int_0^1 \frac{(1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} L(s) d_qs + |\beta_2| \delta_2 \int_0^1 \frac{(1 - qs)^{(\alpha-2)}}{\Gamma_q(\alpha - 1)} L(s) d_qs < 1, \tag{3.7}
 \end{aligned}$$

then the boundary value problem (1.1)-(1.2) has at least one solution on $[0, 1]$.

Proof Consider the set $B_{\bar{r}} = \{x \in \mathcal{C} : \|x\| \leq \bar{r}\}$, where \bar{r} is given in (3.6) and define the operators \mathcal{P} and \mathcal{Q} on $B_{\bar{r}}$ as

$$\begin{aligned}
 (\mathcal{P}x)(t) &= \int_0^t \frac{(t - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} f(s, x(s)) d_qs, \quad t \in [0, 1], \\
 (\mathcal{Q}x)(t) &= \frac{\gamma_1}{\Delta} [(\alpha_2 - \gamma_2)t - (\alpha_2 + \beta_2 - \gamma_2 \eta_2)] \int_0^{\eta_1} \frac{(\eta_1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} f(s, x(s)) d_qs \\
 &\quad - \frac{\gamma_2}{\Delta} [(\alpha_1 - \gamma_1)t + (\beta_1 + \gamma_1 \eta_1)] \int_0^{\eta_2} \frac{(\eta_2 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} f(s, x(s)) d_qs \\
 &\quad + \frac{\alpha_2}{\Delta} [(\alpha_1 - \gamma_1)t + \beta_1 + \gamma_1 \eta_1] \int_0^1 \frac{(1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} f(s, x(s)) d_qs \\
 &\quad + \frac{\beta_2}{\Delta} [(\alpha_1 - \gamma_1)t + \beta_1 + \gamma_1 \eta_1] \int_0^1 \frac{(1 - qs)^{(\alpha-2)}}{\Gamma_q(\alpha - 1)} f(s, x(s)) d_qs, \quad t \in [0, 1].
 \end{aligned}$$

For $x, y \in B_{\bar{r}}$, we find that

$$\begin{aligned}
 |(\mathcal{P}x + \mathcal{Q}y)(t)| &\leq \int_0^t \frac{(t - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} \mu(s) \phi(|x(s)|) d_qs \\
 &\quad + |\gamma_1| \delta_1 \int_0^{\eta_1} \frac{(\eta_1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} \mu(s) \phi(|x(s)|) d_qs \\
 &\quad + \gamma_2 \delta_2 \int_0^{\eta_2} \frac{(\eta_2 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} \mu(s) \phi(|x(s)|) d_qs \\
 &\quad + |\alpha_2| \delta_2 \int_0^1 \frac{(1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} \mu(s) \phi(|x(s)|) d_qs \\
 &\quad + |\beta_2| \delta_2 \int_0^1 \frac{(1 - qs)^{(\alpha-2)}}{\Gamma_q(\alpha - 1)} \mu(s) \phi(|x(s)|) d_qs \\
 &\leq \phi(\bar{r}) \|\mu\| \left\{ \frac{1}{\Gamma_q(\alpha + 1)} + |\gamma_1| \delta_1 \frac{\eta_1^{(\alpha-1)}}{\Gamma_q(\alpha + 1)} + |\gamma_2| \delta_2 \frac{\eta_2^{(\alpha-1)}}{\Gamma_q(\alpha + 1)} \right. \\
 &\quad \left. + |\alpha_2| \delta_2 \frac{1}{\Gamma_q(\alpha + 1)} + |\beta_2| \delta_2 \frac{1}{\Gamma_q(\alpha)} \right\} \\
 &\leq \bar{r}.
 \end{aligned}$$

Thus, $\mathcal{P}x + \mathcal{Q}y \in B_{\bar{r}}$. From (A₁) and (3.7) it follows that \mathcal{Q} is a contraction mapping. Continuity of f implies that the operator \mathcal{P} is continuous. Also, \mathcal{P} is uniformly bounded on $B_{\bar{r}}$

as

$$\|\mathcal{P}x\| \leq \frac{\phi(\bar{r})}{\Gamma_q(\alpha + 1)} \|\mu\|.$$

Now, for any $x \in B_{\bar{r}}$, and $t_1, t_2 \in [0, 1]$ with $t_1 < t_2$, we have

$$\begin{aligned} & |(\mathcal{P}x)(t_2) - (\mathcal{P}x)(t_1)| \\ &= \left| \int_0^{t_2} \frac{(t_2 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} f(s, x(s)) d_qs - \int_0^{t_1} \frac{(t_1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} f(s, x(s)) d_qs \right| \\ &\leq \left| \int_0^{t_1} \frac{(t_2 - qs)^{(\alpha-1)} - (t_1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} f(s, x(s)) d_qs \right| \\ &\quad + \left| \int_{t_1}^{t_2} \frac{(t_2 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} f(s, x(s)) d_qs \right| \\ &\leq \phi(\bar{r}) \|\mu\| \left[\int_0^{t_1} \frac{(t_2 - qs)^{(\alpha-1)} - (t_1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} d_qs + \int_{t_1}^{t_2} \frac{(t_2 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} d_qs \right] \end{aligned}$$

which is independent of x and tends to zero as $t_2 \rightarrow t_1$. Thus, \mathcal{P} is equicontinuous. So \mathcal{P} is relatively compact on $B_{\bar{r}}$. Hence, by the Arzelá-Ascoli theorem, \mathcal{P} is compact on $B_{\bar{r}}$. Thus, all the assumptions of Lemma 3.3 are satisfied. So the conclusion of Lemma 3.3 implies that the boundary value problem (1.1)-(1.2) has at least one solution on $[0, 1]$. \square

In the special case when $\phi(u) \equiv 1$, we see that there always exists a positive r so that (3.6) holds true, thus we have the following corollary.

Corollary 3.5 *Let $f : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function satisfying (A_1) . In addition, we assume that*

$$|f(t, x)| \leq \mu(t), \quad \forall (t, x) \in [0, 1] \times \mathbb{R}, \text{ and } \mu \in C([0, 1], \mathbb{R}^+).$$

If (3.7) holds, then the boundary value problem (1.1)-(1.2) has at least one solution on $[0, 1]$.

The next existence result is based on Leray-Schauder nonlinear alternative.

Lemma 3.6 (Nonlinear alternative for single valued maps [42]) *Let E be a Banach space, C a closed, convex subset of E , U an open subset of C with $0 \in U$. Suppose that $F : \bar{U} \rightarrow C$ is a continuous, compact (that is, $F(\bar{U})$ is a relatively compact subset of C) map. Then either*

- (i) *F has a fixed point in \bar{U} , or*
- (ii) *there is a $u \in \partial U$ (the boundary of U in C) and $\lambda \in (0, 1)$ with $u = \lambda F(u)$.*

Theorem 3.7 *Let $f : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function. Assume that:*

- (A₄) *there exist functions $p_1, p_2 \in L^1([0, 1], \mathbb{R}^+)$, and a nondecreasing function $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that $|f(t, x)| \leq p_1(t)\psi(|x|) + p_2(t)$, for $(t, x) \in [0, 1] \times \mathbb{R}$;*
- (A₅) *there exists a number $M > 0$ such that*

$$\frac{M}{\psi(M)\omega_1 + \omega_2} > 1, \tag{3.8}$$

where

$$\begin{aligned} \omega_i := & (1 + |\alpha_2|\delta_2)(I_q^\alpha p_i)(1) + |\gamma_1|\delta_1(I_q^\alpha p_i)(\eta_1) + |\gamma_2|\delta_2(I_q^\alpha p_i)(\eta_2) \\ & + |\beta_2|\delta_2(I_q^{\alpha-1} p_i)(1), \quad i = 1, 2. \end{aligned}$$

Then the boundary value problem (1.1)-(1.2) has at least one solution on $[0, 1]$.

Proof Consider the operator $F : \mathcal{C} \rightarrow \mathcal{C}$ defined by (2.5). It is easy to show that F is continuous. Next, we show that F maps bounded sets into bounded sets in $C([0, 1], \mathbb{R})$. For a positive number ρ , let $B_\rho = \{x \in C([0, 1], \mathbb{R}) : \|x\| \leq \rho\}$ be a bounded set in $C([0, 1], \mathbb{R})$. Then we have

$$\begin{aligned} |(Fx)(t)| & \leq \int_0^t \frac{(t - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} |f(s, x(s))| d_qs + |\gamma_1|\delta_1 \int_0^{\eta_1} \frac{(\eta_1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} |f(s, x(s))| d_qs \\ & \quad + |\gamma_2|\delta_2 \int_0^{\eta_2} \frac{(\eta_2 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} |f(s, x(s))| d_qs \\ & \quad + |\alpha_2|\delta_2 \int_0^1 \frac{(1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} |f(s, x(s))| d_qs + |\beta_2|\delta_2 \int_0^1 \frac{(1 - qs)^{(\alpha-2)}}{\Gamma_q(\alpha - 1)} |f(s, x(s))| d_qs \\ & \leq \int_0^t \frac{(t - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} [p_1(s)\psi(\|x\|) + p_2(s)] d_qs \\ & \quad + |\gamma_1|\delta_1 \int_0^{\eta_1} \frac{(\eta_1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} [p_1(s)\psi(\|x\|) + p_2(s)] d_qs \\ & \quad + |\gamma_2|\delta_2 \int_0^{\eta_2} \frac{(\eta_2 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} [p_1(s)\psi(\|x\|) + p_2(s)] d_qs \\ & \quad + |\alpha_2|\delta_2 \int_0^1 \frac{(1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} [p_1(s)\psi(\|x\|) + p_2(s)] d_qs \\ & \quad + |\beta_2|\delta_2 \int_0^1 \frac{(1 - qs)^{(\alpha-2)}}{\Gamma_q(\alpha - 1)} [p_1(s)\psi(\|x\|) + p_2(s)] d_qs \\ & \leq \psi(\rho) \left\{ \int_0^1 \frac{(1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} p_1(s) d_qs + |\gamma_1|\delta_1 \int_0^{\eta_1} \frac{(\eta_1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} p_1(s) d_qs \right. \\ & \quad + |\gamma_2|\delta_2 \int_0^{\eta_2} \frac{(\eta_2 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} p_1(s) d_qs + |\alpha_2|\delta_2 \int_0^1 \frac{(1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} p_1(s) d_qs \\ & \quad \left. + |\beta_2|\delta_1 \int_0^1 \frac{(1 - qs)^{(\alpha-2)}}{\Gamma_q(\alpha - 1)} p_1(s) d_qs \right\} \\ & \quad + \left\{ \int_0^1 \frac{(1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} p_2(s) d_qs + |\gamma_1|\delta_1 \int_0^{\eta_1} \frac{(\eta_1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} p_2(s) d_qs \right. \\ & \quad + |\gamma_2|\delta_2 \int_0^{\eta_2} \frac{(\eta_2 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} p_2(s) d_qs + |\alpha_2|\delta_2 \int_0^1 \frac{(1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} p_2(s) d_qs \\ & \quad \left. + |\beta_2|\delta_2 \int_0^1 \frac{(1 - qs)^{(\alpha-2)}}{\Gamma_q(\alpha - 1)} p_2(s) d_qs \right\}. \end{aligned}$$

Thus, for any $x \in B_\rho$ it holds

$$\|Fx\| \leq \psi(\rho)\omega_1 + \omega_2,$$

which proves our assertion.

Now we show that F maps bounded sets into equicontinuous sets of $C([0, 1], \mathbb{R})$. Let $t_1, t_2 \in [0, 1]$ with $t_1 < t_2$ and $x \in B_\rho$, where B_ρ is a bounded set of $C([0, 1], \mathbb{R})$. Then taking into consideration the inequality $(t_2 - qs)^{(\alpha-1)} - (t_1 - qs)^{(\alpha-1)} \leq (t_2 - t_1)$ for $0 < t_1 < t_2$ (see, [33, p.4]), we have

$$\begin{aligned} & |(Fx)(t_2) - (Fx)(t_1)| \\ & \leq \left| \int_0^{t_2} \frac{(t_2 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} f(s, x(s)) d_qs - \int_0^{t_1} \frac{(t_1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} f(s, x(s)) d_qs \right| \\ & \quad + \frac{1}{\Delta} \left\{ |\gamma_1(\alpha_2 - \gamma_2)|(t_2 - t_1) \int_0^{\eta_1} \frac{(\eta_1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} |f(s, x(s))| d_qs \right. \\ & \quad + \gamma_2|\alpha_1 - \gamma_1|(t_2 - t_1) \int_0^{\eta_2} \frac{(\eta_2 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} |f(s, x(s))| d_qs \\ & \quad + |\alpha_2||\alpha_1 - \gamma_1|(t_2 - t_1) \int_0^1 \frac{(1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} |f(s, x(s))| d_qs \\ & \quad \left. + |\beta_2||\alpha_1 - \gamma_1| \int_0^1 \frac{(1 - qs)^{(\alpha-2)}}{\Gamma_q(\alpha - 1)} |f(s, x(s))| d_qs \right\} \\ & \leq \left| \int_0^{t_1} (t_2 - t_1)[p_1(s)\psi(\rho) + p_2(s)] d_qs + \int_{t_1}^{t_2} (t_2 - qs)^{(\alpha-1)} [p_1(s)\psi(\rho) + p_2(s)] d_qs \right| \\ & \quad + \frac{1}{\Delta} \left\{ |\gamma_1(\alpha_2 - \gamma_2)|(t_2 - t_1) \int_0^{\eta_1} \frac{(\eta_1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} [p_1(s)\psi(\rho) + p_2(s)] d_qs \right. \\ & \quad + \gamma_2|\alpha_1 - \gamma_1|(t_2 - t_1) \int_0^{\eta_2} \frac{(\eta_2 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} [p_1(s)\psi(\rho) + p_2(s)] d_qs \\ & \quad + |\alpha_2||\alpha_1 - \gamma_1|(t_2 - t_1) \int_0^1 \frac{(1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} [p_1(s)\psi(\rho) + p_2(s)] d_qs \\ & \quad \left. + |\beta_2||\alpha_1 - \gamma_1|(t_2 - t_1) \int_0^1 \frac{(1 - qs)^{(\alpha-2)}}{\Gamma_q(\alpha - 1)} [p_1(s)\psi(\rho) + p_2(s)] d_qs \right\}. \end{aligned}$$

Obviously, the right-hand side of the above inequality tends to zero independently of $x \in B_\rho$ as $t_2 - t_1 \rightarrow 0$. Therefore, it follows by the Arzelà-Ascoli theorem that $F : C([0, 1], \mathbb{R}) \rightarrow C([0, 1], \mathbb{R})$ is completely continuous.

Thus, the operator F satisfies all the conditions of Lemma 3.6, and hence by its conclusion, either condition (i) or condition (ii) holds. We show that the conclusion (ii) is not possible.

Let $U = \{x \in C([0, 1], \mathbb{R}) : \|x\| < M\}$ with $\psi(M)\omega_1 + \omega_2 < M$ (by (3.8)). Then it can be shown that $\|Fx\| < M$. Indeed, in view of (A₄), we have

$$\begin{aligned} \|Fx\| & \leq \psi(\|x\|) \left\{ \int_0^1 \frac{(1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} p_1(s) d_qs + |\gamma_1|\delta_1 \int_0^{\eta_1} \frac{(\eta_1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} p_1(s) d_qs \right. \\ & \quad \left. + |\gamma_2|\delta_2 \int_0^{\eta_2} \frac{(\eta_2 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} p_1(s) d_qs + |\alpha_2|\delta_2 \int_0^1 \frac{(1 - qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} p_1(s) d_qs \right\} \end{aligned}$$

$$\begin{aligned}
 & + |\beta_2|\delta_2 \int_0^1 \frac{(1-qs)^{(\alpha-2)}}{\Gamma_q(\alpha-1)} p_1(s) d_qs \} \\
 & + \left\{ \int_0^1 \frac{(1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} p_2(s) d_qs + |\gamma_1|\delta_1 \int_0^{\eta_1} \frac{(\eta_1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} p_2(s) d_qs \right. \\
 & + |\gamma_2|\delta_2 \int_0^{\eta_2} \frac{(\eta_2-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} p_2(s) d_qs + |\alpha_2|\delta_2 \int_0^1 \frac{(1-qs)^{(\alpha-1)}}{\Gamma_q(\alpha)} p_2(s) d_qs \\
 & \left. + |\beta_2|\delta_2 \int_0^1 \frac{(1-qs)^{(\alpha-2)}}{\Gamma_q(\alpha-1)} p_2(s) d_qs \right\} \\
 & \leq \psi(M)\omega_1 + \omega_2 < M.
 \end{aligned}$$

Suppose there exists a $x \in \partial U$ and a $\lambda \in (0, 1)$ such that $x = \lambda Fx$. Then for such a choice of x and λ , we have

$$M = \|x\| = \lambda \|Fx\| < \psi(\|x\|)\omega_1 + \omega_2 = \psi(M)\omega_1 + \omega_2 < M,$$

which is a contradiction. Consequently, by the Leray-Schauder alternative (Lemma 3.6), we deduce that F has a fixed point $x \in \bar{U}$ which is a solution of the problem (1.1)-(1.2). This completes the proof. \square

Remark 3.8 If p_1, p_2 in (A_4) are continuous, then $\omega_i \leq A\|p_i\|, i = 1, 2$, where A is defined by (3.5).

Example 3.9 Consider the fractional q -difference nonlocal boundary value problem

$${}^c D_q^{3/2} x(t) = \frac{L}{2} (x + \tan^{-1} x + \sin t), \quad 0 \leq t \leq 1, \tag{3.9}$$

$$x(0) - 1/2 D_q x(0) = x(1/3), \quad \frac{1}{4} x(1) + \frac{3}{4} D_q x(1) = x(2/3). \tag{3.10}$$

Here, $\alpha_1 = 1, \beta_1 = 1/2, \alpha_2 = 1/4, \beta_2 = 3/4, \gamma_1 = 1 = \gamma_2, \eta_1 = 1/3, \eta_2 = 2/3$, and L is a constant to be fixed later on. Moreover, $\Delta = 5/8, \delta_1 = 26/15, \delta_2 = 4/3, |f(t, x) - f(t, y)| \leq L|x - y|$, and

$$k = \frac{L}{\Gamma_{1/2}(3/2)} \left(\frac{2\sqrt{2}(13 + 10\sqrt{2} + 15\sqrt{3})}{15\sqrt{3}(2\sqrt{2} - 1)} + 1 \right).$$

Choosing

$$L < \left[\frac{1}{\Gamma_{1/2}(3/2)} \left(\frac{2\sqrt{2}(13 + 10\sqrt{2} + 15\sqrt{3})}{15\sqrt{3}(2\sqrt{2} - 1)} + 1 \right) \right]^{-1},$$

all the assumptions of Theorem 3.1 are satisfied. Therefore, by Theorem 3.1, problem (3.9)-(3.10) has a unique solution.

Example 3.10 Consider the problem

$${}^c D_q^{3/2} x(t) = \frac{1}{4} \cos t^2 \sin(|x|/2) + \frac{e^{-x^2}(t^2 + 1)}{1 + (t^2 + 1)} + \frac{1}{3}, \quad 0 \leq t \leq 1, \tag{3.11}$$

$$x(0) - 1/2 D_q x(0) = x(1/3), \quad \frac{1}{4} x(1) + \frac{3}{4} D_q x(1) = x(2/3), \tag{3.12}$$

where $\alpha_1 = 1$, $\beta_1 = 1/2$, $\alpha_2 = 1/4$, $\beta_2 = 3/4$, $\gamma_1 = 1 = \gamma_2$, $\eta_1 = 1/3$, $\eta_2 = 2/3$. In a straightforward manner, it can be found that $\Delta = 5/8$, $\delta_1 = 26/15$, $f\delta_2 = 4/3$, and

$$|f(t, x)| = \left| \frac{1}{4} \cos t^2 \sin(|x|/2) + \frac{e^{-x^2}(t^2 + 1)}{1 + (t^2 + 1)} + \frac{1}{3} \right| \leq \frac{1}{8}|x| + 1.$$

Clearly, $p_1 = 1/8$, $p_2 = 1$, $\psi(M) = M$. Consequently, $\omega_1 = 0.567120414$, $\omega_2 = 4.536963312$, and the condition (3.8) implies that $M > 10.48055997$. Thus, all the assumptions of Theorem 3.7 are satisfied. Therefore, the conclusion of Theorem 3.7 applies to the problem (3.11)-(3.12).

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

Each of the authors, BA, SKN, and IKP contributed to each part of this study equally and read and approved the final version of the manuscript.

Author details

¹Department of Mathematics, Faculty of Science, King Abdulaziz University, P.O. Box 80203, Jeddah, 21589, Saudi Arabia.

²Department of Mathematics, University of Ioannina, Ioannina, 451 10, Greece.

Acknowledgements

The research of B. Ahmad was partially supported by Deanship of Scientific Research (DSR), King Abdulaziz University, Jeddah, Saudi Arabia. The authors thank the reviewers for their useful comments.

Received: 7 May 2012 Accepted: 2 August 2012 Published: 8 August 2012

References

1. Podlubny, I: Fractional Differential Equations. Academic Press, San Diego (1999)
2. Kilbas, AA, Srivastava, HM, Trujillo, JJ: Theory and Applications of Fractional Differential Equations. North-Holland Mathematics Studies, vol. 204. Elsevier, Amsterdam (2006)
3. Sabatier, J, Agrawal, OP, Machado, JAT (eds.): Advances in Fractional Calculus: Theoretical Developments and Applications in Physics and Engineering. Springer, Dordrecht (2007)
4. Baleanu, D, Diethelm, K, Scalas, E, Trujillo, JJ: Fractional Calculus Models and Numerical Methods. Series on Complexity, Nonlinearity and Chaos. World Scientific, Boston (2012)
5. Agarwal, RP, Belmekki, M, Benchohra, M: A survey on semilinear differential equations and inclusions involving Riemann-Liouville fractional derivative. *Adv. Differ. Equ.* **2009**, Article ID 981728 (2009)
6. Gafnychuk, V, Datsko, B, Meleshko, V: Mathematical modeling of different types of instabilities in time fractional reaction-diffusion systems. *Comput. Math. Appl.* **59**, 1101-1107 (2010)
7. Ahmad, B, Agarwal, RP: On nonlocal fractional boundary value problems. *Dyn. Contin. Discrete Impuls. Syst.* **18**, 535-544 (2011)
8. Baleanu, D, Mustafa, OG, O'Regan, D: A Nagumo-like uniqueness theorem for fractional differential equations. *J. Phys. A, Math. Theor.* **44**(39), Article ID 392003 (2011)
9. Wang, Y, Liu, L, Wu, Y: Positive solutions for a nonlocal fractional differential equation. *Nonlinear Anal.* **74**, 3599-3605 (2011)
10. Ford, NJ, Morgado, ML: Fractional boundary value problems: analysis and numerical methods. *Fract. Calc. Appl. Anal.* **14**, 554-567 (2011)
11. Ahmad, B, Nieto, JJ, Alsaedi, A, El-Shahed, M: A study of nonlinear Langevin equation involving two fractional orders in different intervals. *Nonlinear Anal., Real World Appl.* **13**, 599-606 (2012)
12. Ahmad, B, Ntouyas, SK: Nonlinear fractional differential equations and inclusions of arbitrary order and multi-strip boundary conditions. *Electron. J. Differ. Equ.* **2012**(98), 1-22 (2012)
13. Chen, F, Nieto, JJ, Zhou, Y: Global attractivity for nonlinear fractional differential equations. *Nonlinear Anal., Real World Appl.* **13**, 287-298 (2012)
14. Jackson, FH: On q -difference equations. *Am. J. Math.* **32**, 305-314 (1910)
15. Jackson, FH: On q -functions and a certain difference operator. *Trans. R. Soc. Edinb.* **46**, 253-281 (1908)
16. Ernst, T: A new notation for q -calculus and a new q -Taylor formula. U.U.D.M. Report 1999:25, Department of Mathematics, Uppsala University (1999). ISSN:1101-3591
17. Ernst, T: The history of q -calculus and a new method. U.U.D.M. Report 2000:16, Department of Mathematics, Uppsala University (2000). ISSN:1101-3591
18. Ahmad, B: Boundary-value problems for nonlinear third-order q -difference equations. *Electron. J. Differ. Equ.* **2011**(94), 1-7 (2011)
19. Ahmad, B, Ntouyas, SK: Boundary value problems for q -difference inclusions. *Abstr. Appl. Anal.* **2011**, Article ID 292860 (2011)

20. Ahmad, B, Alsaedi, A, Ntouyas, SK: A study of second-order q -difference equations with boundary conditions. *Adv. Differ. Equ.* **2012**, 35 (2012)
21. Atici, FM, Eloe, PW: A transform method in discrete fractional calculus. *Int. J. Differ. Equ.* **2**, 165-176 (2007)
22. Goodrich, CS: Solutions to a discrete right-focal fractional boundary value problem. *Int. J. Differ. Equ.* **5**, 195-216 (2010)
23. Goodrich, CS: Continuity of solutions to discrete fractional initial value problems. *Comput. Math. Appl.* **59**, 3489-3499 (2010)
24. Atici, FM, Eloe, PW: Linear systems of fractional nabla difference equation. *Rocky Mt. J. Math.* **41**, 353-360 (2011)
25. Atici, FM, Eloe, PW: Two-point boundary value problems for finite fractional difference equations. *J. Differ. Equ. Appl.* **17**, 445-456 (2011)
26. Bastos, NRO, Ferreira, RAC, Torres, DFM: Discrete-time fractional variational problems. *Signal Process.* **91**, 513-524 (2011)
27. Bastos, NRO, Mozyrska, D, Torres, DFM: Fractional derivatives and integrals on time scales via the inverse generalized Laplace transform. *Int. J. Math. Comput.* **11**, 1-9 (2011)
28. Goodrich, CS: Existence of a positive solution to a system of discrete fractional boundary value problems. *Appl. Math. Comput.* **217**, 4740-4753 (2011)
29. Goodrich, CS: On discrete sequential fractional boundary value problems. *J. Math. Anal. Appl.* **385**, 111-124 (2012)
30. Al-Salamb, WA: Some fractional q -integrals and q -derivatives. *Proc. Edinb. Math. Soc.* **15**, 135-140 (1966-1967)
31. Agarwal, R: Certain fractional q -integrals and q -derivatives. *Proc. Camb. Philos. Soc.* **66**, 365-370 (1969)
32. El-Shahed, M, Al-Askar, F: Positive solutions for boundary value problem of nonlinear fractional q -difference equation. *ISRN Math. Anal.* **2011**, Article ID 385459 (2011)
33. Ferreira, RAC: Positive solutions for a class of boundary value problems with fractional q -differences. *Comput. Math. Appl.* **61**, 367-373 (2011)
34. Ferreira, RAC: Nontrivial solutions for fractional q -difference boundary value problems. *Electron. J. Qual. Theory Differ. Equ.* **2010**(70), 1-10 (2010)
35. Graef, JR, Kong, L: Positive solutions for a class of higher order boundary value problems with fractional q -derivatives. *Appl. Math. Comput.* (2012). doi:10.1016/j.amc.2012.03.006
36. Ma, J, Yang, J: Existence of solutions for multi-point boundary value problem of fractional q -difference equation. *Electron. J. Qual. Theory Differ. Equ.* **2011**(92), 1-10 (2011)
37. Gasper, G, Rahman, M: *Basic Hypergeometric Series*. Cambridge University Press, Cambridge (1990)
38. Kac, V, Cheung, P: *Quantum Calculus*. Springer, New York (2002)
39. Rajkovic, PM, Marinkovic, SD, Stankovic, MS: On q -analogues of Caputo derivative and Mittag-Leffler function. *Fract. Calc. Appl. Anal.* **10**, 359-373 (2007)
40. Rajkovic, PM, Marinkovic, SD, Stankovic, MS: Fractional integrals and derivatives in q -calculus. *Appl. Anal. Discrete Math.* **1**, 311-323 (2007)
41. Smart, DR: *Fixed Point Theorems*. Cambridge University Press, Cambridge (1980)
42. Granas, A, Dugundji, J: *Fixed Point Theory*. Springer, New York (2005)

doi:10.1186/1687-1847-2012-140

Cite this article as: Ahmad et al.: Existence results for nonlocal boundary value problems of nonlinear fractional q -difference equations. *Advances in Difference Equations* 2012 2012:140.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
