

## Research Article

# Elementary Proof of Yu. V. Nesterenko Expansion of the Number Zeta(3) in Continued Fraction

**Leonid Gutnik**

*Moscow State Institute of Electronics and Mathematics, Russia*

Correspondence should be addressed to Leonid Gutnik, gutnik@gutnik.mccme.ru

Received 12 August 2009; Revised 10 December 2009; Accepted 10 January 2010

Academic Editor: Binggen Zhang

Copyright © 2010 Leonid Gutnik. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Yu. V. Nesterenko has proved that  $\zeta(3) = b_0 + a_1/|b_1 + \dots + a_\nu|/|b_\nu + \dots$ ,  $b_0 = b_1 = a_2 = 2$ ,  $a_1 = 1$ ,  $b_2 = 4$ ,  $b_{4k+1} = 2k + 2$ ,  $a_{4k+1} = k(k + 1)$ ,  $b_{4k+2} = 2k + 4$ , and  $a_{4k+2} = (k + 1)(k + 2)$  for  $k \in \mathbb{N}$ ;  $b_{4k+3} = 2k + 3$ ,  $a_{4k+3} = (k + 1)^2$ , and  $b_{4k+4} = 2k + 2$ ,  $a_{4k+4} = (k + 2)^2$  for  $k \in \mathbb{N}_0$ . His proof is based on some properties of hypergeometric functions. We give here an elementary direct proof of this result.

## 1. Foreword

Applications of difference equations to the Number Theory have a long history. For example, one can find in this journal several articles connected with the mentioned applications (see [1–8]). The interest in this area increases after Apéry's discovery of irrationality of the number  $\zeta(3)$ . This paper is inspired by Yu. V. Nesterenko's work [9]. My goal is to give an elementary direct proof of his expansion of the number  $\zeta(3)$  in continued fraction. Let us consider a difference equation

$$x_{\nu+1} - b_{\nu+1}x_\nu - a_{\nu+1}x_{\nu-1} = 0, \quad (1.1)$$

with  $\nu \in \mathbb{N}_0$ . We denote by

$$\{P_\nu(b_0, a_1, b_1, \dots, a_\nu, b_\nu)\}_{\nu=-1}^{+\infty}, \quad \{Q_\nu(b_0, a_1, b_1, \dots, a_\nu, b_\nu)\}_{\nu=-1}^{+\infty} \quad (1.2)$$

the solutions of this equation with initial values

$$P_{-1} = 1, \quad Q_{-1} = 0, \quad P_0(b_0) = b_0, \quad Q_0(b_0) = 1. \quad (1.3)$$

Then

$$\left\{ \frac{P_\nu(b_0, a_1, b_1, \dots, a_\nu, b_\nu)}{Q_\nu(b_0, a_1, b_1, \dots, a_\nu, b_\nu)} \right\}_{\nu=0}^{+\infty} \quad (1.4)$$

is a sequence of convergents of the continued fraction

$$b_0 + \frac{a_1}{|b_1|} + \dots + \frac{a_\nu}{|b_\nu|} + \dots \quad (1.5)$$

According to the famous result of R. Apéry [10],

$$\zeta(3) = \lim_{\nu \rightarrow \infty} \frac{v_\nu}{u_\nu}, \quad (1.6)$$

where  $\{u_\nu\}_{\nu=0}^{+\infty}$  and  $\{v_\nu\}_{\nu=0}^{+\infty}$  are solutions of difference equation

$$(\nu + 1)^3 x_{\nu+1} - (34\nu^3 + 51\nu^2 + 27\nu + 5)x_\nu + \nu^3 x_{\nu-1} = 0 \quad (1.7)$$

with initial values  $u_0 = 1$ ,  $u_1 = 5$ ,  $v_0 = 0$ ,  $v_1 = 6$ . The equality (1.6) is equivalent to the equality

$$\zeta(3) = b_0^\vee + \frac{a_1^\vee}{|b_1^\vee|} + \frac{a_2^\vee}{|b_2^\vee|} + \dots + \frac{a_\nu^\vee}{|b_\nu^\vee|} + \dots \quad (1.8)$$

with

$$b_0^\vee = 0, \quad b_1^\vee = 5, \quad a_1^\vee = 6, \quad b_{\nu+1}^\vee = 34\nu^3 + 51\nu^2 + 27\nu + 5, \quad a_{\nu+1}^\vee = -\nu^6, \quad (1.9)$$

where  $\nu \in \mathbb{N}$ . Nesterenko in [9] has offered the following expansion of the number  $2\zeta(3)$  in continued fraction:

$$2\zeta(3) = 2 + \frac{1}{|2|} + \frac{2}{|4|} + \frac{1}{|3|} + \frac{4}{|2|} \dots, \quad (1.10)$$

with

$$b_0 = b_1 = a_2 = 2, \quad a_1 = 1, \quad b_2 = 4, \quad (1.11)$$

$$b_{4k+1} = 2k + 2, \quad a_{4k+1} = k(k + 1), \quad b_{4k+2} = 2k + 4, \quad a_{4k+2} = (k + 1)(k + 2) \quad (1.12)$$

for  $k \in \mathbb{N}$ ;

$$b_{4k+3} = 2k + 3, \quad a_{4k+3} = (k + 1)^2, \quad b_{4k+4} = 2k + 2, \quad a_{4k+4} = (k + 2)^2 \quad (1.13)$$

for  $k \in \mathbb{N}_0$ .

The halved convergents of continued fraction (1.10) compose a sequence containing convergents of continued fraction (1.8). I give an elementary proof of Yu. V. Nesterenko expansion in Section 2.

## 2. Elementary Proof of Yu. V. Nesterenko Expansion

Instead of expansion (1.10) with (1.11), it is more convenient for us to prove the equivalent expansion

$$\zeta(3) = 1 + \frac{1|}{|4} + \frac{4|}{|4} + \frac{1|}{|3} + \frac{4|}{|2} \dots, \quad (2.1)$$

with

$$b_0 = 1, \quad a_1 = 1, \quad b_1 = a_2 = b_2 = 4. \quad (2.2)$$

Furthermore, to avoid confusion in notations, we denote below  $a_\nu, b_\nu$  for the fraction (2.1) by  $a_\nu^\wedge, b_\nu^\wedge$ . Let  $P_{-1}^\vee = 1, Q_{-1}^\vee = 0$ ,

$$P_\nu^\vee = P_\nu(b_0^\vee, a_1^\vee, b_1^\vee, \dots, a_\nu^\vee, b_\nu^\vee), \quad Q_\nu^\vee = Q_\nu(b_0^\vee, a_1^\vee, b_1^\vee, \dots, a_\nu^\vee, b_\nu^\vee), \quad (2.3)$$

where values  $a_\nu^\vee, b_\nu^\vee$  are specified in (1.9), and  $\nu \in \mathbb{N}_0$ . Then

$$\begin{aligned} Q_0^\vee = 1, \quad P_0^\vee = b_0^\vee = 0, \quad Q_1^\vee = b_1^\vee = 5, \quad P_1^\vee = a_1^\vee = 6, \quad b_2^\vee = 117, \quad a_2^\vee = -1, \\ P_2^\vee = b_2^\vee P_1^\vee + a_2^\vee P_0^\vee = 702, \quad Q_2^\vee = b_2^\vee Q_1^\vee + a_2^\vee Q_0^\vee = 584. \end{aligned} \quad (2.4)$$

Let  $P_{-1}^\wedge = 1, Q_{-1}^\wedge = 0$ ,

$$P_\nu^\wedge = P_\nu(b_0^\wedge, a_1^\wedge, b_1^\wedge, \dots, a_\nu^\wedge, b_\nu^\wedge), \quad Q_\nu^\wedge = Q_\nu(b_0^\wedge, a_1^\wedge, b_1^\wedge, \dots, a_\nu^\wedge, b_\nu^\wedge), \quad (2.5)$$

where  $\nu \in \mathbb{N}_0, a_\nu^\wedge := a_\nu, b_\nu^\wedge := b_\nu$ , and values  $a_\nu, b_\nu$  are specified in (2.2), (1.12), and (1.13). We calculate first  $P_k^\wedge$  and  $Q_k^\wedge$  for  $k = 0, \dots, 6$ .

Since  $P_{-1}^\wedge = 1$ ,  $Q_{-1}^\wedge = 0$ , it follows from (2.2) that

$$\begin{aligned} P_0^\wedge &= b_0 = 1, & Q_0^\wedge &= 1, \\ P_1^\wedge &= b_1^\wedge P_0^\wedge + a_1^\wedge P_{-1}^\wedge = 5, & Q_1^\wedge &= b_1^\wedge Q_0^\wedge + a_1^\wedge Q_{-1}^\wedge = 4, \end{aligned} \quad (2.6)$$

$$\begin{aligned} P_2^\wedge &= b_2^\wedge P_1^\wedge + a_2^\wedge P_0^\wedge = 24 = 4P_1^\vee, \\ Q_2^\wedge &= b_2^\wedge Q_1 + a_2^\wedge Q_0 = 20 = 4Q_1^\vee, \end{aligned} \quad (2.7)$$

$$\begin{aligned} P_3^\wedge &= b_3^\wedge P_2^\wedge + a_3^\wedge P_1^\wedge = 77, & Q_3^\wedge &= b_3^\wedge Q_2^\wedge + a_3^\wedge Q_1^\wedge = 64, \\ P_4^\wedge &= b_4^\wedge P_3^\wedge + a_4^\wedge P_2^\wedge = 250, & Q_4^\wedge &= b_4^\wedge Q_3^\wedge + a_4^\wedge Q_2^\wedge = 208, \end{aligned} \quad (2.8)$$

$$\begin{aligned} P_5^\wedge &= b_5^\wedge P_4^\wedge + a_5^\wedge P_3^\wedge = 1154, & Q_5^\wedge &= b_5^\wedge Q_4^\wedge + a_5^\wedge Q_3^\wedge = 960, \\ P_6^\wedge &= b_6^\wedge P_5^\wedge + a_6^\wedge P_4^\wedge = 12 \times 702 = 12P_2^\vee, \end{aligned} \quad (2.9)$$

$$Q_6^\wedge = b_6^\wedge Q_5^\wedge + a_6^\wedge Q_4^\wedge = 12 \times 584 = 12Q_2^\vee. \quad (2.10)$$

Let  $k \in \mathbb{N}$ ,  $k \geq 2$ ,

$$P_k^* = \frac{P_{4k-2}^\wedge}{2(k+1)!}, \quad Q_k^* = \frac{Q_{4k-2}^\wedge}{2(k+1)!}. \quad (2.11)$$

We want to prove that if  $k \in \mathbb{N}$ , then

$$P_k^* = P_k^\vee, \quad Q_k^* = Q_k^\vee. \quad (2.12)$$

Note that if  $k = 1, 2$ , then (2.12) follows from (2.6)–(2.10). Therefore, we can consider only  $k \in [3, +\infty) \cap \mathbb{Z}$ . Let us consider the following difference equations:

$$x_{\nu+1} - b_{\nu+1}^\vee x_\nu - a_{\nu+1}^\vee x_{\nu-1} = 0, \quad (2.13)$$

$$x_{\nu+1} - b_{\nu+1}^\wedge x_\nu - a_{\nu+1}^\wedge x_{\nu-1} = 0, \quad (2.14)$$

with  $\nu \in \mathbb{N}_0$ . Then  $x_\nu = P_\nu^\vee$ ,  $x_\nu = Q_\nu^\vee$ , with  $\nu \in (-1, +\infty) \cap \mathbb{Z}$  representing a fundamental system of solutions of (2.13), and  $x_\nu = P_\nu^\wedge$ ,  $x_\nu = Q_\nu^\wedge$  with  $\nu \in (-1, +\infty) \cap \mathbb{Z}$  representing a fundamental

system of solutions of (2.14). Making use of standard interpretation of a difference equation as a difference system, we rewrite the equalities (2.13) and (2.14), respectively in the form

$$X_{\nu+1} = A_{\nu}^{\vee} X_{\nu}, \tag{2.15}$$

$$X_{\nu+1} = A_{\nu}^{\wedge} X_{\nu}, \tag{2.16}$$

where

$$X_{\nu} = \begin{pmatrix} x_{\nu-1} \\ x_{\nu} \end{pmatrix}, \tag{2.17}$$

$$A_{\nu}^{\vee} = \begin{pmatrix} 0 & 1 \\ a_{1+\nu}^{\vee} & b_{1+\nu}^{\vee} \end{pmatrix}, \quad A_{\nu}^{\wedge} = \begin{pmatrix} 0 & 1 \\ a_{1+\nu}^{\wedge} & b_{1+\nu}^{\wedge} \end{pmatrix}, \tag{2.18}$$

and  $\nu \in \mathbb{N}_0$ . Let

$$U_{\nu}^{\vee} = \begin{pmatrix} P_{\nu-1}^{\vee} & Q_{\nu-1}^{\vee} \\ P_{\nu}^{\vee} & Q_{\nu}^{\vee} \end{pmatrix}, \tag{2.19}$$

$$U_{\nu}^{\wedge} = \begin{pmatrix} P_{\nu-1}^{\wedge} & Q_{\nu-1}^{\wedge} \\ P_{\nu}^{\wedge} & Q_{\nu}^{\wedge} \end{pmatrix}, \tag{2.20}$$

with  $\nu \in \mathbb{N}_0$  be fundamental matrices of solutions of systems (2.15) and (2.16), respectively. Therefore,

$$U_{\nu}^{\wedge} = A_{\nu-1}^{\wedge} U_{\nu-1}^{\wedge}, \quad U_{\nu}^{\vee} = A_{\nu-1}^{\vee} U_{\nu-1}^{\vee} \tag{2.21}$$

for  $\nu \in \mathbb{N}$ . In view of (2.18) and (2.21),  $\det(U_{\nu}) = -a_{\nu} \det(U_{\nu-1})$ , and therefore,

$$\det(U_{\nu}^{\wedge}) = (-1)^{\nu} \det(U_0^{\wedge}) \prod_{k=1}^{\nu} a_k^{\wedge} = (-1)^{\nu} \prod_{k=1}^{\nu} a_k^{\wedge}. \tag{2.22}$$

Hence

$$\frac{P_{\nu-1}^{\wedge}}{Q_{\nu-1}^{\wedge}} - \frac{P_{\nu}^{\wedge}}{Q_{\nu}^{\wedge}} = (-1)^{\nu} \frac{\prod_{k=1}^{\nu} a_k^{\wedge}}{Q_{\nu}^{\wedge} Q_{\nu-1}^{\wedge}} \tag{2.23}$$

(see [11]).

Further, we have

$$\begin{aligned} U_0^\vee &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, & U_1^\vee &= \begin{pmatrix} 0 & 1 \\ 6 & 5 \end{pmatrix}, & U_2^\vee &= \begin{pmatrix} 6 & 5 \\ 702 & 584 \end{pmatrix}, \\ U_0^\wedge &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, & U_1^\wedge &= \begin{pmatrix} 1 & 1 \\ 5 & 4 \end{pmatrix}, & U_2^\wedge &= \begin{pmatrix} 5 & 4 \\ 24 & 20 \end{pmatrix}, \end{aligned} \quad (2.24)$$

$$U_3^\wedge = \begin{pmatrix} 24 & 20 \\ 77 & 64 \end{pmatrix}, \quad U_4^\wedge = \begin{pmatrix} 77 & 64 \\ 250 & 208 \end{pmatrix},$$

$$U_5^\wedge = \begin{pmatrix} 250 & 208 \\ 1154 & 960 \end{pmatrix}, \quad U_6^\wedge = \begin{pmatrix} 1154 & 960 \\ 8424 & 7008 \end{pmatrix},$$

$$(U_1^\vee)(U_2^\wedge)^{-1} = \frac{1}{4} \begin{pmatrix} -24 & 5 \\ 0 & 1 \end{pmatrix}, \quad (2.25)$$

$$(U_2^\vee)(U_6^\wedge)^{-1} = \frac{1}{96} \begin{pmatrix} -36 & 5 \\ 0 & 8 \end{pmatrix}. \quad (2.26)$$

Let  $k \in \mathbb{N}, k \geq 2$ . Then, in view of (2.20),

$$\begin{aligned} A_{4k-6}^\wedge &= \begin{pmatrix} 0 & 1 \\ a_{4(k-2)+3}^\wedge & b_{4(k-2)+3}^\wedge \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ (k-1)^2 & 2k-1 \end{pmatrix}, \\ A_{4k-5}^\wedge &= \begin{pmatrix} 0 & 1 \\ a_{4(k-2)+4}^\wedge & b_{4(k-2)+4}^\wedge \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ k^2 & 2k-2 \end{pmatrix}, \\ A_{4k-4}^\wedge &= \begin{pmatrix} 0 & 1 \\ a_{4(k-1)+1}^\wedge & b_{4(k-1)+1}^\wedge \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ k^2 - k & 2k \end{pmatrix}, \\ A_{4k-3}^\wedge &= \begin{pmatrix} 0 & 1 \\ a_{4(k-1)+2}^\wedge & b_{4(k-1)+2}^\wedge \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ k^2 + k & 2k + 2 \end{pmatrix}. \end{aligned} \quad (2.27)$$

Let  $Y_k = X_{4k-6}$  for  $k \in [2, +\infty) \cap \mathbb{Z}$ . In view of (2.16) and (2.18),

$$Y_{k+1} = B_k^\wedge Y_k, \quad (2.28)$$

$$U_{4k-2}^\wedge = B_k^\wedge U_{4k-6}^\wedge, \quad (2.29)$$

where, as before,  $k \in [2, +\infty) \cap \mathbb{Z}$ ,

$$B_k^\wedge = A_{4k-3}^\wedge A_{4k-4}^\wedge A_{4k-5}^\wedge A_{4k-6}^\wedge = \begin{pmatrix} 5k(k-1)^3 & k(12k^2 - 15k + 5) \\ 12k(k+1)(k-1)^3 & k(k+1)(29k^2 - 36k + 12) \end{pmatrix}. \quad (2.30)$$

In view of (2.22), (2.2), (1.12), (1.13), (2.29), and (2.28), the matrix  $U_{4k-6}^\wedge$  is a fundamental matrix of solutions of system (2.28). The substitution  $Z_k = C_k Y_k$ , with  $\det(C_k) \neq 0$  for  $k \in [2, +\infty) \cap \mathbb{Z}$ , transforms the system (2.28) into the system

$$Z_{k+1} = D_k Z_k, \tag{2.31}$$

with  $D_k = C_{k+1} B_k^\wedge (C_k)^{-1}$  for  $k \in [2, +\infty) \cap \mathbb{Z}$ . We prove now that if we take  $k \in [3, +\infty) \cap \mathbb{Z}$ , and  $C_k = H_{k-1}$ , where

$$H_1 = \frac{1}{4} \begin{pmatrix} -24 & 5 \\ 0 & 1 \end{pmatrix}, \tag{2.32}$$

$$H_k = \begin{pmatrix} 12(k+2)(k+1)c(k+1) & -5(k+2)c(k+1) \\ 0 & -(k-1)^3 c(k) \end{pmatrix}, \tag{2.33}$$

with  $k \in [2, +\infty) \cap \mathbb{Z}$  and  $c(k) = (-2(k-1)^3(k+1)!)^{-1}$ , then we obtain the equality  $D_k = A_{k-1}^\vee$ . So, let  $k \in [3, +\infty) \cap \mathbb{Z}$ . Then, in view of (2.33),

$$H_{k-1} = \begin{pmatrix} 12(k+1)kc(k) & -5(k+1)c(k) \\ 0 & -(k-2)^3 c(k-1) \end{pmatrix}. \tag{2.34}$$

In view of (1.9)

$$b_k^\vee = 34(k-1)^3 + 51(k-1)^2 + 27(k-1) + 5 = 34k^3 - 51k^2 + 27k - 5, \quad a_k^\vee = -(k-1)^6, \tag{2.35}$$

where  $k \in [3, +\infty) \cap \mathbb{Z}$ . Hence, in view of (2.19),

$$A_{k-1}^\vee = \begin{pmatrix} 0 & 1 \\ -(k-1)^6 & 34k^3 - 51k^2 + 27k - 5 \end{pmatrix}. \tag{2.36}$$

In view of (2.34)–(2.36),

$$\begin{aligned} A_{k-1}^\vee H_{k-1} &= \begin{pmatrix} 0 & 1 \\ -(k-1)^6 & 34k^3 - 51k^2 + 27k - 5 \end{pmatrix} \times \begin{pmatrix} 12(k+1)kc(k) & -5(k+1)c(k) \\ 0 & -(k-2)^3 c(k-1) \end{pmatrix} \\ &= \begin{pmatrix} 0 & -(k-2)^3 c(k-1) \\ -(k-1)^6 12(k+1)kc(k) & (k-1)^6 5(k+1)c(k) - b_k^\vee (k-2)^6 c(k-1) \end{pmatrix}. \end{aligned} \tag{2.37}$$

In view of (2.30) and (2.33),

$$\begin{aligned}
 H_k B_k^\wedge &= \begin{pmatrix} 12(k+2)(k+1)c(k+1) & -5(k+2)c(k+1) \\ 0 & -(k-1)^3 c(k) \end{pmatrix} \\
 &\times \begin{pmatrix} 5k(k-1)^3 & k(12k^2 - 15k + 5) \\ 12k(k+1)(k-1)^3 & k(k+1)(29k^2 - 36k + 12) \end{pmatrix} \\
 &= \begin{pmatrix} 0 & (k+2)c(k+1)k(k+1)(-k^2) \\ -c(k)12k(k+1)(k-1)^6 & -(k-1)^3 c(k)k(k+1)(29k^2 - 36k + 12) \end{pmatrix}.
 \end{aligned} \tag{2.38}$$

Since

$$\begin{aligned}
 &-(k+2)(k+1)c(k+1)k^3 = -c(k-1)(k-2)^3, \\
 &-(k-1)^3 c(k)k(k+1)(29k^2 - 36k + 12) - (k-1)^6 5(k+1)c(k) \\
 &= -(34k^3 - 51k^2 + 27k - 5)(k-1)^3(k+1)c(k) \\
 &= -(34k^3 - 51k^2 + 27k - 5)(k-2)^3 c(k-1),
 \end{aligned} \tag{2.39}$$

it follows from (2.35), (2.37), and (2.38) that

$$A_{k-1}^\vee H_{k-1} = H_k B_k^\wedge \tag{2.40}$$

for  $k \in [3, +\infty) \cap \mathbb{Z}$ . We prove by induction now the following equality:

$$U_k^\vee = H_k U_{4k-2}^\wedge, \tag{2.41}$$

for any  $k \in \mathbb{N}$ . In view of (2.25) and (2.32), the equality (2.41) holds for  $k = 1$ . In view of (2.26) and (2.33), the equality (2.41) hold for  $k = 2$ . Let  $k \in [3, +\infty) \cap \mathbb{Z}$  and (2.41) holds for  $k - 1$ . Then, in view of (2.29), (2.40), and (2.21),

$$H_k U_{4k-2}^\wedge = H_k B_k U_{4k-6}^\wedge = A_{k-1}^\vee H_{k-1} U_{4k-6}^\wedge = A_{k-1}^\vee U_{k-1}^\vee = U_k^\vee. \tag{2.42}$$

So, the equality (2.41) holds for any  $k \in \mathbb{N}$ . In view of (2.41),

$$P_k^\vee = (2(k+1)!)^{-1} P_{4k-2}^\wedge, \quad Q_k^\vee = (2(k+1)!)^{-1} Q_{4k-2}^\wedge \tag{2.43}$$

for  $k \in [2, +\infty) \cap \mathbb{Z}$ . Since

$$P_\nu^\vee = (\nu!)^3 v_\nu, \quad Q_\nu^\vee = (\nu!)^3 u_\nu \tag{2.44}$$

for  $v$ , and  $u$ , in (1.6) and  $\nu \in \mathbb{N}_0$ , it follows from (2.43) and (2.44), that

$$P_{4k-2}^\wedge = 2(k+1)(k!)^4 v_k, \quad Q_{4k-2}^\wedge = 2(k+1)(k!)^4 u_k. \tag{2.45}$$

As it is well known, for any  $\varepsilon > 0$  there exist  $C_1(\varepsilon) > 0$  and  $C_2(\varepsilon) > 0$  such that

$$C_1(\varepsilon)(1 + \sqrt{2})^{4k(1-\varepsilon)} < |u_k| < C_2(\varepsilon)(1 + \sqrt{2})^{4k(1+\varepsilon)}, \tag{2.46}$$

$$C_1(\varepsilon)(1 + \sqrt{2})^{4k(1-\varepsilon)} < |v_k| < C_2(\varepsilon)(1 + \sqrt{2})^{4k(1+\varepsilon)}, \tag{2.47}$$

$$\frac{C_1(\varepsilon)}{(1 + \sqrt{2})^{8k(1+\varepsilon)}} < \left| \zeta(3) - \frac{v_k}{u_k} \right| < \frac{C_2(\varepsilon)}{(1 + \sqrt{2})^{8k(1-\varepsilon)}}. \tag{2.48}$$

We apply (2.23) now. Let  $k \in [2, +\infty) \cap \mathbb{Z}$ . In view of (2.2), (1.12)–(1.13), and (2.45), if  $\eta = 1, 2, 3$ , then

$$\begin{aligned} 0 &\leq \prod_{\kappa=1}^{4k-2+\eta} a_\kappa \leq \prod_{\kappa=1}^{4k+1} a_\kappa \leq a_{4k-1} a_{4k} a_{4k+1} \times k^3 (k+1)^3 \prod_{\kappa=1}^{4k-2} a_\kappa \\ &= 4k^3 (k+1)^3 \prod_{\kappa=2}^k a_{4\kappa-5} a_{4\kappa-4} a_{4\kappa-3} a_{4\kappa-2} \end{aligned} \tag{2.49}$$

$$\begin{aligned} &= 4k^3 (k+1)^3 \prod_{\kappa=2}^k (\kappa-1)^2 \kappa^2 (\kappa-1) \kappa \kappa (\kappa+1) = 2(k!)^8 (k+1)^4, \\ &4(k+1)^2 (k!)^8 u_k^2 = (Q_{4k-2})^2 < Q_{4k-3+\eta} Q_{4k-2+\eta}. \end{aligned} \tag{2.50}$$

In view of (2.23), (2.50), and (2.49), if  $\theta = 1, 2, 3$

$$\begin{aligned} \left| \frac{P_{4k-2}}{Q_{4k-2}} - \frac{P_{4k-2+\theta}}{Q_{4k-1+\theta}} \right| &\leq \sum_{\eta=1}^{\theta} \left| \frac{P_{4k-3+\eta}}{Q_{4k-3+\eta}} - \frac{P_{4k-2+\eta}}{Q_{4k-2+\eta}} \right| \\ &\leq \sum_{\eta=1}^3 \left| \frac{P_{4k-3+\eta}}{Q_{4k-3+\eta}} - \frac{P_{4k-2+\eta}}{Q_{4k-2+\eta}} \right| \leq 3 \frac{(k+1)^2}{2u_k^2} \leq (1 + \sqrt{2})^{8k(-1+o(1))}, \end{aligned} \tag{2.51}$$

when  $k \rightarrow +\infty$ . In view of (2.45), (2.48), and (2.51), there exist  $C_3(\varepsilon) > 0$  and  $C_4(\varepsilon) > 0$  such that

$$\frac{C_3(\varepsilon)}{(1 + \sqrt{2})^{8k(1+\varepsilon)}} < \left| \zeta(3) - \frac{P_{4k-2+\theta}^\wedge}{Q_{4k-2}^\wedge} \right| < \frac{C_4(\varepsilon)}{(1 + \sqrt{2})^{8k(1-\varepsilon)}}, \quad (2.52)$$

where  $\theta = 0, 1, 2, 3$ . So, the equality (2.1) is proved. In view of (2.23),

$$\zeta(3) - \frac{P_0^\wedge}{Q_0^\wedge} = \sum_{v=1}^{\infty} (-1)^{v-1} d_v, \quad (2.53)$$

where

$$0 < d_v = \frac{\prod_{k=1}^v a_k^\wedge}{(Q_v^\wedge Q_{v-1}^\wedge)}. \quad (2.54)$$

Further, we have

$$\frac{d_{v+1}}{d_v} = \frac{a_{v+1}^\wedge Q_{v-1}^\wedge}{b_{v+1}^\wedge Q_v^\wedge + a_{v+1}^\wedge Q_{v-1}^\wedge} < 1. \quad (2.55)$$

Hence, the series (2.53) is the series of Leibnitz type. Therefore,  $P_{2k-1}^\wedge / Q_{2k-1}^\wedge$  decreases, when  $k$  increases in  $\mathbb{N}$ , and  $P_{2k}^\wedge / Q_{2k}^\wedge$  increases, when  $k$  increases in  $\mathbb{N}$ .

## Acknowledgment

The author would like to express his thanks to the reviewer of this article for his efforts, his criticism, his advices, and indications of misprints. Ravi P. Agarwal had expressed a useful suggestion, which the author realized in foreword and references. He is grateful to him in this connection.

## References

- [1] T. Kim, K.-W. Hwang, and Y.-H. Kim, "Symmetry properties of higher-order Bernoulli polynomials," *Advances in Difference Equations*, vol. 2009, Article ID 318639, 6 pages, 2009.
- [2] T. Kim, K.-W. Hwang, and B. Lee, "A note on the  $q$ -Euler measures," *Advances in Difference Equations*, vol. 2009, Article ID 956910, 8 pages, 2009.
- [3] K.-H. Park and Y.-H. Kim, "On some arithmetical properties of the Genocchi numbers and polynomials," *Advances in Difference Equations*, vol. 2008, Article ID 195049, 14 pages, 2008.
- [4] Y. Simsek, I. N. Cangul, V. Kurt, V. Curt, and D. Kim, " $q$ -Genocchi numbers and polynomials associated with  $q$ -Genocchi-type  $l$ -functions," *Advances in Difference Equations*, vol. 2008, Article ID 815750, 12 pages, 2008.
- [5] L. C. Jang, "Multiple twisted  $q$ -Euler numbers and polynomials associated with  $p$ -adic  $q$ -integrals," *Advances in Difference Equations*, vol. 2008, Article ID 738603, 11 pages, 2008.

- [6] T. Kim, " $q$ -Bernoulli numbers associated with  $q$ -Stirling numbers," *Advances in Difference Equations*, vol. 2008, Article ID 743295, 10 pages, 2008.
- [7] M. Rachidi and O. Saeki, "Extending generalized Fibonacci sequences and their Binet-type formula," *Advances in Difference Equations*, vol. 2006, Article ID 23849, 11 pages, 2006.
- [8] J. H. Jaroma, "On the appearance of primes in linear recursive sequences," *Advances in Difference Equations*, vol. 2005, no. 2, pp. 145–151, 2005.
- [9] Yu. V. Nesterenko, "Some remarks on  $\zeta(3)$ ," *Rossiiskaya Akademiya Nauk. Matematicheskie Zametki*, vol. 59, no. 6, pp. 865–880, 1996.
- [10] R. Apéry, "Interpolation des fractions continues et irrationalité de certaines constantes," *Bulletin de la Section des Sciences du C.T.H.*, vol. 3, pp. 37–53, 1981.
- [11] O. Perron, *Die Lehre von den Kettenbrüchen. Bd I. Elementare Kettenbrüche*, B. G. Teubner Verlagsgesellschaft, Stuttgart, Germany, 1954.