

## Research Article

# A Study on the $p$ -Adic Integral Representation on $\mathbb{Z}_p$ Associated with Bernstein and Bernoulli Polynomials

Lee-Chae Jang,<sup>1</sup> Won-Joo Kim,<sup>2</sup> and Yilmaz Simsek<sup>3</sup>

<sup>1</sup> Department of Mathematics and Computer Science, Konkuk University, Chungju 138-701, Republic of Korea

<sup>2</sup> The Research Institute of Natural Sciences, Konkuk University, Seoul 138-701, Republic of Korea

<sup>3</sup> Department of Mathematics, Faculty of Arts and Science, University of Akdeniz, Antalya, Turkey

Correspondence should be addressed to Lee-Chae Jang, leechae.jang@kku.ac.kr

Received 13 August 2010; Accepted 15 September 2010

Academic Editor: Toka Diagana

Copyright © 2010 Lee-Chae Jang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We consider the Bernstein polynomials on  $\mathbb{Z}_p$  and investigate some interesting properties of Bernstein polynomials related to Stirling numbers and Bernoulli numbers.

## 1. Introduction

Let  $C[0, 1]$  denote the set of continuous function on  $[0, 1]$ . Then, Bernstein operator for  $f \in C[0, 1]$  is defined as

$$B_n(f)(x) = \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} x^k (1-x)^{n-k} = \sum_{k=0}^n f\left(\frac{k}{n}\right) B_{k,n}(x), \quad (1.1)$$

for  $k, n \in \mathbb{Z}$ , where  $B_{k,n}(x) = \binom{n}{k} x^k (1-x)^{n-k}$  is called Bernstein polynomial of degree  $n$ . Some researchers have studied the Bernstein polynomials in the area of approximation theory (see [1–6]).

Let  $p$  be a fixed prime number. Throughout this paper  $\mathbb{Z}_p$ ,  $\mathbb{Q}_p$ ,  $\mathbb{C}$ , and  $\mathbb{C}_p$  will, respectively, denote the ring of  $p$ -adic rational integers, the field of  $p$ -adic rational numbers, the complex number field, and the completion of algebraic closure of  $\mathbb{Q}_p$ . Let  $UD(\mathbb{Z}_p)$  be the

set of uniformly differentiable function on  $\mathbb{Z}_p$ . For  $f \in UD(\mathbb{Z}_p)$ , the  $p$ -adic  $q$ -integral on  $\mathbb{Z}_p$  is defined by

$$\int_{\mathbb{Z}_p} f(x) d\mu(x) = \lim_{N \rightarrow \infty} \frac{1}{p^N} \sum_{x=0}^{p^N-1} f(x) \quad (1.2)$$

(see [4, 7–15]).

In the special case, if we set  $f(x) = x^n$  in (1.2), we have

$$B_n = \int_{\mathbb{Z}_p} x^n d\mu(x). \quad (1.3)$$

In this paper, we consider Bernstein polynomials on  $\mathbb{Z}_p$  and we investigate some interesting properties of Bernstein polynomials related to Stirling numbers and Bernoulli numbers.

## 2. Bernstein Polynomials Related to Stirling Numbers and Bernoulli Numbers

In this section, for  $f \in UD(\mathbb{Z}_p)$ , we consider Bernstein type operator on  $\mathbb{Z}_p$  as follows:

$$B_n(f)(x) = \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} x^k (1-x)^{n-k} = \sum_{k=0}^n f\left(\frac{k}{n}\right) B_k(x), \quad (2.1)$$

for  $n \in \mathbb{Z}_+$ , where  $B_{k,n}(x) = \binom{n}{k} x^k (1-x)^{n-k}$  is called Bernstein polynomial of degree  $n$ . We consider Newton's forward difference operator as follows:

$$\begin{aligned} \Delta f(x) &= f(x+1) - f(x), \\ \Delta^n f(x) &= \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} f(x+k) = \sum_{k=0}^n \binom{n}{k} (-1)^k f(x+n-k). \end{aligned} \quad (2.2)$$

For  $x = 0$ ,

$$\Delta^n f(0) = \sum_{k=0}^n \binom{n}{k} (-1)^k f(n-k) = \sum_{n=0}^{\infty} \binom{n}{k} (-1)^{n-k} f(k). \quad (2.3)$$

Then, we have

$$f(n) = (1 + \Delta)^n f(0) = \sum_{l=0}^n \binom{n}{l} \Delta^l f(0). \quad (2.4)$$

From (2.4), we note that

$$f(x) = \sum_{n=0}^{\infty} \binom{x}{n} \Delta^n f(0), \tag{2.5}$$

where

$$\Delta^n f(0) = \sum_{k=0}^n \binom{n}{k} (-1)^k f(n-k). \tag{2.6}$$

The Stirling number of the first kind is defined by

$$\prod_{k=1}^n (1+kz) = \sum_{k=0}^n S_1(n,k) z^k, \tag{2.7}$$

and the Stirling number of the second kind is also defined by

$$\prod_{k=1}^n \left( \frac{1}{1+kz} \right) = \sum_{k=0}^n S_2(n,k) z^k. \tag{2.8}$$

By (2.5), (2.6), (2.7), and (2.8), we see that

$$S_2(n,k) = \frac{1}{k!} \sum_{j=0}^k (-1)^j \binom{k}{j} (k-j)^n, \tag{2.9}$$

where  $\Delta^n 0^m = \sum_{k=0}^n \binom{n}{k} (-1)^k (n-k)^m$ . Note that, for  $k \in \mathbb{Z}_+$  and  $x \in [0, 1]$ ,

$$\begin{aligned} F^{(k)}(t, x) &= \frac{t^k e^{(1-x)t} x^k}{k!} = x^k \sum_{n=0}^{\infty} \binom{n+k}{k} (1-x)^n \frac{t^{n+k}}{(n+k)!} \\ &= \sum_{n=k}^{\infty} \binom{n}{k} x^k (1-x)^{n-k} \frac{t^n}{(n)!} = \sum_{n=0}^{\infty} B_{k,n}(x) \frac{t^n}{n!}. \end{aligned} \tag{2.10}$$

Thus, we note that  $t^k e^{(1-x)t} x^k / k!$  is the generating function of Bernstein polynomial. It is easy to show that

$$\frac{1}{\binom{n}{k}} \int_{\mathbb{Z}_p} B_{k,n}(x) d\mu(x) = \sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l \int_{\mathbb{Z}_p} x^{l+k} d\mu(x) = \sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l B_{n+k}. \tag{2.11}$$

By (2.11), we obtain the following theorem.

**Theorem 2.1.** For  $n, k \in \mathbb{Z}_+$  with  $n \geq k$ , one has

$$\frac{1}{\binom{n}{k}} \int_{\mathbb{Z}_p} B_{k,n}(x) d\mu(x) = \sum_{m=0}^{\infty} \sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l B_{n+k}, \quad (2.12)$$

where  $B_n$  are the  $n$ th Bernoulli numbers.

In [12], it is known that

$$x^n = \sum_{k=0}^n \binom{x}{k} k! S_2(n, k), \quad (2.13)$$

$$\sum_{k=i-1}^n \frac{\binom{k}{i}}{\binom{n}{i}} B_{k,n}(x) = x^i, \quad (2.14)$$

for  $i \in \mathbb{N}$ . By (1.1) and (2.14), we see that

$$\begin{aligned} x^i &= \sum_{m=0}^{\infty} \binom{n-i+m-1}{m} (-1)^m x^{n-i-m} (1-x)^m \sum_{k=i-1}^n \frac{\binom{k}{i}}{\binom{n}{i}} B_{k,n}(x) \\ &= \sum_{m=0}^{\infty} \sum_{k=i-1}^n \frac{\binom{k}{i}}{\binom{n}{i}} \binom{n-i+m-1}{m} \binom{n}{k} (-1)^m x^{n-i-m+k} (1-x)^{n+m-k} \\ &= \sum_{m=0}^{\infty} \sum_{k=i}^n \sum_{l=0}^{n+m-k} \binom{n-i+m-1}{m} \binom{n+m-k}{l} \binom{n}{k} \\ &\quad \times (-1)^{l+m} x^{l+n-i-m+k}, \end{aligned} \quad (2.15)$$

for  $i \in \mathbb{N}$ . By (2.15), we obtain the following theorem.

**Theorem 2.2.** For  $n, k \in \mathbb{Z}_+$ , and  $i \in \mathbb{N}$ , one has

$$B_i = \sum_{m=0}^{\infty} \sum_{k=i}^n \sum_{l=0}^{m+n-k} \binom{n-i+m-1}{m} \binom{m+n-k}{l} \binom{n}{k} (-1)^{l+m} B_{l+n-i-m+k}. \quad (2.16)$$

From (2.13) and (2.14), we note that

$$\sum_{k=i-1}^n \frac{\binom{k}{i}}{\binom{n}{i}} B_{k,n}(x) = \sum_{k=0}^i \binom{x}{k} k! S_2(i, k). \quad (2.17)$$

In [16], it is known that

$$\int_{\mathbb{Z}_p} \binom{x}{n} d\mu(x) = \frac{1}{n+1}. \quad (2.18)$$

By (2.17), (2.18), and Theorem 2.2, we have

$$B_n = \sum_{k=0}^n \frac{k!}{k+1} (-1)^k S_2(k, n-k). \quad (2.19)$$

From the definition of the Stirling numbers of the first kind, we drive that

$$\binom{x}{n} n! = (x)_n = \sum_{k=0}^n S_1(n, k) x^k. \quad (2.20)$$

By (2.17), (2.19), and (2.20), we obtain the following theorem.

**Theorem 2.3.** For  $k, n \in \mathbb{Z}_+$  and  $i \in \mathbb{N}$ , one has

$$\sum_{k=i-1}^n \frac{\binom{k}{i}}{\binom{n}{i}} B_{k,n}(x) = \sum_{k=0}^i \sum_{l=0}^k S_1(n, l) S_2(i, k) x^l. \quad (2.21)$$

By Theorems 2.2 and 2.3, we obtain the following corollary.

**Corollary 2.4.** For  $k \in \mathbb{N}$ , one has

$$B_i(x) = \sum_{k=0}^i \sum_{l=0}^k S_1(n, l) S_2(i, k) B_l, \quad (2.22)$$

where  $B_i$  are the  $i$ th Bernoulli numbers.

## Acknowledgment

The present investigation was supported by the Scientific Research Project Administration of Akdeniz University.

## References

- [1] M. Acikgoz and S. Araci, "A study on the integral of the product of several type Bernstein polynomials," *IST Transaction of Applied Mathematics-Modelling and Simulation*. In press.
- [2] M. Acikgoz and S. Araci, "On the generating function of the Bernstein polynomials," in *Proceedings of the 8th International Conference of Numerical Analysis and Applied Mathematics (ICNAAM '10)*, AIP, Rhodes, Greece, March 2010.
- [3] S. Bernstein, "Demonstration du theoreme de Weierstrass, fondee sur le calcul des probabilities," *Communications of the Kharkov Mathematical Society*, vol. 13, pp. 1-2, 1913.
- [4] T. Kim, L. C. Jang, and H. Yi, "A note on the modified  $q$ -Bernstein polynomials," *Discrete Dynamics in Nature and Society*, vol. 2010, Article ID 706483, 12 pages, 2010.
- [5] G. M. Phillips, "Bernstein polynomials based on the  $q$ -integers," *Annals of Numerical Mathematics*, vol. 4, no. 1-4, pp. 511-518, 1997.
- [6] M. S. Kim, D. Kim, and T. Kim, "On the  $q$ -Euler numbers related to modified  $q$ -Bernstein polynomials," <http://arxiv.org/abs/1007.3317>.

- [7] T. Kim, " $q$ -Volkenborn integration," *Russian Journal of Mathematical Physics*, vol. 9, no. 3, pp. 288–299, 2002.
- [8] T. Kim, " $q$ -Bernoulli numbers and polynomials associated with Gaussian binomial coefficients," *Russian Journal of Mathematical Physics*, vol. 15, no. 1, pp. 51–57, 2008.
- [9] T. Kim, "Note on the Euler  $q$ -zeta functions," *Journal of Number Theory*, vol. 129, no. 7, pp. 1798–1804, 2009.
- [10] T. Kim, "On a  $q$ -analogue of the  $p$ -adic log gamma functions and related integrals," *Journal of Number Theory*, vol. 76, no. 2, pp. 320–329, 1999.
- [11] T. Kim, "Barnes-type multiple  $q$ -zeta functions and  $q$ -Euler polynomials," *Journal of Physics A*, vol. 43, no. 25, Article ID 255201, 2010.
- [12] T. Kim, "Power series and asymptotic series associated with the  $q$ -analog of the two-variable  $p$ -adic  $L$ -function," *Russian Journal of Mathematical Physics*, vol. 12, no. 2, pp. 186–196, 2005.
- [13] T. Kim, J. Choi, and Y.-H. Kim, "Some identities on the  $q$ -Bernstein polynomials,  $g$ -Stirling numbers and  $q$ -Bernoulli numbers," *Advanced Studies in Contemporary Mathematics*, vol. 20, no. 3, pp. 335–341, 2010.
- [14] T. Kim, "Some identities on the  $q$ -Euler polynomials of higher order and  $q$ -Stirling numbers by the fermionic  $p$ -adic integral on  $\mathbb{Z}_p$ ," *Russian Journal of Mathematical Physics*, vol. 16, no. 4, pp. 484–491, 2009.
- [15] T. Kim, "On  $p$ -adic interpolating function for  $q$ -Euler numbers and its derivatives," *Journal of Mathematical Analysis and Applications*, vol. 339, no. 1, pp. 598–608, 2008.
- [16] T. Kim, D.-W. Park, and S.-H. Rim, "On multivariate  $p$ -adic  $q$ -integrals," *Journal of Physics A*, vol. 34, no. 37, pp. 7633–7638, 2001.