

Research Article

Existence of Positive Solutions for Semipositone Higher-Order BVPS on Time Scales

Yuguo Lin and Minghe Pei

Department of Mathematics, Beihua University, JiLin City 132013, China

Correspondence should be addressed to Minghe Pei, peiminghe@ynu.ac.kr

Received 4 December 2009; Revised 16 March 2010; Accepted 29 March 2010

Academic Editor: Alberto Cabada

Copyright © 2010 Y. Lin and M. Pei. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We offer conditions on semipositone function $f(t, u_0, u_1, \dots, u_{n-2})$ such that the boundary value problem, $u^{\Delta^n}(t) + f(t, u(\sigma^{n-1}(t)), u^{\Delta}(\sigma^{n-2}(t)), \dots, u^{\Delta^{n-2}}(\sigma(t))) = 0$, $t \in (0, 1) \cap \mathbb{T}$, $n \geq 2$, $u^{\Delta^i}(0) = 0$, $i = 0, 1, \dots, n-3$, $\alpha u^{\Delta^{n-2}}(0) - \beta u^{\Delta^{n-1}}(0) = 0$, $\gamma u^{\Delta^{n-2}}(\sigma(1)) + \delta u^{\Delta^{n-1}}(\sigma(1)) = 0$, has at least one positive solution, where \mathbb{T} is a time scale and $f(t, u_0, u_1, \dots, u_{n-2}) \in C([0, 1] \times \mathbb{R}[0, \infty)^{n-1}, \mathbb{R}(-\infty, \infty))$ is continuous with $f(t, u_0, u_1, \dots, u_{n-2}) \geq -M$ for some positive constant M .

1. Introduction

Throughout this paper, let \mathbb{T} be a time scale, for any $a, b \in \mathbb{R} = (-\infty, +\infty)$ ($b > a$), the interval $[a, b]$ defined as $[a, b] = \{t \in \mathbb{T} \mid a \leq t \leq b\}$. Analogous notations for open and half-open intervals will also be used in the paper. We also use the notation $\mathbb{R}[c, d]$ to denote the real interval $\{t \in \mathbb{R} \mid c \leq t \leq d\}$. To understand further knowledge about dynamic equations on time scales, the reader may refer to [1–3] for an introduction to the subject.

In this paper, we present results governing the existence of positive solutions to the differential equation on time scales of the form

$$u^{\Delta^n}(t) + f\left(t, u\left(\sigma^{n-1}(t)\right), u^{\Delta}\left(\sigma^{n-2}(t)\right), \dots, u^{\Delta^{n-2}}\left(\sigma(t)\right)\right) = 0, \quad t \in (0, 1), \quad n \geq 2 \quad (1.1)$$

subject to the two-point boundary conditions

$$\begin{aligned} u^{\Delta^i}(0) &= 0, \quad i = 0, 1, \dots, n-3, \\ \alpha u^{\Delta^{n-2}}(0) - \beta u^{\Delta^{n-1}}(0) &= 0, \\ \gamma u^{\Delta^{n-2}}(\sigma(1)) + \delta u^{\Delta^{n-1}}(\sigma(1)) &= 0, \end{aligned} \quad (1.2)$$

where $\alpha, \gamma, \beta, \delta \geq 0, d := \alpha\delta + \gamma\beta + \alpha\gamma\sigma(1) > 0$ and

$$\delta + \gamma(\sigma(1) - \sigma^2(1)) \geq 0. \quad (1.3)$$

Throughout, we assume that $f(t, u_0, u_1, \dots, u_{n-2}) \in C([0, 1] \times \mathbb{R}[0, \infty)^{n-1}, \mathbb{R}(-\infty, \infty))$ is continuous with $f(t, u_0, u_1, \dots, u_{n-2}) \geq -M$ for some positive constant M .

Let $C_{rd}^n[0, 1]$ denote the space of functions

$$C_{rd}^n[0, 1] = \left\{ u : u \in C[0, \sigma^n(1)], \dots, u^{\Delta^{n-1}} \in C[0, \sigma(1)], u^{\Delta^n} \in C_{rd}[0, 1] \right\}. \quad (1.4)$$

We say that $u(t)$ is a positive solution of BVP (1.1) and (1.2), if $u(t) \in C_{rd}^n[0, 1]$ is a solution of BVP (1.1) and (1.2) and $u^{\Delta^i}(t) > 0, t \in (0, \sigma^{n-i}(1)), i = 0, 1, \dots, n-2$.

Various cases of BVP (1.1) and (1.2) have attracted a lot of attention in the literature. When $n = 2$, BVP (1.1) and (1.2) has been studied by many specialists. For example, Agarwal et al. [4] have established the existence of positive solutions for continuous case of the semipositone Sturm-Liouville BVPs. Erbe and Peterson [5] and Hao et al. [6] dealt with Sturm-Liouville BVPs on time scale of positone nonlinear term. In addition, Agarwal and O'Regan [7] obtained positive solution of second-order right focal BVPs on time scale by using nonlinear alternative of Leray-Schauder type. In 2005, Chyan and Wong [8] obtained triple solutions of the same BVPs with [7]. Recently, Sun and Li [9, 10] investigated semipositone Dirichlet BVPs on time scale. For higher-order BVPs, continuous case of BVP (1.1) and (1.2) have been investigated by Agarwal and Wong [11], Wong and Agarwal [12] and Wong [13]. The discrete positone case of BVP (1.1) and (1.2) has been tackled by using a fixed point theorem for mappings that are decreasing with respect to a cone in [14]. Especially, time-scale case of (1.1) with four-point boundary condition has been studied by Liu and Sang [15]. Besides, BVP (1.1) and (1.2) of nonlinear positone term $f(t, u(t), u'(t), \dots, u^{(n-1)}(t))$ which satisfied Nagumo-type conditions have been dealt with in [16]. Motivated by the works mentioned above, the purpose of this paper is to tackle semipositone BVP (1.1) and (1.2). In fact, BVPs appeared in [7–14] can be looked at as special case of BVP (1.1) and (1.2) in this paper. For other related works, we also refer to [17–19].

The paper is outlined as follows. In Section 2, we will present some notations and lemmas which will be used later. In Section 3, by using Krasnoselskii's fixed point theorem in a cone, we offer criteria for the existence of positive solution of BVP (1.1) and (1.2).

2. Preliminary

In this section, we offer some notations and lemmas, which will be used in main results. Throughout this paper, we always use the following notations:

(C₁) $K(t, s)$ is the Green's function of the differential equation $-u^{\Delta^n}(t) = 0, t \in (0, 1)$ subject to the boundary conditions (1.2);

(C₂) $k(t, s)$ is the Green's function of the differential equation $-u^{\Delta\Delta}(t) = 0, t \in (0, 1)$ subject to the boundary conditions

$$\alpha u(0) - \beta u^{\Delta}(0) = 0, \quad \gamma u(\sigma(1)) + \delta u^{\Delta}(\sigma(1)) = 0; \quad (2.1)$$

(C₃) Define $T_i : [0, 1] \rightarrow \mathbb{R}, i = 0, 1, \dots, n - 2$ as

$$T_0(t) \equiv q(t), \quad T_i(t) = \int_0^t T_{i-1}(\tau) \Delta \tau, \quad i = 1, 2, \dots, n - 2, \tag{2.2}$$

where

$$q(t) := \min_{t \in [0, \sigma^2(1)]} \left\{ \frac{\alpha t + \beta}{\alpha \sigma^2(1) + \beta} \frac{\gamma(\sigma(1) - t) + \delta}{\gamma \sigma(1) + \delta} \right\}. \tag{2.3}$$

Lemma 2.1. For the Green's function $k(t, s)$ the following hold:

$$0 \leq q(t)k(\sigma(s), s) \leq k(t, s) \leq k(\sigma(s), s), \quad (t, s) \in [0, \sigma^2(1)] \times [0, 1]. \tag{2.4}$$

Proof. It is clear that

$$k(t, s) = K^{\Delta_i^{n-2}}(t, s) = \begin{cases} \frac{1}{d} \{ \alpha t + \beta \} \{ \gamma(\sigma(1) - \sigma(s)) + \delta \}, & t \leq s, \\ \frac{1}{d} \{ \alpha \sigma(s) + \beta \} \{ \gamma(\sigma(1) - t) + \delta \}, & \sigma(s) \leq t. \end{cases} \tag{2.5}$$

From the expression of $k(t, s)$, we can easily obtain

$$0 \leq q(t)k(\sigma(s), s) \leq k(t, s) \leq k(\sigma(s), s), \quad (t, s) \in [0, \sigma^2(1)] \times [0, 1]. \tag{2.6}$$

□

Lemma 2.2. Let $w(t)$ be the solution of BVP

$$\begin{aligned} -u^{\Delta^n}(t) &= M, \quad t \in [0, 1], \\ u^{\Delta^i}(0) &= 0, \quad i = 0, 1, \dots, n - 3, \\ \alpha u^{\Delta^{n-2}}(0) - \beta u^{\Delta^{n-1}}(0) &= 0, \\ \gamma u^{\Delta^{n-2}}(\sigma(1)) + \delta u^{\Delta^{n-1}}(\sigma(1)) &= 0. \end{aligned} \tag{2.7}$$

Then

$$0 \leq w^{\Delta^i}(t) \leq cMT_{n-2-i}(t), \quad t \in [0, \sigma^{n-i}(1)], \quad i = 0, 1, \dots, n - 2, \tag{2.8}$$

where

$$c := \frac{(\gamma \sigma(1) + \delta)(\alpha \sigma^2(1) + \beta)}{d} \sigma(1), \tag{2.9}$$

and $M \in \mathbb{R}(0, \infty)$ is a positive constant.

Proof. For $t \leq s$,

$$\begin{aligned} k(t, s) &= \frac{1}{d} \{at + \beta\} \{\gamma(\sigma(1) - \sigma(s)) + \delta\} \leq \frac{1}{d} \{at + \beta\} \{\gamma(\sigma(1) - t) + \delta\} \\ &= \frac{at + \beta}{\alpha\sigma^2(1) + \beta} \cdot \frac{\gamma(\sigma(1) - t) + \delta}{\gamma\sigma(1) + \delta} \cdot \frac{(\gamma\sigma(1) + \delta)(\alpha^2\sigma(1) + \beta)}{d} \\ &\leq \frac{cq(t)}{\sigma(1)}. \end{aligned} \quad (2.10)$$

For $\sigma(s) \leq t$,

$$\begin{aligned} k(t, s) &= \frac{1}{d} \{\alpha\sigma(s) + \beta\} \{\gamma(\sigma(1) - t) + \delta\} \leq \frac{1}{d} \{at + \beta\} \{\gamma(\sigma(1) - t) + \delta\} \\ &= \frac{\gamma(\sigma(1) - t) + \delta}{\gamma\sigma(1) + \delta} \cdot \frac{at + \beta}{\alpha\sigma^2(1) + \beta} \cdot \frac{(\gamma\sigma(1) + \delta)(\alpha\sigma^2(1) + \beta)}{d} \\ &\leq \frac{cq(t)}{\sigma(1)}. \end{aligned} \quad (2.11)$$

So

$$0 \leq k(t, s) \leq \frac{cq(t)}{\sigma(1)}, \quad (t, s) \in [0, \sigma^2(1)] \times [0, 1]. \quad (2.12)$$

By defining $w(t)$ as $w(t) = \int_0^{\sigma(1)} K(t, s)M ds, t \in [0, \sigma^n(1)]$, it is clear that

$$w^{\Delta^{n-2}}(t) = \int_0^{\sigma(1)} k(t, s)M ds, \quad t \in [0, \sigma^2(1)]. \quad (2.13)$$

Then

$$0 \leq w^{\Delta^{n-2}}(t) \leq \frac{cq(t)}{\sigma(1)} \int_0^{\sigma(1)} M \Delta s = cMq(t), \quad t \in [0, \sigma^2(1)]. \quad (2.14)$$

Further, since $w^{\Delta^i}(t) = 0, i = 0, 1, \dots, n-3$, we get

$$0 \leq w^{\Delta^i}(t) \leq cMT_{n-2-i}(t), \quad t \in [0, \sigma^{n-i}(1)], \quad i = 0, 1, \dots, n-2. \quad (2.15)$$

□

Lemma 2.3 (see [20]). *Let E be a Banach space, and let $C \subset E$ be a cone in E . Assume that Ω_1, Ω_2 are open subsets of E with $0 \in \Omega_1 \subset \overline{\Omega}_1 \subset \Omega_2$, and let $T : C \cap (\overline{\Omega}_2 \setminus \Omega_1) \rightarrow C$ be a completely continuous operator such that either*

- (i) $\|Tu\| \leq \|u\|, u \in C \cap \partial\Omega_1; \|Tu\| \geq \|u\|, u \in C \cap \partial\Omega_2$ or
- (ii) $\|Tu\| \geq \|u\|, u \in C \cap \partial\Omega_1; \|Tu\| \leq \|u\|, u \in C \cap \partial\Omega_2$.

Then, T has a fixed point in $C \cap (\overline{\Omega_2} \setminus \Omega_1)$.

3. Main Results

In this section, by using Lemma 2.3, we offer criteria for the existence of positive solution of BVP (1.1) and (1.2).

Let E denote the space of functions

$$E = \left\{ u : u \in C[0, \sigma^n(1)], \dots, u^{\Delta^{n-3}} \in C[0, \sigma^3(1)], u^{\Delta^{n-2}} \in C[0, \sigma^2(1)] \right\}. \tag{3.1}$$

Let $\mathcal{B} = \{u \in E : u^{\Delta^i}(0) = 0, i = 0, 1, \dots, n - 3\}$ be a Banach space with the norm $\|u\| = \sup_{t \in [0, \sigma^2(1)]} |u^{\Delta^{n-2}}(t)|$, and let

$$C = \left\{ u \in \mathcal{B} : u^{\Delta^{n-2}}(t) \geq q(t)\|u\|, t \in [0, \sigma^2(1)] \right\}. \tag{3.2}$$

It is obvious that C is a cone in \mathcal{B} . From $u^{\Delta^i}(0) = 0, i = 0, 1, \dots, n - 3$, it follows that for all $u \in C$,

$$T_{n-2-i}(t)\|u\| \leq u^{\Delta^i}(t) \leq \sigma_0\|u\|, \quad t \in [0, \sigma^{n-i}(1)], \quad i = 0, 1, \dots, n - 2, \tag{3.3}$$

where

$$\sigma_0 := [\sigma^n(1)]^{n-2}. \tag{3.4}$$

Throughout the rest of the section, we assume that the set $[0, \sigma(1)]$ is such that

$$\xi = \min \left\{ \tau \in \mathbb{T} : \tau \geq \frac{\sigma(1)}{4} \right\}, \quad \zeta = \min \left\{ \tau \in \mathbb{T} : \tau \leq \frac{3\sigma(1)}{4} \right\} \tag{3.5}$$

exist and satisfy

$$\frac{\sigma(1)}{4} \leq \xi < \zeta \leq \frac{3\sigma(1)}{4}. \tag{3.6}$$

In addition, we denote that

$$\eta_i = \min_{t \in [\xi, \sigma^{n-i-1}(\zeta)]} T_{n-2-i}(t), \quad i = 0, 1, \dots, n - 2. \tag{3.7}$$

In order to obtain positive solutions of BVP (1.1) and (1.2), we need to consider the following boundary value problem:

$$\begin{aligned} u^{\Delta^n}(t) + f^*(t, v(\sigma^{n-1}(t)), v^{\Delta}(\sigma^{n-2}(t)), \dots, v^{\Delta^{n-2}}(\sigma(t))) &= 0, \quad t \in (0, 1), \\ u^{\Delta^i}(0) &= 0, \quad i = 0, 1, \dots, n-3, \\ \alpha u^{\Delta^{n-2}}(0) - \beta u^{\Delta^{n-1}}(0) &= 0, \quad \gamma u^{\Delta^{n-2}}(\sigma(1)) + \delta u^{\Delta^{n-1}}(\sigma(1)) = 0, \end{aligned} \quad (3.8)$$

where

$$\begin{aligned} v(t) &= u(t) - w(t), \quad (w(t) \text{ is as in Lemma 2.2}), \\ f^*(t, u_0, u_1, \dots, u_{n-2}) &= f(t, \rho_0, \rho_1, \dots, \rho_{n-2}) + M \end{aligned} \quad (3.9)$$

and for all $i = 0, 1, \dots, n-2$,

$$\rho_i = \begin{cases} u_i, & u_i \geq 0, \\ 0, & u_i < 0. \end{cases} \quad (3.10)$$

Let the operator $S : C \rightarrow \mathcal{B}$ be defined by

$$\begin{aligned} (Su)(t) &= \int_0^{\sigma(1)} K(t, s) f^*(s, v(\sigma^{n-1}(s)), v^{\Delta}(\sigma^{n-2}(s)), \dots, v^{\Delta^{n-2}}(\sigma(s))) \Delta s, \quad t \in [0, \sigma^n(1)], \\ (Su)^{\Delta^{n-2}}(t) &= \int_0^{\sigma(1)} k(t, s) f^*(s, v(\sigma^{n-1}(s)), v^{\Delta}(\sigma^{n-2}(s)), \dots, v^{\Delta^{n-2}}(\sigma(s))) \Delta s, \quad t \in [0, \sigma^2(1)]. \end{aligned} \quad (3.11)$$

Lemma 3.1. *The operator S maps C into C .*

Proof. From Lemma 2.1, we know that for $t \in [0, \sigma^2(1)]$,

$$\begin{aligned} (Su)^{\Delta^{n-2}}(t) &= \int_0^{\sigma(1)} k(t, s) f^*(s, v(\sigma^{n-1}(s)), v^{\Delta}(\sigma^{n-2}(s)), \dots, v^{\Delta^{n-2}}(\sigma(s))) \Delta s \\ &\leq \int_0^{\sigma(1)} k(\sigma(s), s) f^*(s, v(\sigma^{n-1}(s)), v^{\Delta}(\sigma^{n-2}(s)), \dots, v^{\Delta^{n-2}}(\sigma(s))) \Delta s. \end{aligned} \quad (3.12)$$

So for $t \in [0, \sigma^2(1)]$,

$$\|Su\| \leq \int_0^{\sigma(1)} k(\sigma(s), s) f^*(s, v(\sigma^{n-1}(s)), v^{\Delta}(\sigma^{n-2}(s)), \dots, v^{\Delta^{n-2}}(\sigma(s))) \Delta s. \quad (3.13)$$

From Lemma 2.1 again, it follows that for $t \in [0, \sigma^2(1)]$,

$$\begin{aligned} (Su)^{\Delta^{n-2}}(t) &= \int_0^{\sigma(1)} k(t,s) f^* \left(s, v \left(\sigma^{n-1}(s) \right), v^{\Delta} \left(\sigma^{n-2}(s) \right), \dots, v^{\Delta^{n-2}} \left(\sigma(s) \right) \right) \Delta s \\ &\geq \int_0^{\sigma(1)} q(t) k(\sigma(s), s) f^* \left(s, v \left(\sigma^{n-1}(s) \right), v^{\Delta} \left(\sigma^{n-2}(s) \right), \dots, v^{\Delta^{n-2}} \left(\sigma(s) \right) \right) \Delta s \\ &\geq q(t) \|Su\|. \end{aligned} \tag{3.14}$$

Hence, S maps C into C . □

Lemma 3.2. *The operator $S : C \rightarrow C$ is completely continuous.*

Proof. First we shall prove that the operator S is continuous. Let $u_m, u \in C$ be such that $\lim_{m \rightarrow \infty} \|u_m - u\| = 0$. From $u^{\Delta^i}(0) = 0, i = 0, 1, \dots, n - 3$, we have

$$\sup_{t \in [0, \sigma^{n-i}(1)]} |u_m^{\Delta^i}(t) - u^{\Delta^i}(t)| \rightarrow 0, \quad i = 0, 1, \dots, n - 2. \tag{3.15}$$

Then, it is easy to see that as $m \rightarrow \infty$

$$\begin{aligned} \rho_m := \sup_{s \in [0,1]} & \left| f^* \left(s, u_m \left(\sigma^{n-1}(s) \right) - w \left(\sigma^{n-1}(s) \right), \dots, u_m^{\Delta^{n-2}} \left(\sigma(s) \right) - w^{\Delta^{n-2}} \left(\sigma(s) \right) \right) \right. \\ & \left. - f^* \left(s, u \left(\sigma^{n-1}(s) \right) - w \left(\sigma^{n-1}(s) \right), \dots, u^{\Delta^{n-2}} \left(\sigma(s) \right) - w^{\Delta^{n-2}} \left(\sigma(s) \right) \right) \right| \rightarrow 0. \end{aligned} \tag{3.16}$$

Hence, we get from Lemma 2.1 that for $t \in [0, \sigma^2(1)]$,

$$\begin{aligned} & \left| (Su_m)^{\Delta^{n-2}}(t) - (Su)^{\Delta^{n-2}}(t) \right| \\ &= \left| \int_0^{\sigma(1)} k(t,s) \left[f^* \left(s, u_m \left(\sigma^{n-1}(s) \right) - w \left(\sigma^{n-1}(s) \right), \dots, u_m^{\Delta^{n-2}} \left(\sigma(s) \right) - w^{\Delta^{n-2}} \left(\sigma(s) \right) \right) \right. \right. \\ & \quad \left. \left. - f^* \left(s, u \left(\sigma^{n-1}(s) \right) - w \left(\sigma^{n-1}(s) \right), \dots, u^{\Delta^{n-2}} \left(\sigma(s) \right) - w^{\Delta^{n-2}} \left(\sigma(s) \right) \right) \right] \Delta s \right| \\ &\leq \rho_m \int_0^{\sigma(1)} k(t,s) \Delta s \leq \rho_m \int_0^{\sigma(1)} k(\sigma(s), s) \Delta s \rightarrow 0, \quad \text{as } m \rightarrow \infty. \end{aligned} \tag{3.17}$$

This shows that $S : C \rightarrow C$ is continuous.

Next, to show complete continuity, we will apply Arzela-Ascoli theorem. Let Ω be a bounded subset of C . Then there exists $L > 0$ such that for all $u \in \Omega$,

$$\sup |u^{\Delta^{n-2}}| \leq L, \quad \sup |u^{\Delta^i}| \leq \sigma_0 L, \quad i = 0, 1, \dots, n - 3, \tag{3.18}$$

where σ_0 is given in (3.4). Let

$$\widetilde{M} = \sup_{(s, \rho_0, \rho_1, \dots, \rho_{n-2}) \in [0,1] \times \mathbb{R} [0, \sigma_0 L]^{n-2} \times \mathbb{R} [0, L]} f(s, \rho_0, \rho_1, \dots, \rho_{n-2}) + M. \quad (3.19)$$

Clearly, we have for $t \in [0, \sigma^2(1)]$,

$$\left| (Su)^{\Delta_{n-2}}(t) \right| \leq \widetilde{M} \int_0^{\sigma(1)} k(t, s) \Delta s \leq \widetilde{M} \int_0^{\sigma(1)} k(\sigma(s), s) \Delta s \quad (3.20)$$

and for $t, t' \in [0, \sigma^2(1)]$,

$$\left| (Su)^{\Delta_{n-2}}(t) - (Su)^{\Delta_{n-2}}(t') \right| \leq \widetilde{M} \int_0^{\sigma(1)} |k(t, s) - k(t', s)| \Delta s. \quad (3.21)$$

The Arzela-Ascoli theorem guarantees that $S\Omega$ is relatively compact, so $S : C \rightarrow C$ is completely continuous. \square

Theorem 3.3. *Assume that the following conditions hold:*

- (i) *there exist $r \in \mathbb{R}(cM, \infty)$ such that for any $(u_0, u_1, \dots, u_{n-2}) \in \Gamma_r := \mathbb{R}[0, \sigma_0 r]^{n-2} \times \mathbb{R}[0, r]$,*

$$A(u_0, u_1, \dots, u_{n-2}) := \int_0^{\sigma(1)} k(\sigma(s), s) [f(s, u_0, u_1, \dots, u_{n-2}) + M] \Delta s \leq r, \quad (3.22)$$

- (ii) *there exist $R \in \mathbb{R}(cM, \infty)$ with $R \neq r$ such that for any $(u_0, u_1, \dots, u_{n-2}) \in \Gamma_R := \mathbb{R}[\epsilon \eta_0 R, \sigma_0 R] \times \mathbb{R}[\epsilon \eta_1 R, \sigma_0 R] \times \dots \times \mathbb{R}[\epsilon \eta_{n-2} R, R]$,*

$$B(u_0, u_1, \dots, u_{n-2}) := \eta_{n-2} \int_{\xi}^{\zeta} k(\sigma(s), s) [f(s, u_0, u_1, \dots, u_{n-2}) + M] \Delta s \geq R, \quad (3.23)$$

where σ_0 is given in (3.4), ξ, ζ are given in (3.5), $\eta_i, i = 0, 1, \dots, n-2$ are given in (3.7), and

$$\epsilon = 1 - \frac{cM}{R}. \quad (3.24)$$

Then BVP (1.1) and (1.2) has a positive solution.

Proof. Without loss of generality, we assume that $r < R$. Now we seek positive solutions of BVP (3.8). Let

$$\Omega_1 = \{u \in B : \|u\| \leq r\}. \quad (3.25)$$

For $u \in \partial\Omega_1 \cap C$, it follows from (3.3) that

$$0 \leq u^{\Delta^i}(t) \leq \sigma_0 r, \quad t \in [0, \sigma^{n-i}(1)], \quad i = 0, 1, \dots, n-3. \quad (3.26)$$

From (i), we obtain that for $u \in \partial\Omega_1 \cap C$,

$$\begin{aligned} (Su)^{\Delta^{n-2}}(t) &= \int_0^{\sigma(1)} k(t, s) f^* \left(s, v(\sigma^{n-1}(s)), v^{\Delta}(\sigma^{n-2}(s)), \dots, v^{\Delta^{n-2}}(\sigma(s))) \right) \Delta s \\ &\leq \int_0^{\sigma(1)} k(\sigma(s), s) f^* \left(s, v(\sigma^{n-1}(s)), v^{\Delta}(\sigma^{n-2}(s)), \dots, v^{\Delta^{n-2}}(\sigma(s))) \right) \Delta s \\ &\leq r. \end{aligned} \quad (3.27)$$

So

$$\|Su\| \leq \|u\|, \quad u \in \partial\Omega_1 \cap C. \quad (3.28)$$

Let

$$\Omega_2 = \{u \in B : \|u\| \leq R\}. \quad (3.29)$$

For $u \in \partial\Omega_2 \cap C$, it follows from Lemma 2.2 and (3.3) that for $s \in [0, \sigma(1)]$,

$$\begin{aligned} v^{\Delta^i}(\sigma^{n-i-1}(s)) &= u^{\Delta^i}(\sigma^{n-i-1}(s)) - w^{\Delta^i}(\sigma^{n-i-1}(s)) \\ &\geq u^{\Delta^i}(\sigma^{n-i-1}(s)) - cMT_{n-2-i}(\sigma^{n-i-1}(s)) \\ &= u^{\Delta^i}(\sigma^{n-i-1}(s)) - \frac{cMT_{n-2-i}(\sigma^{n-i-1}(s))\|u\|}{R} \\ &\geq u^{\Delta^i}(\sigma^{n-i-1}(s)) - \frac{cMu^{\Delta^i}(\sigma^{n-i-1}(s))}{R} \\ &= \epsilon u^{\Delta^i}(\sigma^{n-i-1}(s)) \geq \epsilon RT_{n-2-i}(\sigma^{n-i-1}(s)), \quad i = 0, 1, \dots, n-2. \end{aligned} \quad (3.30)$$

So

$$v^{\Delta^i}(\sigma^{n-i-1}(s)) \geq \epsilon \eta_i R, \quad s \in [\xi, \zeta], \quad i = 0, 1, \dots, n-2, \quad (3.31)$$

where η_i is given in (3.7) and ϵ is given in (3.24).

Combining Lemma 2.1, (3.3), and (ii) with (3.31), we obtain that for $u \in \partial\Omega_2 \cap C$,

$$\begin{aligned}
 (Su)^{\Delta^{n-2}}(t) &= \int_0^{\sigma(1)} k(t,s) f^*(s, v(\sigma^{n-1}(s)), v^\Delta(\sigma^{n-2}(s)), \dots, v^{\Delta^{n-2}}(\sigma(s))) \Delta s \\
 &\geq \int_\xi^\zeta k(t,s) f^*(s, v(\sigma^{n-1}(s)), v^\Delta(\sigma^{n-2}(s)), \dots, v^{\Delta^{n-2}}(\sigma(s))) \Delta s \\
 &\geq \eta_{n-2} \int_\xi^\zeta k(\sigma(s), s) f^*(s, v(\sigma^{n-1}(s)), v^\Delta(\sigma^{n-2}(s)), \dots, v^{\Delta^{n-2}}(\sigma(s))) \Delta s \\
 &\geq R.
 \end{aligned} \tag{3.32}$$

So

$$\|Su\| \geq \|u\|, \quad u \in \partial\Omega_2 \cap C. \tag{3.33}$$

Therefore, it follows from Lemma 2.3 that BVP (3.8) has a solution $u_1 \in C$ such that $r \leq \|u_1\| \leq R$.

Finally, we will prove that $u_1(t) - w(t)$ is a positive solution of BVP (1.1) and (1.2). Let $u(t) = u_1(t) - w(t)$, then we have from Lemma 2.2 and (3.3) that for $i = 0, 1, \dots, n-2$,

$$\begin{aligned}
 u^{\Delta^i}(t) &= u_1^{\Delta^i}(t) - w^{\Delta^i}(t) \geq u_1^{\Delta^i}(t) - cMT_{n-2-i}(t) \\
 &\geq u_1^{\Delta^i}(t) - \frac{cMT_{n-2-i}(t)\|u_1\|}{r} \\
 &\geq u_1^{\Delta^i}(t) - \frac{cMu_1^{\Delta^i}(t)}{r} \\
 &= \left(1 - \frac{cM}{r}\right)u_1^{\Delta^i}(t) \geq (r - cM)T_{n-2-i}(t) > 0, \quad t \in (0, \sigma^{n-i}(1)).
 \end{aligned} \tag{3.34}$$

In addition,

$$\begin{aligned}
 u^{\Delta^n}(t) &= u_1^{\Delta^n}(t) + M \\
 &= -f^*(t, u_1(\sigma^{n-1}(t)) - w(\sigma^{n-1}(t)), \dots, u_1^{\Delta^{n-2}}(\sigma(t)) - w^{\Delta^{n-2}}(\sigma(t))) + M \\
 &= -f(t, u_1(\sigma^{n-1}(t)) - w(\sigma^{n-1}(t)), \dots, u_1^{\Delta^{n-2}}(\sigma(t)) - w^{\Delta^{n-2}}(\sigma(t))) \\
 &= -f(t, u(\sigma^{n-1}(t)), \dots, u^{\Delta^{n-2}}(\sigma(t))).
 \end{aligned} \tag{3.35}$$

So, $u(t) = u_1(t) - w(t)$ is a positive solution of BVP (1.1) and (1.2). This completes the proof. \square

Corollary 3.4. *Assume that*

(a) *for any $(t, u_0, u_1, \dots, u_{n-2}) \in [0, 1] \times \mathbb{R}[0, \infty)^{n-1}$,*

$$f(t, u_0, u_1, \dots, u_{n-2}) + M \leq \mu(t)g(u_0, u_1, \dots, u_{n-2}), \tag{3.36}$$

where $g : \mathbb{R}[0, \infty)^{n-1} \rightarrow \mathbb{R}[0, \infty)$ is a continuous function which is nondecreasing in u_j for each fixed $(u_0, \dots, u_{j-1}, u_{j+1}, \dots, u_{n-2})$ and $\mu(t)$ is a continuous nonnegative function on $[0, 1]$,

(b) *for any $(t, u_0, u_1, \dots, u_{n-2}) \in [\xi, \zeta] \times \mathbb{R}[0, \infty)^{n-1}$,*

$$f(t, u_0, u_1, \dots, u_{n-2}) + M \geq \nu(t)h(u_0, u_1, \dots, u_{n-2}), \tag{3.37}$$

where $h : \mathbb{R}[0, \infty)^{n-1} \rightarrow \mathbb{R}[0, \infty)$ is a continuous function which is nondecreasing in u_j for each fixed $(u_0, \dots, u_{j-1}, u_{j+1}, \dots, u_{n-2})$ and $\nu(t)$ is a continuous nonnegative function on $[0, 1]$,

(c) *there exists $r \in \mathbb{R}(cM, \infty)$ such that*

$$g(\sigma_0 r, \dots, \sigma_0 r, r) \int_0^{\sigma(1)} k(\sigma(s), s) \mu(s) \Delta s \leq r, \tag{3.38}$$

(d) *there exists $R \in \mathbb{R}(cM, \infty)$ with $R \neq r$ such that*

$$h(\epsilon \eta_0 R, \epsilon \eta_1 R, \dots, \epsilon \eta_{n-2} R) \eta_{n-2} \int_{\xi}^{\zeta} k(\sigma(s), s) \nu(s) \Delta s \geq R. \tag{3.39}$$

Then BVP (1.1) and (1.2) has a positive solution.

Proof. From (a) and (c), we obtain that for $(u_0, u_1, \dots, u_{n-2}) \in \Gamma_r$,

$$\begin{aligned} A(u_0, u_1, \dots, u_{n-2}) &= \int_0^{\sigma(1)} k(\sigma(s), s) [f(s, u_0, u_1, \dots, u_{n-2}) + M] \Delta s \\ &\leq \int_0^{\sigma(1)} k(\sigma(s), s) \mu(s) g(u_0, u_1, \dots, u_{n-2}) \Delta s \\ &\leq g(\sigma_0 r, \dots, \sigma_0 r, r) \int_0^{\sigma(1)} k(\sigma(s), s) \mu(s) \Delta s \leq r. \end{aligned} \tag{3.40}$$

So, condition (i) of Theorem 3.3 is satisfied. From (b) and (d), we obtain that for $(u_0, u_1, \dots, u_{n-2}) \in \Gamma_R$,

$$\begin{aligned} B(u_0, u_1, \dots, u_{n-2}) &= \eta_{n-2} \int_{\xi}^{\zeta} k(\sigma(s), s) [f(s, u_0, u_1, \dots, u_{n-2}) + M] \Delta s \\ &\geq \eta_{n-2} \int_{\xi}^{\zeta} k(\sigma(s), s) v(s) h(u_0, u_1, \dots, u_{n-2}) \Delta s \\ &\geq h(\epsilon \eta_0 R, \epsilon \eta_1 R, \dots, \epsilon \eta_{n-2} R) \eta_{n-2} \int_{\xi}^{\zeta} k(\sigma(s), s) v(s) \Delta s \geq R. \end{aligned} \quad (3.41)$$

So, condition (ii) of Theorem 3.3 is satisfied.

Therefore, from Theorem 3.3, BVP(1.1) and (1.2) has a positive solution. \square

Corollary 3.5. *Assume that conditions (a) and (c) of Corollary 3.4 and the following condition hold:*

$$\lim_{u_0+u_1+\dots+u_{n-2} \rightarrow \infty} \min_{t \in [\xi, \zeta]} \frac{f(t, u_0, u_1, \dots, u_{n-2}) + M}{u_0 + u_1 + \dots + u_{n-2}} \in \mathbb{R} \left(\frac{D_1}{\epsilon \sum_{i=0}^{n-2} \eta_i}, \infty \right), \quad (3.42)$$

where $D_1 = [\eta_{n-2} \int_{\xi}^{\zeta} k(\sigma(s), s) \Delta s]^{-1}$. Then BVP (1.1) and (1.2) has one positive solution.

Proof. We only need to prove that (3.42) implies condition (ii) of Theorem 3.3. From (3.42), we know that there exists R (R may be chosen arbitrary large) such that for $(u_0, u_1, \dots, u_{n-2}) \in \mathbb{R}[\epsilon \eta_0 R, \infty) \times \dots \times \mathbb{R}[\epsilon \eta_{n-2} R, \infty)$,

$$\min_{t \in [\xi, \zeta]} \frac{f(t, u_0, u_1, \dots, u_{n-2}) + M}{u_0 + u_1 + \dots + u_{n-2}} \geq \frac{D_1}{\epsilon \sum_{i=0}^{n-2} \eta_i}. \quad (3.43)$$

Hence, for $(t, u_0, u_1, \dots, u_{n-2}) \in [\xi, \zeta] \times \Gamma_R$,

$$f(t, u_0, u_1, \dots, u_{n-2}) + M \geq \frac{D_1}{\epsilon \sum_{i=0}^{n-2} \eta_i} \sum_{i=0}^{n-2} u_i \geq \frac{D_1}{\epsilon \sum_{i=0}^{n-2} \eta_i} \cdot \epsilon \sum_{i=0}^{n-2} \eta_i R = D_1 R. \quad (3.44)$$

Thus, it follows that

$$\begin{aligned} B(u_0, u_1, \dots, u_{n-2}) &= \eta_{n-2} \int_{\xi}^{\zeta} k(\sigma(s), s) [f(s, u_0, u_1, \dots, u_{n-2}) + M] \Delta s \\ &\geq \eta_{n-2} \int_{\xi}^{\zeta} k(\sigma(s), s) D_1 R \Delta s = R. \end{aligned} \quad (3.45)$$

So, condition (ii) of Theorem 3.3 is satisfied. \square

Finally we present an example to illustrate our result.

Example 3.6. Consider the following boundary value problem:

$$\begin{aligned}
 u^{\Delta^3}(t) + \sin[u^{\Delta}(\sigma(t))] + \frac{\exp((u(\sigma^2(t)) + u^{\Delta}(\sigma(t)))/5)}{5(5 + u(\sigma^2(t)))} &= 0, \quad t \in (0, 1) \cap \mathbb{T}, \\
 u(0) = 0, \quad u^{\Delta}(0) - u^{\Delta^2}(0) = 0, \quad u^{\Delta^2}(\sigma(1)) = 0, &
 \end{aligned}
 \tag{3.46}$$

where $\mathbb{T} = 0 \cup \{t/8 : t \in \mathbb{N}\}$, $f(t, u_0, u_1) = \sin u_1 + \exp((u_0 + u_1)/5)/5(5 + u_0)$, $M = 1$, $\alpha = \beta = \delta = 1$, and $\gamma = 0$. Obviously,

$$d = \alpha\delta + \gamma\beta + \alpha\gamma\sigma(1) = 1, \quad \delta + \gamma(\sigma(1) - \sigma^2(1)) = 1 \geq 0, \quad \xi = \frac{3}{8}, \quad \zeta = \frac{6}{8}. \tag{3.47}$$

Let $\mu(t) = \nu(t) = 1$, $g(u_0, u_1) = 2 + \exp((u_0 + u_1)/5)/5(5 + u_0)$, and $h(u_0, u_1) = \exp((u_0 + u_1)/5)/5(5 + u_0)$. So conditions (a) and (b) in Corollary 3.4 are satisfied. By direct calculation, we obtain that $c = 81/32$, $\sigma_0 = 11/8$, $T_0(t) = q(t) = (4/9)(t + 1)$, $t \in [0, 10/8]$, and $T_1(t) = \int_0^t q(\tau)\Delta\tau = \sum_{\tau \in [0, t]} [\sigma(\tau) - \tau]q(\tau)$. Since $T_i(t)$, $i = 0, 1$ are nondecreasing, $\eta_0 = \min_{t \in [3/8, 1]} T_1(t) = T_1(3/8) = 3/16$, $\eta_1 = \min_{t \in [3/8, 7/8]} q(t) = q(3/8) = 11/18$. In addition, $\int_0^{\sigma(1)} k(\sigma(s), s)\Delta s = 117/64$, $\int_{\xi}^{\zeta} k(\sigma(s), s)\Delta s = 39/64$. Take $r = 5$, $R = 63$. So

$$\begin{aligned}
 &g(\sigma_0 r, r) \int_0^{\sigma(1)} k(\sigma(s), s)\mu(s)\Delta s \\
 &= \left(2 + \frac{\exp([\sigma_0 + 1]r/5)}{5(5 + \sigma_0 r)}\right) \int_0^{\sigma(1)} k(\sigma(s), s)\Delta s \approx 3.99 < 5 = r, \\
 &h(\epsilon\eta_0 R, \epsilon\eta_1 R)\eta_1 \int_{\xi}^{\zeta} k(\sigma(s), s)\nu(s)\Delta s \\
 &= \frac{\exp((R - cM)(\eta_0 + \eta_1)/5)}{5[5 + (R - cM)\eta_0]}\eta_1 \int_{\xi}^{\zeta} k(\sigma(s), s)\Delta s \approx 71.25 > 60 = R.
 \end{aligned}
 \tag{3.48}$$

Hence, conditions (c) and (d) in Corollary 3.4 are satisfied. Therefore from Corollary 3.4, (3.46) has at least one positive solution.

Remark 3.7. In Example 3.6, because nonlinear term f may attain negative value, the result in [15] is not applicable.

Acknowledgment

The authors thank the referee for valuable suggestions which led to improvement of the original manuscript.

References

- [1] R. P. Agarwal and M. Bohner, "Basic calculus on time scales and some of its applications," *Results in Mathematics. Resultate der Mathematik*, vol. 35, no. 1-2, pp. 3–22, 1999.
- [2] M. Bohner and A. Peterson, *Dynamic Equations on Time Scales: An Introduction with Application*, Birkhäuser, Boston, Mass, USA, 2001.
- [3] M. Bohner and A. Peterson, *Advances in Dynamic Equations on Time Scales*, Birkhäuser, Boston, Mass, USA, 2003.
- [4] R. P. Agarwal, H.-L. Hong, and C.-C. Yeh, "The existence of positive solutions for the Sturm-Liouville boundary value problems," *Computers & Mathematics with Applications*, vol. 35, no. 9, pp. 89–96, 1998.
- [5] L. Erbe and A. Peterson, "Positive solutions for a nonlinear differential equation on a measure chain," *Mathematical and Computer Modelling*, vol. 32, no. 5-6, pp. 571–585, 2000.
- [6] Z.-C. Hao, T.-J. Xiao, and J. Liang, "Existence of positive solutions for singular boundary value problem on time scales," *Journal of Mathematical Analysis and Applications*, vol. 325, no. 1, pp. 517–528, 2007.
- [7] R. P. Agarwal and D. O'Regan, "Nonlinear boundary value problems on time scales," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 44, no. 4, pp. 527–535, 2001.
- [8] C. J. Chyan and P. J. Y. Wong, "Triple solutions of focal boundary value problems on time scale," *Computers & Mathematics with Applications*, vol. 49, no. 7-8, pp. 963–979, 2005.
- [9] J.-P. Sun and W.-T. Li, "Existence of positive solutions to semipositone Dirichlet BVPs on time scales," *Dynamic Systems and Applications*, vol. 16, no. 3, pp. 571–578, 2007.
- [10] J.-P. Sun and W.-T. Li, "Solutions and positive solutions to semipositone Dirichlet BVPs on time scales," *Dynamic Systems and Applications*, vol. 17, no. 2, pp. 303–311, 2008.
- [11] R. P. Agarwal and F.-H. Wong, "Existence of positive solutions for non-positive higher-order BVPs," *Journal of Computational and Applied Mathematics*, vol. 88, no. 1, pp. 3–14, 1998.
- [12] P. J. Y. Wong and R. P. Agarwal, "On eigenvalue intervals and twin eigenfunctions of higher-order boundary value problems," *Journal of Computational and Applied Mathematics*, vol. 88, no. 1, pp. 15–43, 1998.
- [13] F.-H. Wong, "An application of Schauder's fixed point theorem with respect to higher order BVPs," *Proceedings of the American Mathematical Society*, vol. 126, no. 8, pp. 2389–2397, 1998.
- [14] P. J. Y. Wong and R. P. Agarwal, "On the existence of solutions of singular boundary value problems for higher order difference equations," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 28, no. 2, pp. 277–287, 1997.
- [15] J. Liu and Y. Sang, "Existence results for higher-order boundary value problems on time scales," *Advances in Difference Equations*, vol. 2009, Article ID 209707, 18 pages, 2009.
- [16] M. R. Grossinho and F. M. Minhós, "Upper and lower solutions for higher order boundary value problems," *Nonlinear Studies*, vol. 12, no. 2, pp. 165–176, 2005.
- [17] R. P. Agarwal, *Boundary Value Problems for Higher Order Differential Equations*, World Scientific, Singapore, 1986.
- [18] Y. Lin and M. Pei, "Positive solutions for two-point semipositone right focal eigenvalue problem," *Boundary Value Problems*, vol. 2007, Article ID 23108, 12 pages, 2007.
- [19] Y. Lin and M. Pei, "Positive solutions of two-point right focal eigenvalue problems on time scales," *Advances in Difference Equations*, vol. 2007, Article ID 87818, 15 pages, 2007.
- [20] D. Guo and V. Lakshmikantham, *Nonlinear Problems in Abstract Cones*, vol. 5 of *Notes and Reports in Mathematics in Science and Engineering*, Academic Press, Boston, Mass, USA, 1988.