

Research Article

Existence of Solutions for a Class of Damped Vibration Problems on Time Scales

Yongkun Li and Jianwen Zhou

Department of Mathematics, Yunnan University, Kunming, Yunnan 650091, China

Correspondence should be addressed to Yongkun Li, yklike@ynu.edu.cn

Received 3 June 2010; Revised 20 November 2010; Accepted 24 November 2010

Academic Editor: Kanishka Perera

Copyright © 2010 Y. Li and J. Zhou. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We present a recent approach via variational methods and critical point theory to obtain the existence of solutions for a class of damped vibration problems on time scale \mathbb{T} , $u^{\Delta^2}(t) + w(t)u^\Delta(\sigma(t)) = \nabla F(\sigma(t), u(\sigma(t)))$, Δ -a.e. $t \in [0, T]_{\mathbb{T}}^{\kappa}$, $u(0) - u(T) = 0$, $u^\Delta(0) - u^\Delta(T) = 0$, where $u^\Delta(t)$ denotes the delta (or Hilger) derivative of u at t , $u^{\Delta^2}(t) = (u^\Delta)^\Delta(t)$, σ is the forward jump operator, T is a positive constant, $w \in \mathcal{R}^+([0, T]_{\mathbb{T}}, \mathbb{R})$, $e_w(T, 0) = 1$, and $F : [0, T]_{\mathbb{T}} \times \mathbb{R}^N \rightarrow \mathbb{R}$. By establishing a proper variational setting, three existence results are obtained. Finally, three examples are presented to illustrate the feasibility and effectiveness of our results.

1. Introduction

Consider the damped vibration problem on time-scale \mathbb{T}

$$\begin{aligned} u^{\Delta^2}(t) + w(t)u^\Delta(\sigma(t)) &= \nabla F(\sigma(t), u(\sigma(t))), \quad \Delta\text{-a.e. } t \in [0, T]_{\mathbb{T}}^{\kappa}, \\ u(0) - u(T) &= 0, \quad u^\Delta(0) - u^\Delta(T) = 0, \end{aligned} \quad (1.1)$$

where $u^\Delta(t)$ denotes the delta (or Hilger) derivative of u at t , $u^{\Delta^2}(t) = (u^\Delta)^\Delta(t)$, σ is the forward jump operator, T is a positive constant, $w \in \mathcal{R}^+([0, T]_{\mathbb{T}}, \mathbb{R})$, $e_w(T, 0) = 1$, and $F : [0, T]_{\mathbb{T}} \times \mathbb{R}^N \rightarrow \mathbb{R}$ satisfies the following assumption.

- (A) $F(t, x)$ is Δ -measurable in t for every $x \in \mathbb{R}^N$ and continuously differentiable in x for $t \in [0, T]_{\mathbb{T}}$ and there exist $a \in C(\mathbb{R}^+, \mathbb{R}^+)$, $b \in L^1_{\Delta}([0, T]_{\mathbb{T}}, \mathbb{R}^+)$ such that

$$|F(t, x)| \leq a(|x|)b(t), \quad |\nabla F(t, x)| \leq a(|x|)b(t) \quad (1.2)$$

for all $x \in \mathbb{R}^N$ and $t \in [0, T]_{\mathbb{T}}$, where $\nabla F(t, x)$ denotes the gradient of $F(t, x)$ in x .

Problem (1.1) covers the second-order damped vibration problem (for when $\mathbb{T} = \mathbb{R}$)

$$\begin{aligned} \ddot{u}(t) + w(t)\dot{u}(t) &= \nabla F(t, u(t)), \quad \text{a.e. } t \in [0, T], \\ u(0) - u(T) &= 0, \quad \dot{u}(0) - \dot{u}(T) = 0, \end{aligned} \quad (1.3)$$

as well as the second-order discrete damped vibration problem (for when $\mathbb{T} = \mathbb{Z}, T \geq 2$)

$$\begin{aligned} \Delta^2(t) + w(t)\Delta u(t+1) &= \nabla F(t+1, u(t+1)), \quad t \in [0, T-1] \cap \mathbb{Z}, \\ u(0) - u(T) &= 0, \quad \Delta u(0) - \Delta u(T) = 0. \end{aligned} \quad (1.4)$$

The calculus of time-scales was initiated by Stefan Hilger in his Ph.D. thesis in 1988 in order to create a theory that can unify discrete and continuous analysis. A time-scale \mathbb{T} is an arbitrary nonempty closed subset of the real numbers, which has the topology inherited from the real numbers with the standard topology. The two most popular examples are $\mathbb{T} = \mathbb{R}$ and $\mathbb{T} = \mathbb{Z}$. The time-scales calculus has a tremendous potential for applications in some mathematical models of real processes and phenomena studied in physics, chemical technology, population dynamics, biotechnology and economics, neural networks, and social sciences (see [1]). For example, it can model insect populations that are continuous while in season (and may follow a difference scheme with variable step-size), die out in winter, while their eggs are incubating or dormant, and then hatch in a new season, giving rise to a nonoverlapping population.

In recent years, dynamic equations on time-scales have received much attention. We refer the reader to the books [2–7] and the papers [8–15]. In this century, some authors have begun to discuss the existence of solutions of boundary value problems on time-scales (see [16–22]). There have been many approaches to study the existence and the multiplicity of solutions for differential equations on time-scales, such as methods of lower and upper solutions, fixed-point theory, and coincidence degree theory. In [14], the authors have used the fixed-point theorem of strict-set-contraction to study the existence of positive periodic solutions for functional differential equations with impulse effects on time-scales. However, the study of the existence and the multiplicity of solutions for differential equations on time-scales using variational method has received considerably less attention (see, e.g., [19, 23]). Variational method is, to the best of our knowledge, novel and it may open a new approach to deal with nonlinear problems on time-scales.

When $w(t) \equiv 0$, (1.1) is the second-order Hamiltonian system on time-scale \mathbb{T}

$$\begin{aligned} u^{\Delta^2}(t) &= \nabla F(\sigma(t), u(\sigma(t))), \quad \Delta\text{-a.e. } t \in [0, T]_{\mathbb{T}}^{\kappa}, \\ u(0) - u(T) &= 0, \quad u^{\Delta}(0) - u^{\Delta}(T) = 0. \end{aligned} \quad (1.5)$$

Zhou and Li in [23] studied the existence of solutions for (1.5) by critical point theory on the Sobolev spaces on time-scales that they established.

When $w(t) \neq 0$, to the best of our knowledge, the existence of solutions for problems (1.1) have not been studied yet. Our purpose of this paper is to study the variational structure of problem (1.1) in an appropriate space of functions and the existence of solutions for problem (1.1) by some critical point theorems.

This paper is organized as follows. In Section 2, we present some fundamental definitions and results from the calculus on time-scales and Sobolev’s spaces on time-scales. In Section 3, we make a variational structure of (1.1). From this variational structure, we can reduce the problem of finding solutions of problem (1.1) to one of seeking the critical points of a corresponding functional. Section 4 is the existence of solutions. Section 5 is the conclusion of this paper.

2. Preliminaries and Statements

In this section, we present some basic definitions and results from the calculus on time-scales and Sobolev’s spaces on time-scales that will be required below. We first briefly recall some basic definitions and results concerning time-scales. Further general details can be found in [3–5, 7, 10, 13].

Through this paper, we assume that $0, T \in \mathbb{T}$. We start by the definitions of the forward and backward jump operators.

Definition 2.1 (see [3, Definition 1.1]). Let \mathbb{T} be a time-scale, for $t \in \mathbb{T}$, the forward jump operator $\sigma : \mathbb{T} \rightarrow \mathbb{T}$ is defined by

$$\sigma(t) = \inf\{s \in \mathbb{T}, s > t\}, \quad \forall t \in \mathbb{T}, \tag{2.1}$$

while the backward jump operator $\rho : \mathbb{T} \rightarrow \mathbb{T}$ is defined by

$$\rho(t) = \sup\{s \in \mathbb{T}, s < t\}, \quad \forall t \in \mathbb{T} \tag{2.2}$$

(supplemented by $\inf \emptyset = \sup \mathbb{T}$ and $\sup \emptyset = \inf \mathbb{T}$, where \emptyset denotes the empty set). A point $t \in \mathbb{T}$ is called right scattered, left scattered, if $\sigma(t) > t, \rho(t) < t$ hold, respectively. Points that are right scattered and left scattered at the same time are called isolated. Also, if $t < \sup \mathbb{T}$ and $\sigma(t) = t$, then t is called right-dense, and if $t > \inf \mathbb{T}$ and $\rho(t) = t$, then t is called left dense. Points that are right-dense and left dense at the same time are called dense. The set \mathbb{T}^κ which is derived from the time-scale \mathbb{T} as follows: if \mathbb{T} has a left scattered maximum m , the $\mathbb{T}^\kappa = \mathbb{T} - \{m\}$, otherwise, $\mathbb{T}^\kappa = \mathbb{T}$. Finally, the graininess function $\mu : \mathbb{T} \rightarrow [0, \infty)$ is defined by

$$\mu(t) = \sigma(t) - t. \tag{2.3}$$

When $a, b \in \mathbb{T}, a < b$, we denote the intervals $[a, b]_{\mathbb{T}}, [a, b)_{\mathbb{T}}$, and $(a, b]_{\mathbb{T}}$ in \mathbb{T} by

$$[a, b]_{\mathbb{T}} = [a, b] \cap \mathbb{T}, \quad [a, b)_{\mathbb{T}} = [a, b) \cap \mathbb{T}, \quad (a, b]_{\mathbb{T}} = (a, b] \cap \mathbb{T}, \tag{2.4}$$

respectively. Note that $[a, b]_{\mathbb{T}}^\kappa = [a, b]_{\mathbb{T}}$ if b is left dense and $[a, b]_{\mathbb{T}}^\kappa = [a, b)_{\mathbb{T}} = [a, \rho(b)]_{\mathbb{T}}$ if b is left scattered. We denote $[a, b]_{\mathbb{T}}^{\kappa^2} = ([a, b]_{\mathbb{T}}^\kappa)^\kappa$, therefore $[a, b]_{\mathbb{T}}^{\kappa^2} = [a, b]_{\mathbb{T}}$ if b is left dense and $[a, b]_{\mathbb{T}}^{\kappa^2} = [a, \rho(b)]_{\mathbb{T}}^\kappa$ if b is left scattered.

Definition 2.2 (see [3, Definition 1.10]). Assume that $f : \mathbb{T} \rightarrow \mathbb{R}$ is a function and let $t \in \mathbb{T}^\kappa$. Then we define $f^\Delta(t)$ to be the number (provided it exists) with the property that given any $\epsilon > 0$, there is a neighborhood U of t (i.e., $U = (t - \delta, t + \delta) \cap \mathbb{T}$ for some $\delta > 0$) such that

$$\left| [f(\sigma(t)) - f(s)] - f^\Delta(t)[\sigma(t) - s] \right| \leq \epsilon |\sigma(t) - s| \quad \forall s \in U. \quad (2.5)$$

We call $f^\Delta(t)$ the delta (or Hilger) derivative of f at t . The function f is delta (or Hilger) differentiable on \mathbb{T}^κ provided $f^\Delta(t)$ exists for all $t \in \mathbb{T}^\kappa$. The function $f^\Delta : \mathbb{T}^\kappa \rightarrow \mathbb{R}$ is then called the delta derivative of f on \mathbb{T}^κ .

Definition 2.3 (see [23, Definition 2.3]). Assume that $f : \mathbb{T} \rightarrow \mathbb{R}^N$ is a function, $f(t) = (f^1(t), f^2(t), \dots, f^N(t))$ and let $t \in \mathbb{T}^\kappa$. Then we define $f^\Delta(t) = (f^{1\Delta}(t), f^{2\Delta}(t), \dots, f^{N\Delta}(t))$ (provided it exists). We call $f^\Delta(t)$ the delta (or Hilger) derivative of f at t . The function f is delta (or Hilger) differentiable provided $f^\Delta(t)$ exists for all $t \in \mathbb{T}^\kappa$. The function $f^\Delta : \mathbb{T}^\kappa \rightarrow \mathbb{R}^N$ is then called the delta derivative of f on \mathbb{T}^κ .

Definition 2.4 (see [3, Definition 2.7]). For a function $f : \mathbb{T} \rightarrow \mathbb{R}$ we will talk about the second derivative f^{Δ^2} provided f^Δ is differentiable on $\mathbb{T}^{\kappa^2} = (\mathbb{T}^\kappa)^\kappa$ with derivative $f^{\Delta^2} = (f^\Delta)^\Delta : \mathbb{T}^{\kappa^2} \rightarrow \mathbb{R}$.

Definition 2.5 (see [23, Definition 2.5]). For a function $f : \mathbb{T} \rightarrow \mathbb{R}^N$ we will talk about the second derivative f^{Δ^2} provided f^Δ is differentiable on $\mathbb{T}^{\kappa^2} = (\mathbb{T}^\kappa)^\kappa$ with derivative $f^{\Delta^2} = (f^\Delta)^\Delta : \mathbb{T}^{\kappa^2} \rightarrow \mathbb{R}^N$.

Definition 2.6 (see [23, Definition 2.6]). A function $f : \mathbb{T} \rightarrow \mathbb{R}^N$ is called rd-continuous provided it is continuous at right-dense points in \mathbb{T} and its left sided limits exist (finite) at left dense points in \mathbb{T} .

Definition 2.7 (see [3, Definition 2.25]). We say that a function $w : \mathbb{T} \rightarrow \mathbb{R}$ is regressive provided

$$1 + \mu(t)w(t) \neq 0 \quad \forall t \in \mathbb{T}^\kappa \quad (2.6)$$

holds. The set of all regressive and rd-continuous functions $w : \mathbb{T} \rightarrow \mathbb{R}$ is denoted by

$$\begin{aligned} \mathcal{R} &= \mathcal{R}(\mathbb{T}) = \mathcal{R}(\mathbb{T}, \mathbb{R}), \\ \mathcal{R}^+(\mathbb{T}, \mathbb{R}) &= \{w \in \mathcal{R} : 1 + \mu(t)w(t) > 0 \quad \forall t \in \mathbb{T}\}. \end{aligned} \quad (2.7)$$

Definition 2.8 (see [7, Definition 8.2.18]). If $w \in \mathcal{R}$ and $t_0 \in \mathbb{T}$, then the unique solution of the initial value problem

$$y^\Delta = w(t)y, \quad y(t_0) = 1 \quad (2.8)$$

is called the exponential function and denoted by $e_w(\cdot, t_0)$.

The exponential function has some important properties.

Lemma 2.9 (see [3, Theorem 2.36]). *If $w \in \mathcal{R}$, then*

$$e_0(t, s) \equiv 1, \quad e_w(t, t) \equiv 1. \tag{2.9}$$

Throughout this paper, we will use the following notations:

$$\begin{aligned} C_{rd}(\mathbb{T}) &= C_{rd}(\mathbb{T}, \mathbb{R}^N) = \{f : \mathbb{T} \rightarrow \mathbb{R}^N : f \text{ is rd-continuous}\}, \\ C_{rd}^1(\mathbb{T}) &= C_{rd}^1(\mathbb{T}, \mathbb{R}^N) = \{f : \mathbb{T} \rightarrow \mathbb{R}^N : f \text{ is differentiable on } \mathbb{T}^\kappa \text{ and } f^\Delta \in C_{rd}(\mathbb{T}^\kappa)\}, \\ C_{T,rd}^1([0, T]_{\mathbb{T}}, \mathbb{R}^N) &= \{f \in C_{rd}^1([0, T]_{\mathbb{T}}, \mathbb{R}^N) : f(0) = f(T)\}. \end{aligned} \tag{2.10}$$

The Δ -measure m_Δ and Δ -integration are defined as those in [10].

Definition 2.10 (see [23, Definition 2.7]). Assume that $f : \mathbb{T} \rightarrow \mathbb{R}^N$ is a function, $f(t) = (f^1(t), f^2(t), \dots, f^N(t))$ and let A be a Δ -measurable subset of \mathbb{T} . f is integrable on A if and only if f^i ($i = 1, 2, \dots, N$) are integrable on A , and

$$\int_A f(t) \Delta t = \left(\int_A f^1(t) \Delta t, \int_A f^2(t) \Delta t, \dots, \int_A f^N(t) \Delta t \right). \tag{2.11}$$

Definition 2.11 (see [13, Definition 2.3]). Let $B \subset \mathbb{T}$. B is called Δ -null set if $m_\Delta(B) = 0$. Say that a property P holds Δ -almost everywhere (Δ -a.e.) on B , or for Δ -almost all (Δ -a.a.) $t \in B$ if there is a Δ -null set $E_0 \subset B$ such that P holds for all $t \in B \setminus E_0$.

For $p \in \mathbb{R}$, $p \geq 1$, we set the space

$$L_\Delta^p([0, T]_{\mathbb{T}}, \mathbb{R}^N) = \left\{ u : [0, T]_{\mathbb{T}} \rightarrow \mathbb{R}^N : \int_{[0, T]_{\mathbb{T}}} |f(t)|^p \Delta t < +\infty \right\} \tag{2.12}$$

with the norm

$$\|f\|_{L_\Delta^p} = \left(\int_{[0, T]_{\mathbb{T}}} |f(t)|^p \Delta t \right)^{1/p}. \tag{2.13}$$

We have the following lemma.

Lemma 2.12 (see [23, Theorem 2.1]). *Let $p \in \mathbb{R}$ be such that $p \geq 1$. Then the space $L_\Delta^p([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ is a Banach space together with the norm $\|\cdot\|_{L_\Delta^p}$. Moreover, $L_\Delta^2([a, b]_{\mathbb{T}}, \mathbb{R}^N)$ is a Hilbert space together with the inner product given for every $(f, g) \in L_\Delta^p([a, b]_{\mathbb{T}}, \mathbb{R}^N) \times L_\Delta^p([a, b]_{\mathbb{T}}, \mathbb{R}^N)$ by*

$$\langle f, g \rangle_{L_\Delta^2} = \int_{[a, b]_{\mathbb{T}}} (f(t), g(t)) \Delta t, \tag{2.14}$$

where (\cdot, \cdot) denotes the inner product in \mathbb{R}^N .

As we know from general theory of Sobolev spaces, another important class of functions is just the absolutely continuous functions on time-scales.

Definition 2.13 (see [13, Definition 2.9]). A function $f : [a, b]_{\mathbb{T}} \rightarrow \mathbb{R}$ is said to be absolutely continuous on $[a, b]_{\mathbb{T}}$ (i.e., $f \in AC([a, b]_{\mathbb{T}}, \mathbb{R})$), if for every $\epsilon > 0$, there exists $\delta > 0$ such that if $\{[a_k, b_k]_{\mathbb{T}}\}_{k=1}^n$ is a finite pairwise disjoint family of subintervals of $[a, b]_{\mathbb{T}}$ satisfying $\sum_{k=1}^n (b_k - a_k) < \delta$, then $\sum_{k=1}^n |f(b_k) - f(a_k)| < \epsilon$.

Definition 2.14 (see [23, Definition 2.11]). A function $f : [a, b]_{\mathbb{T}} \rightarrow \mathbb{R}^N$, $f(t) = (f^1(t), f^2(t), \dots, f^N(t))$. We say that f is absolutely continuous on $[a, b]_{\mathbb{T}}$ (i.e., $f \in AC([a, b]_{\mathbb{T}}, \mathbb{R}^N)$), if for every $\epsilon > 0$, there exists $\delta > 0$ such that if $\{[a_k, b_k]_{\mathbb{T}}\}_{k=1}^n$ is a finite pairwise disjoint family of subintervals of $[a, b]_{\mathbb{T}}$ satisfying $\sum_{k=1}^n (b_k - a_k) < \delta$, then $\sum_{k=1}^n |f(b_k) - f(a_k)| < \epsilon$.

Absolutely continuous functions have the following properties.

Lemma 2.15 (see [23, Theorem 2.2]). *A function $f : [a, b]_{\mathbb{T}} \rightarrow \mathbb{R}^N$ is absolutely continuous on $[a, b]_{\mathbb{T}}$ if and only if f is delta differentiable Δ -a.e. on $[a, b]_{\mathbb{T}}$ and*

$$f(t) = f(a) + \int_{[a,t]_{\mathbb{T}}} f^{\Delta}(s) \Delta s, \quad \forall t \in [a, b]_{\mathbb{T}}. \quad (2.15)$$

Lemma 2.16 (see [23, Theorem 2.3]). *Assume that functions $f, g : [a, b]_{\mathbb{T}} \rightarrow \mathbb{R}^N$ are absolutely continuous on $[a, b]_{\mathbb{T}}$. Then fg is absolutely continuous on $[a, b]_{\mathbb{T}}$ and the following equality is valid:*

$$\begin{aligned} \int_{[a,b]_{\mathbb{T}}} \left((f^{\Delta}(t), g(t)) + (f^{\sigma}(t), g^{\Delta}(t)) \right) \Delta t &= (f(b), g(b)) - (f(a), g(a)) \\ &= \int_{[a,b]_{\mathbb{T}}} \left((f(t), g^{\Delta}(t)) + (f^{\Delta}(t), g^{\sigma}(t)) \right) \Delta t. \end{aligned} \quad (2.16)$$

Now, we recall the definition and properties of the Sobolev space on $[0, T]_{\mathbb{T}}$ in [23]. For the sake of convenience, in the sequel, we will let $u^{\sigma}(t) = u(\sigma(t))$.

Definition 2.17 (see [23, Definition 2.12]). Let $p \in \mathbb{R}$ be such that $p \geq 1$ and $u : [0, T]_{\mathbb{T}} \rightarrow \mathbb{R}^N$. We say that $u \in W_{\Delta, T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ if and only if $u \in L_{\Delta}^p([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ and there exists $g : [0, T]_{\mathbb{T}}^{\kappa} \rightarrow \mathbb{R}^N$ such that $g \in L_{\Delta}^p([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ and

$$\int_{[0,T]_{\mathbb{T}}} (u(t), \phi^{\Delta}(t)) \Delta t = - \int_{[0,T]_{\mathbb{T}}} (g(t), \phi^{\sigma}(t)) \Delta t, \quad \forall \phi \in C_{T, \text{rd}}^1([0, T]_{\mathbb{T}}, \mathbb{R}^N). \quad (2.17)$$

For $p \in \mathbb{R}, p \geq 1$, we denote

$$V_{\Delta, T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N) = \left\{ x \in AC([0, T]_{\mathbb{T}}, \mathbb{R}^N) : x^{\Delta} \in L_{\Delta}^p([0, T]_{\mathbb{T}}, \mathbb{R}^N), x(0) = x(T) \right\}. \quad (2.18)$$

Remark 2.18 (see [23, Remark 2.2]). $V_{\Delta, T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N) \subset W_{\Delta, T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ is true for every $p \in \mathbb{R}$ with $p \geq 1$.

Lemma 2.19 (see [23, Theorem 2.5]). *Suppose that $u \in W_{\Delta,T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ for some $p \in \mathbb{R}$ with $p \geq 1$, and that (2.17) holds for $g \in L_{\Delta}^p([0, T]_{\mathbb{T}}, \mathbb{R}^N)$. Then, there exists a unique function $x \in V_{\Delta,T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ such that the equalities*

$$x = u, \quad x^{\Delta} = g \quad \Delta\text{-a.e. on } [0, T]_{\mathbb{T}} \tag{2.19}$$

are satisfied and

$$\int_{[0, T]_{\mathbb{T}}} g(t) \Delta t = 0. \tag{2.20}$$

Lemma 2.20 (see [3, Theorem 1.16]). *Assume that $f : \mathbb{T} \rightarrow \mathbb{R}$ is a function and let $t \in \mathbb{T}^{\kappa}$. Then, one has the following.*

- (i) *If f is differentiable at t , then f is continuous at t .*
- (ii) *If f is differentiable at t , then*

$$f(\sigma(t)) = f(t) + \mu(t) f^{\Delta}(t). \tag{2.21}$$

By identifying $u \in W_{\Delta,T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ with its absolutely continuous representative $x \in V_{\Delta,T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ for which (2.19) holds, the set $W_{\Delta,T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ can be endowed with the structure of Banach space.

Theorem 2.21. *Assume $p \in \mathbb{R}$ and $p \geq 1$. The set $W_{\Delta,T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ is a Banach space together with the norm defined as*

$$\|u\|_{W_{\Delta,T}^{1,p}} = \left(\int_{[0, T]_{\mathbb{T}}} |u^{\sigma}(t)|^p \Delta t + \int_{[0, T]_{\mathbb{T}}} |u^{\Delta}(t)|^p \Delta t \right)^{1/p} \quad \forall u \in W_{\Delta,T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N). \tag{2.22}$$

Moreover, the set $H_{\Delta,T}^1 = W_{\Delta,T}^{1,2}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ is a Hilbert space together with the inner product

$$\langle u, v \rangle_{H_{\Delta,T}^1} = \int_{[0, T]_{\mathbb{T}}} (u^{\sigma}(t), v^{\sigma}(t)) \Delta t + \int_{[0, T]_{\mathbb{T}}} (u^{\Delta}(t), v^{\Delta}(t)) \Delta t \quad \forall u, v \in H_{\Delta,T}^1. \tag{2.23}$$

Proof. Let $\{u_n\}_{n \in \mathbb{N}}$ be a Cauchy sequence in $W_{\Delta,T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$. That is, $\{u_n\}_{n \in \mathbb{N}} \subset L_{\Delta}^p([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ and there exist $g_n : [0, T]_{\mathbb{T}} \rightarrow \mathbb{R}^N$ such that $\{g_n\}_{n \in \mathbb{N}} \subset L_{\Delta}^p([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ and

$$\int_{[0, T]_{\mathbb{T}}} (u_n(t), \phi^{\Delta}(t)) \Delta t = - \int_{[0, T]_{\mathbb{T}}} (g_n(t), \phi^{\sigma}(t)) \Delta t, \quad \forall \phi \in C_{T,rd}^1([0, T]_{\mathbb{T}}, \mathbb{R}^N). \tag{2.24}$$

Thus, by Lemma 2.19, there exists $\{x_n\}_{n \in \mathbb{N}} \subset V_{\Delta,T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ such that

$$x_n = u_n, \quad x_n^{\Delta} = g_n \quad \Delta\text{-a.e. on } [0, T]_{\mathbb{T}}. \tag{2.25}$$

Combining (2.24) and (2.25), we obtain

$$\int_{[0,T]_{\mathbb{T}}} (x_n(t), \phi^\Delta(t)) \Delta t = - \int_{[0,T]_{\mathbb{T}}} (x_n^\Delta(t), \phi^\sigma(t)) \Delta t, \quad \forall \phi \in C_{T,\text{rd}}^1([0, T]_{\mathbb{T}}, \mathbb{R}^N). \quad (2.26)$$

Since $\{u_n\}_{n \in \mathbb{N}}$ is a Cauchy sequence in $W_{\Delta, T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$, by (2.22), one has

$$\int_{[0,T]_{\mathbb{T}}} |u_n^\sigma(t) - u_m^\sigma(t)|^2 \Delta t \longrightarrow 0 \quad (m, n \longrightarrow \infty), \quad (2.27)$$

$$\int_{[0,T]_{\mathbb{T}}} |u_n^\Delta(t) - u_m^\Delta(t)|^2 \Delta t \longrightarrow 0 \quad (m, n \longrightarrow \infty). \quad (2.28)$$

It follows from Lemma 2.20, (2.27), and (2.28) that

$$\begin{aligned} \int_{[0,T]_{\mathbb{T}}} |u_n(t) - u_m(t)|^2 \Delta t &= \int_{[0,T]_{\mathbb{T}}} \left| (u_n^\sigma(t) - u_m^\sigma(t)) - \mu(t)(u_n^\Delta(t) - u_m^\Delta(t)) \right|^2 \Delta t \\ &\leq 2 \int_{[0,T]_{\mathbb{T}}} |u_n^\sigma(t) - u_m^\sigma(t)|^2 \Delta t + 2(\sigma(T))^2 \int_{[0,T]_{\mathbb{T}}} |u_n^\Delta(t) - u_m^\Delta(t)|^2 \Delta t \\ &\longrightarrow 0 \quad (m, n \longrightarrow \infty). \end{aligned} \quad (2.29)$$

By Lemma 2.12, (2.28) and (2.29), there exist $u, g \in L_\Delta^p([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ such that

$$\|u_n - u\|_{L_\Delta^p} \longrightarrow 0 \quad (n \longrightarrow \infty), \quad \left\| u_n^\Delta - g \right\|_{L_\Delta^p} \longrightarrow 0 \quad (n \longrightarrow \infty). \quad (2.30)$$

From (2.26) and (2.30), one has

$$\int_{[0,T]_{\mathbb{T}}} (u(t), \phi^\Delta(t)) \Delta t = - \int_{[0,T]_{\mathbb{T}}} (g(t), \phi^\sigma(t)) \Delta t, \quad \forall \phi \in C_{T,\text{rd}}^1([0, T]_{\mathbb{T}}, \mathbb{R}^N). \quad (2.31)$$

From (2.31), we conclude that $u \in W_{\Delta,T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$. Moreover, by Lemma 2.20 and (2.30), one has

$$\begin{aligned} \int_{[0,T]_{\mathbb{T}}} |u_n^\sigma(t) - u^\sigma(t)|^2 \Delta t &= \int_{[0,T]_{\mathbb{T}}} \left| (u_n(t) - u(t)) + \mu(t)(u_n^\Delta(t) - u^\Delta(t)) \right|^2 \Delta t \\ &= \int_{[0,T]_{\mathbb{T}}} \left| (u_n(t) - u(t)) + \mu(t)(u_n^\Delta(t) - g(t)) \right|^2 \Delta t \\ &\leq 2 \int_{[0,T]_{\mathbb{T}}} |u_n(t) - u(t)|^2 \Delta t + 2(\sigma(T))^2 \int_{[0,T]_{\mathbb{T}}} |u_n^\Delta(t) - g(t)|^2 \Delta t \\ &\rightarrow 0 \quad (n \rightarrow \infty). \end{aligned} \tag{2.32}$$

Thereby, it follows from Remark 2.18, (2.30), (2.32), and Lemma 2.19 that there exists $x \in V_{\Delta,T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N) \subset W_{\Delta,T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ such that

$$\|u_n - x\|_{W_{\Delta,T}^{1,p}} \rightarrow 0 \quad (n \rightarrow \infty). \tag{2.33}$$

Obviously, the set $H_{\Delta,T}^1$ is a Hilbert space together with the inner product

$$\langle u, v \rangle_{H_{\Delta,T}^1} = \int_{[0,T]_{\mathbb{T}}} (u^\sigma(t), v^\sigma(t)) \Delta t + \int_{[0,T]_{\mathbb{T}}} (u^\Delta(t), v^\Delta(t)) \Delta t \quad \forall u, v \in H_{\Delta,T}^1. \tag{2.34}$$

□

We will derive some properties of the Banach space $W_{\Delta,T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$.

Lemma 2.22 (see [10, Theorem A.2]). *Let $f : [a, b]_{\mathbb{T}} \rightarrow \mathbb{R}$ be a continuous function on $[a, b]_{\mathbb{T}}$ which is delta differentiable on $[a, b]_{\mathbb{T}}$. Then there exist $\xi, \tau \in [a, b]_{\mathbb{T}}$ such that*

$$f^\Delta(\tau) \leq \frac{f(b) - f(a)}{b - a} \leq f^\Delta(\xi). \tag{2.35}$$

Theorem 2.23. *There exists $K = K(p) > 0$ such that the inequality*

$$\|u\|_\infty \leq K \|u\|_{W_{\Delta,T}^{1,p}} \tag{2.36}$$

holds for all $u \in W_{\Delta,T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$, where $\|u\|_\infty = \max_{t \in [0, T]_{\mathbb{T}}} |u(t)|$.
Moreover, if $\int_{[0, T]_{\mathbb{T}}} u(t) \Delta t = 0$, then

$$\|u\|_\infty \leq K \left\| u^\Delta \right\|_{L_\Delta^p}. \tag{2.37}$$

Proof. Going to the components of u , we can assume that $N = 1$. If $u \in W_{\Delta, T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R})$, by Lemma 2.19, $U(t) = \int_{[0, t]_{\mathbb{T}}} u(s) \Delta s$ is absolutely continuous on $[a, b]_{\mathbb{T}}$. It follows from Lemma 2.22 that there exists $\zeta \in [a, b]_{\mathbb{T}}$ such that

$$u(\zeta) \leq \frac{U(T) - U(0)}{T} = \frac{1}{T} \int_{[0, T]_{\mathbb{T}}} u(s) \Delta s. \quad (2.38)$$

Hence, for $t \in [a, b]_{\mathbb{T}}$, using Lemma 2.15, (2.38), and Hölder's inequality, one has that

$$\begin{aligned} |u(t)| &= \left| u(\zeta) + \int_{[\zeta, t]_{\mathbb{T}}} u^{\Delta}(s) \Delta s \right| \\ &\leq |u(\zeta)| + \int_{[0, T]_{\mathbb{T}}} |u^{\Delta}(s)| \Delta s \\ &\leq \frac{1}{T} \left| \int_{[0, T]_{\mathbb{T}}} u(s) \Delta s \right| + T^{1/q} \left(\int_{[0, T]_{\mathbb{T}}} |u^{\Delta}(s)|^p \Delta s \right)^{1/p}, \end{aligned} \quad (2.39)$$

where $1/p + 1/q = 1$. If $\int_{[0, T]_{\mathbb{T}}} u(t) \Delta t = 0$, by (2.39), we obtain (2.37). In the general case, for $t \in [a, b]_{\mathbb{T}}$, by Lemma 2.20 and Hölder's inequality, we get

$$\begin{aligned} |u(t)| &\leq \frac{1}{T} \left| \int_{[0, T]_{\mathbb{T}}} u(s) \Delta s \right| + T^{1/q} \left(\int_{[0, T]_{\mathbb{T}}} |u^{\Delta}(s)|^p \Delta s \right)^{1/p} \\ &\leq \frac{1}{T} \int_{[0, T]_{\mathbb{T}}} |u(s)| \Delta s + T^{1/q} \left(\int_{[0, T]_{\mathbb{T}}} |u^{\Delta}(s)|^p \Delta s \right)^{1/p} \\ &= \frac{1}{T} \int_{[0, T]_{\mathbb{T}}} |u^{\sigma}(s) - \mu(s)u^{\Delta}(s)| \Delta s + T^{1/q} \left(\int_{[0, T]_{\mathbb{T}}} |u^{\Delta}(s)|^p \Delta s \right)^{1/p} \\ &= \frac{1}{T} \int_{[0, T]_{\mathbb{T}}} |u^{\sigma}(s)| \Delta s + \frac{1}{T} \sigma(T) \int_{[0, T]_{\mathbb{T}}} |u^{\Delta}(s)| \Delta s + T^{1/q} \left(\int_{[0, T]_{\mathbb{T}}} |u^{\Delta}(s)|^p \Delta s \right)^{1/p} \\ &\leq T^{(-1/p)} \left(\int_{[0, T]_{\mathbb{T}}} |u^{\sigma}(s)|^p \Delta s \right)^{1/p} + T^{(-1/p)} \sigma(T) \left(\int_{[0, T]_{\mathbb{T}}} |u^{\Delta}(s)|^p \Delta s \right)^{1/p} \\ &\quad + T^{1/q} \left(\int_{[0, T]_{\mathbb{T}}} |u^{\Delta}(s)|^p \Delta s \right)^{1/p} \\ &\leq \left(T^{(-1/p)} + T^{(-1/p)} \sigma(T) + T^{1/q} \right) \|u\|_{W_{\Delta, T}^{1,p}}. \end{aligned} \quad (2.40)$$

From (2.40), (2.36) holds. \square

Remark 2.24. It follows from Theorem 2.23 that $W_{\Delta, T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ is continuously immersed into $C([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ with the norm $\|\cdot\|_{\infty}$.

Theorem 2.25. *If the sequence $\{u_k\}_{k \in \mathbb{N}} \subset W_{\Delta, T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ converges weakly to u in $W_{\Delta, T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$, then $\{u_k\}_{k \in \mathbb{N}}$ converges strongly in $C([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ to u .*

Proof. Since $u_k \rightharpoonup u$ in $W_{\Delta, T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$, $\{u_k\}_{k \in \mathbb{N}}$ is bounded in $W_{\Delta, T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ and, hence, in $C([0, T]_{\mathbb{T}}, \mathbb{R}^N)$. It follows from Remark 2.24 that $u_k \rightharpoonup u$ in $C([0, T]_{\mathbb{T}}, \mathbb{R}^N)$. For $t_1, t_2 \in [0, T]_{\mathbb{T}}$, $t_1 \leq t_2$, there exists $C_1 > 0$ such that

$$\begin{aligned} |u_k(t_2) - u_k(t_1)| &\leq \int_{[t_1, t_2]_{\mathbb{T}}} |u_k^\Delta(s)| \Delta s \\ &\leq (t_2 - t_1)^{1/q} \left(\int_{[t_1, t_2]_{\mathbb{T}}} |u_k^\Delta(s)|^p \Delta s \right)^{1/p} \\ &\leq (t_2 - t_1)^{1/q} \|u_k\|_{W_{\Delta, T}^{1,p}} \\ &\leq C_1 (t_2 - t_1)^{1/q}. \end{aligned} \tag{2.41}$$

Hence, the sequence $\{u_k\}_{k \in \mathbb{N}}$ is equicontinuous. By Ascoli-Arzelà theorem, $\{u_k\}_{k \in \mathbb{N}}$ is relatively compact in $C([0, T]_{\mathbb{T}}, \mathbb{R}^N)$. By the uniqueness of the weak limit in $C([0, T]_{\mathbb{T}}, \mathbb{R}^N)$, every uniformly convergent subsequence of $\{u_k\}_{k \in \mathbb{N}}$ converges to u . Thus, $\{u_k\}_{k \in \mathbb{N}}$ converges strongly in $C([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ to u . \square

Remark 2.26. By Theorem 2.25, the immersion $W_{\Delta, T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N) \hookrightarrow C([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ is compact.

Theorem 2.27. *Let $L : [0, T]_{\mathbb{T}} \times \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$, $(t, x, y) \rightarrow L(t, x, y)$ be Lebesgue Δ -measurable in t for each $(x, y) \in \mathbb{R}^N \times \mathbb{R}^N$ and continuously differentiable in (x, y) for every $t \in [0, T]_{\mathbb{T}}$. If there exist $a \in C(\mathbb{R}^+, \mathbb{R}^+)$, $b, c \in [0, T]_{\mathbb{T}} \rightarrow \mathbb{R}^+$, $b^\sigma \in L_{\Delta}^1([0, T]_{\mathbb{T}}, \mathbb{R}^+)$ and $c^\sigma \in L_{\Delta}^q([0, T]_{\mathbb{T}}, \mathbb{R}^+)$ ($1 < q < +\infty$) such that for Δ -almost $t \in [0, T]_{\mathbb{T}}$ and every $(x, y) \in \mathbb{R}^N \times \mathbb{R}^N$, one has*

$$\begin{aligned} |L(t, x, y)| &\leq a(|x|)(b(t) + |y|^p), \\ |L_x(t, x, y)| &\leq a(|x|)(b(t) + |y|^p), \\ |L_y(t, x, y)| &\leq a(|x|)(c(t) + |y|^{p-1}), \end{aligned} \tag{2.42}$$

where $1/p + 1/q = 1$, then the functional $\Phi : W_{\Delta, T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N) \rightarrow \mathbb{R}$ defined as

$$\Phi(u) = \int_{[0, T]_{\mathbb{T}}} L(\sigma(t), u^\sigma(t), u^\Delta(t)) \Delta t \tag{2.43}$$

is continuously differentiable on $W_{\Delta,T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ and

$$\langle \Phi'(u), v \rangle = \int_{[0, T]_{\mathbb{T}}} \left[\left(L_x(\sigma(t), u^\sigma(t), u^\Delta(t)), v^\sigma(t) \right) + \left(L_y(\sigma(t), u^\sigma(t), u^\Delta(t)), v^\Delta(t) \right) \right] \Delta t. \quad (2.44)$$

Proof. It suffices to prove that Φ has at every point u a directional derivative $\Phi'(u) \in (W_{\Delta,T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N))^*$ given by (2.44) and that the mapping

$$\Phi' : W_{\Delta,T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N) \longrightarrow (W_{\Delta,T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N))^* \quad (2.45)$$

is continuous.

Firstly, it follows from (2.42) that Φ is everywhere finite on $W_{\Delta,T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$. We define, for u and v fixed in $W_{\Delta,T}^{1,p}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$, $t \in [0, T]_{\mathbb{T}}$, $\lambda \in [-1, 1]$,

$$\begin{aligned} G(\lambda, t) &= L(\sigma(t), u^\sigma(t) + \lambda v^\sigma(t), u^\Delta(t) + \lambda v^\Delta(t)), \\ \Psi(\lambda) &= \int_{[0, T]_{\mathbb{T}}} G(\lambda, t) \Delta t = \Phi(u + \lambda v). \end{aligned} \quad (2.46)$$

From (2.42), one has

$$\begin{aligned} |D_\lambda G(\lambda, t)| &\leq \left| \left(D_x L(\sigma(t), u^\sigma(t) + \lambda v^\sigma(t), u^\Delta(t) + \lambda v^\Delta(t)), v^\sigma(t) \right) \right| \\ &\quad + \left| \left(D_y L(\sigma(t), u^\sigma(t) + \lambda v^\sigma(t), u^\Delta(t) + \lambda v^\Delta(t)), v^\Delta(t) \right) \right| \\ &\leq a(|u^\sigma(t) + \lambda v^\sigma(t)|) \left(b^\sigma(t) + |u^\Delta(t) + \lambda v^\Delta(t)|^p \right) |v^\sigma(t)| \\ &\quad + a(|u^\sigma(t) + \lambda v^\sigma(t)|) \left(c^\sigma(t) + |u^\Delta(t) + \lambda v^\Delta(t)|^{p-1} \right) |v^\Delta(t)| \\ &\leq \bar{a} \left(b^\sigma(t) + (|u^\Delta(t)| + |v^\Delta(t)|)^p \right) |v^\sigma(t)| \\ &\quad + \bar{a} \left(c^\sigma(t) + (|u^\Delta(t)| + |v^\Delta(t)|)^{p-1} \right) |v^\Delta(t)| \\ &\triangleq d(t), \end{aligned} \quad (2.47)$$

where

$$\bar{a} = \max_{(\lambda, t) \in [-1, 1] \times [0, T]_{\mathbb{T}}} a(|u(t) + \lambda v(t)|), \quad (2.48)$$

thus, $d \in L^1_\Delta([0, T]_{\mathbb{T}}, \mathbb{R}^+)$. Since $b^\sigma \in L^1_\Delta([0, T]_{\mathbb{T}}, \mathbb{R}^+)$, $(|u^\Delta| + |v^\Delta|)^p \in L^1_\Delta([0, T]_{\mathbb{T}}, \mathbb{R})$, $c^\sigma \in L^q_\Delta([0, T]_{\mathbb{T}}, \mathbb{R}^+)$, one has

$$|D_\lambda G(\lambda, t)| \leq d(t), \tag{2.49}$$

$$\begin{aligned} \Psi'(0) &= \int_{[0, T]_{\mathbb{T}}} D_\lambda G(0, t) \Delta t \\ &= \int_{[0, T]_{\mathbb{T}}} \left[\left(D_x L(\sigma(t), u^\sigma(t), u^\Delta(t)), v^\sigma(t) \right) + \left(D_y L(\sigma(t), u^\sigma(t), u^\Delta(t)), v^\Delta(t) \right) \right] \Delta t. \end{aligned} \tag{2.50}$$

On the other hand, it follows from (2.42) that

$$\begin{aligned} \left| D_x L(\sigma(t), u^\sigma(t), u^\Delta(t)) \right| &\leq a(|u^\sigma(t)|) \left(b^\sigma(t) + |u^\Delta(t)|^p \right) \triangleq \psi_1(t), \\ \left| D_y L(\sigma(t), u^\sigma(t), u^\Delta(t)) \right| &\leq a(|u^\sigma(t)|) \left(c^\sigma(t) + |u^\Delta(t)|^{p-1} \right) \triangleq \psi_2(t), \end{aligned} \tag{2.51}$$

thus $\psi_1 \in L^1_\Delta([0, T]_{\mathbb{T}}, \mathbb{R}^+)$, $\psi_2 \in L^q_\Delta([0, T]_{\mathbb{T}}, \mathbb{R}^+)$. Thereby, by Theorem 2.23, (2.50), and (2.51), there exist positive constants C_2, C_3, C_4 such that

$$\begin{aligned} &\int_{[0, T]_{\mathbb{T}}} \left[\left(D_x L(\sigma(t), u^\sigma(t), u^\Delta(t)), v^\sigma(t) \right) + \left(D_y L(\sigma(t), u^\sigma(t), u^\Delta(t)), v^\Delta(t) \right) \right] \Delta t \\ &\leq C_2 \|v\|_\infty + C_3 \|v^\Delta\|_{L^p_\Delta} \\ &\leq C_4 \|v\|_{W^{1,p}_{\Delta, T}} \end{aligned} \tag{2.52}$$

and Φ has a directional derivative at u and $\Phi'(u) \in (W^{1,p}_{\Delta, T}([0, T]_{\mathbb{T}}, \mathbb{R}^N))^*$ given by (2.44).

Moreover, (2.42) implies that the mapping from $W^{1,p}_{\Delta, T}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ into $L^1_\Delta([0, T]_{\mathbb{T}}, \mathbb{R}^N) \times L^q_\Delta([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ defined by

$$u \longrightarrow \left(D_x L(\cdot, u^\sigma, u^\Delta), D_y L(\cdot, u^\sigma, u^\Delta) \right) \tag{2.53}$$

is continuous, so that Φ' is continuous from $W^{1,p}_{\Delta, T}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ into $(W^{1,p}_{\Delta, T}([0, T]_{\mathbb{T}}, \mathbb{R}^N))^*$. \square

3. Variational Setting

In this section, in order to apply the critical point theory, we make a variational structure. From this variational structure, we can reduce the problem of finding solutions of problem (1.1) to one of seeking the critical points of a corresponding functional.

By Theorem 2.21, the space $H_{\Delta,T}^1 = W_{\Delta,T}^{1,2}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ with the inner product

$$\langle u, v \rangle_{H_{\Delta,T}^1} = \int_{[0,T]_{\mathbb{T}}} (u^\sigma(t), v^\sigma(t)) \Delta t + \int_{[0,T]_{\mathbb{T}}} (u^\Delta(t), v^\Delta(t)) \Delta t \quad (3.1)$$

and the induced norm

$$\|u\|_{H_{\Delta,T}^1} = \left(\int_{[0,T]_{\mathbb{T}}} |u^\sigma(t)|^2 \Delta t + \int_{[0,T]_{\mathbb{T}}} |u^\Delta(t)|^2 \Delta t \right)^{1/2} \quad (3.2)$$

is Hilbert space.

Since $w \in \mathcal{R}^+([0, T]_{\mathbb{T}}, \mathbb{R})$, by Theorem 2.44 in [3], one has that

$$e_w(t, 0) > 0 \quad \forall t \in [0, T]_{\mathbb{T}}, \quad (3.3)$$

in $H_{\Delta,T}^1$, we also consider the inner product

$$\langle u, v \rangle = \int_{[0,T]_{\mathbb{T}}} e_w(t, 0) (u^\sigma(t), v^\sigma(t)) \Delta t + \int_{[0,T]_{\mathbb{T}}} e_w(t, 0) (u^\Delta(t), v^\Delta(t)) \Delta t \quad (3.4)$$

and the induced norm

$$\|u\| = \left(\int_{[0,T]_{\mathbb{T}}} e_w(t, 0) |u^\sigma(t)|^2 \Delta t + \int_{[0,T]_{\mathbb{T}}} e_w(t, 0) |u^\Delta(t)|^2 \Delta t \right)^{1/2}, \quad (3.5)$$

we prove the following theorem.

Theorem 3.1. *The norm $\|\cdot\|$ and $\|\cdot\|_{H_{\Delta,T}^1}$ are equivalent.*

Proof. Since $e_w(\cdot, 0)$ is continuous on $[0, T]_{\mathbb{T}}$ and

$$e_w(t, 0) > 0 \quad \forall t \in [0, T]_{\mathbb{T}}, \quad (3.6)$$

there exist two positive constants M_1 and M_2 such that

$$M_1 = \min_{t \in [0, T]_{\mathbb{T}}} e_w(t, 0), \quad M_2 = \max_{t \in [0, T]_{\mathbb{T}}} e_w(t, 0). \quad (3.7)$$

Hence, one has

$$\sqrt{M_1} \|u\|_{H_{\Delta,T}^1} \leq \|u\| \leq \sqrt{M_2} \|u\|_{H_{\Delta,T}^1}, \quad \forall u \in H_{\Delta,T}^1. \quad (3.8)$$

Consequently, the norm $\|\cdot\|$ and $\|\cdot\|_{H_{\Delta,T}^1}$ are equivalent. \square

Consider the functional $\varphi : H_{\Delta,T}^1 \rightarrow \mathbb{R}$ defined by

$$\varphi(u) = \frac{1}{2} \int_{[0,T]_{\mathbb{T}}} e_w(t,0) |u^\Delta(t)|^2 \Delta t + \int_{[0,T]_{\mathbb{T}}} e_w(t,0) F(\sigma(t), u^\sigma(t)) \Delta t. \quad (3.9)$$

We have the following facts.

Theorem 3.2. *The functional φ is continuously differentiable on $H_{\Delta,T}^1$ and*

$$\langle \varphi'(u), v \rangle = \int_{[0,T]_{\mathbb{T}}} e_w(t,0) (u^\Delta(t), v^\Delta(t)) \Delta t + \int_{[0,T]_{\mathbb{T}}} e_w(t,0) (\nabla F(\sigma(t), u^\sigma(t)), v^\sigma(t)) \Delta t \quad (3.10)$$

for all $v \in H_{\Delta,T}^1$.

Proof. Let $L(t, x, y) = e_w(\rho(t), 0) [(1/2)|y|^2 + F(t, x)]$ for all $x, y \in \mathbb{R}^N$ and $t \in [0, T]_{\mathbb{T}}$. Then, by condition (A), $L(t, x, y)$ satisfies all assumptions of Theorem 2.27. Hence, by Theorem 2.27, we know that the functional φ is continuously differentiable on $H_{\Delta,T}^1$ and

$$\langle \varphi'(u), v \rangle = \int_{[0,T]_{\mathbb{T}}} e_w(t,0) (u^\Delta(t), v^\Delta(t)) \Delta t + \int_{[0,T]_{\mathbb{T}}} e_w(t,0) (\nabla F(\sigma(t), u^\sigma(t)), v^\sigma(t)) \Delta t \quad (3.11)$$

for all $v \in H_{\Delta,T}^1$. □

Theorem 3.3. *If $u \in H_{\Delta,T}^1$ is a critical point of φ in $H_{\Delta,T}^1$, that is, $\varphi'(u) = 0$, then u is a solution of problem (1.1).*

Proof. Since $\varphi'(u) = 0$, it follows from Theorem 3.2 that

$$\int_{[0,T]_{\mathbb{T}}} e_w(t,0) (u^\Delta(t), v^\Delta(t)) \Delta t + \int_{[0,T]_{\mathbb{T}}} e_w(t,0) (\nabla F(\sigma(t), u^\sigma(t)), v^\sigma(t)) \Delta t = 0 \quad (3.12)$$

for all $v \in H_{\Delta,T}^1$, that is,

$$\int_{[0,T]_{\mathbb{T}}} e_w(t,0) (u^\Delta(t), v^\Delta(t)) \Delta t = - \int_{[0,T]_{\mathbb{T}}} e_w(t,0) (\nabla F(\sigma(t), u^\sigma(t)), v^\sigma(t)) \Delta t \quad (3.13)$$

for all $v \in H_{\Delta,T}^1$. From condition (A) and Definition 2.17, one has that $e_w(\cdot, 0)u^\Delta \in H_{\Delta,T}^1$. By Lemma 2.19 and (2.20), there exists a unique function $x \in V_{\Delta,T}^{1,2}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ such that

$$x = u, \quad (e_w(t,0)x^\Delta(t))^\Delta = e_w(t,0)\nabla F(\sigma(t), u^\sigma(t)) \quad \Delta\text{-a.e. on } [0, T]_{\mathbb{T}}^k, \quad (3.14)$$

$$\int_{[0,T]_{\mathbb{T}}} e_w(t,0)\nabla F(\sigma(t), u^\sigma(t)) \Delta t = 0. \quad (3.15)$$

By (3.14), one has

$$e_w(t,0)x^{\Delta^2}(t) + w(t)e_w(t,0)x^\Delta(\sigma(t)) = e_w(t,0)\nabla F(\sigma(t), u^\sigma(t)) \quad \Delta\text{-a.e. on } [0, T]_{\mathbb{T}}^{\kappa}. \quad (3.16)$$

Combining (3.14), (3.15), (3.16), and Lemma 2.19, we obtain

$$\begin{aligned} x^{\Delta^2}(t) + w(t)x^\Delta(\sigma(t)) &= \nabla F(\sigma(t), u^\sigma(t)) \quad \Delta\text{-a.e. on } [0, T]_{\mathbb{T}}^{\kappa}, \\ x(0) - x(T) &= 0, \quad x^\Delta(0) - x^\Delta(T) = 0. \end{aligned} \quad (3.17)$$

We identify $u \in H_{\Delta, T}^1$ with its absolutely continuous representative $x \in V_{\Delta, T}^{1,2}([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ for which (3.14) holds. Thus u is a solution of problem (1.1). \square

Theorem 3.4. *The functional φ is weakly lower semicontinuous on $H_{\Delta, T}^1$.*

Proof. Let

$$\begin{aligned} \varphi_1(u) &= \frac{1}{2} \int_{[0, T]_{\mathbb{T}}} e_w(t,0) \left| u^\Delta(t) \right|^2 \Delta t, \\ \varphi_2(u) &= \int_{[0, T]_{\mathbb{T}}} e_w(t,0) F(\sigma(t), u^\sigma(t)) \Delta t. \end{aligned} \quad (3.18)$$

Then, φ_1 is continuous and convex. Hence, φ_1 is weakly lower semicontinuous. On the other hand, let $\{u_n\}_{n \in \mathbb{N}} \subset H_{\Delta, T}^1$, $u_n \rightharpoonup u$ in $H_{\Delta, T}^1$. By Theorem 2.25, $\{u_k\}_{k \in \mathbb{N}}$ converges strongly in $C([0, T]_{\mathbb{T}}, \mathbb{R}^N)$ to u . By condition (A), one has

$$\begin{aligned} |\varphi_2(u_n) - \varphi_2(u)| &= \left| \int_{[0, T]_{\mathbb{T}}} e_w(t,0) F(\sigma(t), u_n^\sigma(t)) \Delta t - \int_{[0, T]_{\mathbb{T}}} e_w(t,0) F(\sigma(t), u^\sigma(t)) \Delta t \right| \\ &\leq M_2 \int_{[0, T]_{\mathbb{T}}} |F(\sigma(t), u_n^\sigma(t)) - F(\sigma(t), u^\sigma(t))| \Delta t \\ &\longrightarrow 0. \end{aligned} \quad (3.19)$$

Thus, φ_2 is weakly continuous. Consequently, $\varphi = \varphi_1 + \varphi_2$ is weakly lower semicontinuous. \square

To prove the existence of solutions for problem (1.1), we need the following definitions.

Definition 3.5 (see [23, page 81]). Let X be a real Banach space, $I \in C^1(X, \mathbb{R})$ and $c \in \mathbb{R}$. I is said to satisfy $(PS)_c$ -condition on X if the existence of a sequence $\{x_n\} \subseteq X$ such that $I(x_n) \rightarrow c$ and $I'(x_n) \rightarrow 0$ as $n \rightarrow \infty$, implies that c is a critical value of I .

Definition 3.6 (see [23, page 81]). Let X be a real Banach space and $I \in C^1(X, \mathbb{R})$. I is said to satisfy P.S. condition on X if any sequence $\{x_n\} \subseteq X$ for which $I(x_n)$ is bounded and $I'(x_n) \rightarrow 0$ as $n \rightarrow \infty$, possesses a convergent subsequence in X .

Remark 3.7. It is clear that the P.S. condition implies the $(PS)_c$ -condition for each $c \in \mathbb{R}$.

We also need the following result to prove our main results of this paper.

Lemma 3.8 (see [24, Theorem 1.1]). *If φ is weakly lower semicontinuous on a reflexive Banach space X and has a bounded minimizing sequence, then φ has a minimum on X .*

Lemma 3.9 (see [24, Theorem 4.7]). *Let X be a Banach space and let $J \in C^1(X, \mathbb{R})$, $R > 0$. Assume that X splits into a direct sum of closed subspace $X = X^- \oplus X^+$ with $\dim X^- < \infty$ and $\sup_{S_R^-} J < \inf_{X^+} J$, where $S_R^- = \{u \in X^- : \|u\| = R\}$. Let $B_R^- = \{u \in X^- : \|u\| \leq R\}$, $M = \{h \in C(B_R^-, X) : h(s) = s \text{ if } s \in S_R^-\}$ and $c = \inf_{h \in M} \max_{s \in B_R^-} J(h(s))$. Then, if J satisfies the $(PS)_c$ condition, c is a critical value of J .*

Lemma 3.10 (see [24, Proposition 1.4]). *Let $G \in C^1(\mathbb{R}^N, \mathbb{R})$ be a convex function. Then, for all $x, y \in \mathbb{R}^N$ one has*

$$G(x) \geq G(y) + (\nabla G(y), x - y). \tag{3.20}$$

4. Existence of Solutions

For $u \in H^1_{\Delta, T}$, let $\bar{u} = (1/T) \int_{[0, T]_{\mathbb{T}}} u(t) \Delta t$ and $\tilde{u}(t) = u(t) - \bar{u}$, then $\int_{[0, T]_{\mathbb{T}}} \tilde{u}(t) \Delta t = 0$. We have the following existence results.

Theorem 4.1. *Assume that (A) and the following conditions are satisfied.*

- (i) *There exist $f, g : [0, T]_{\mathbb{T}} \rightarrow \mathbb{R}^+$ and $\alpha \in [0, 1)$ such that $f^\sigma, g^\sigma \in L^1_{\Delta}([0, T]_{\mathbb{T}}, \mathbb{R}^+)$ and*

$$|\nabla F(t, x)| \leq f(t)|x|^\alpha + g(t) \tag{4.1}$$

for all $x \in \mathbb{R}^N$ and Δ -a.e. $t \in [0, T]_{\mathbb{T}}$.

- (ii) $|x|^{-2\alpha} \int_{[0, T]_{\mathbb{T}}} e_w(t, 0) F(\sigma(t), x) \Delta t \rightarrow +\infty$ as $|x| \rightarrow \infty$.

Then problem (1.1) has at least one solution which minimizes the function φ .

Proof. By Theorem 2.23, there exists $C_5 > 0$ such that

$$\|\tilde{u}\|_\infty^2 \leq C_5 \int_{[0, T]_{\mathbb{T}}} |u^\Delta(t)|^2 \Delta t. \tag{4.2}$$

It follows from (i), Theorem 2.23 and (4.2) that

$$\begin{aligned}
& \left| \int_{[0,T]_{\mathbb{T}}} e_w(t,0)(F(\sigma(t), u^\sigma(t)) - F(\sigma(t), \bar{u})) \Delta t \right| \\
& \leq \left| \int_{[0,T]_{\mathbb{T}}} e_w(t,0) \left(\int_0^1 (\nabla F(\sigma(t), \bar{u} + s\tilde{u}^\sigma(t)), \tilde{u}^\sigma(t)) ds \right) \Delta t \right| \\
& \leq M_2 \int_{[0,T]_{\mathbb{T}}} \left(\int_0^1 f^\sigma(t) |\bar{u} + s\tilde{u}^\sigma(t)|^\alpha |\tilde{u}^\sigma(t)| ds \right) \Delta t + M_2 \int_{[0,T]_{\mathbb{T}}} \left(\int_0^1 g^\sigma(t) |\tilde{u}^\sigma(t)| ds \right) \Delta t \\
& \leq 2M_2 (|\bar{u}|^\alpha + \|\tilde{u}\|_\infty^\alpha) \|\tilde{u}\|_\infty \int_{[0,T]_{\mathbb{T}}} f^\sigma(t) \Delta t + M_2 \|\tilde{u}\|_\infty \int_{[0,T]_{\mathbb{T}}} g^\sigma(t) \Delta t \\
& \leq \frac{M_1}{4C_5} \|\tilde{u}\|_\infty^2 + \frac{4M_2^2 C_5}{M_1} |\bar{u}|^{2\alpha} \left(\int_{[0,T]_{\mathbb{T}}} f^\sigma(t) \Delta t \right)^2 \\
& \quad + 2M_2 \|\tilde{u}\|_\infty^{\alpha+1} \int_{[0,T]_{\mathbb{T}}} f^\sigma(t) dt + M_2 \|\tilde{u}\|_\infty \int_{[0,T]_{\mathbb{T}}} g^\sigma(t) \Delta t \\
& \leq \frac{M_1}{4} \int_{[0,T]_{\mathbb{T}}} |u^\Delta(t)|^2 \Delta t + C_6 |\bar{u}|^{2\alpha} + C_7 \left(\int_{[0,T]_{\mathbb{T}}} |u^\Delta(t)|^2 \Delta t \right)^{(\alpha+1)/2} \\
& \quad + C_8 \left(\int_{[0,T]_{\mathbb{T}}} |u^\Delta(t)|^2 \Delta t \right)^{1/2}
\end{aligned} \tag{4.3}$$

for all $u \in H_{\Delta, T}^1$, where $C_6 = (4M_2^2 C_5 / M_1) \left(\int_{[0,T]_{\mathbb{T}}} f^\sigma(t) dt \right)^2$, $C_7 = 2M_2(C_5)^{(\alpha+1)/2} \int_{[0,T]_{\mathbb{T}}} f^\sigma(t) \Delta t$, $C_8 = M_2(C_5)^{1/2} \int_{[0,T]_{\mathbb{T}}} g^\sigma(t) \Delta t$. Therefore, one has

$$\begin{aligned}
\varphi(u) &= \frac{1}{2} \int_{[0,T]_{\mathbb{T}}} e_w(t,0) |u^\Delta(t)|^2 \Delta t + \int_{[0,T]_{\mathbb{T}}} e_w(t,0) F(\sigma(t), u^\sigma(t)) \Delta t \\
&= \frac{1}{2} \int_{[0,T]_{\mathbb{T}}} e_w(t,0) |u^\Delta(t)|^2 \Delta t + \int_{[0,T]_{\mathbb{T}}} e_w(t,0) (F(\sigma(t), u^\sigma(t)) - F(\sigma(t), \bar{u})) \Delta t \\
& \quad + \int_{[0,T]_{\mathbb{T}}} e_w(t,0) F(\sigma(t), \bar{u}) \Delta t \\
&\geq \frac{1}{4} M_1 \int_{[0,T]_{\mathbb{T}}} |u^\Delta(t)|^2 \Delta t + |\bar{u}|^{2\alpha} \left(|\bar{u}|^{-2\alpha} \int_{[0,T]_{\mathbb{T}}} e_w(t,0) F(\sigma(t), \bar{u}) \Delta t - C_6 \right) \\
& \quad - C_7 \left(\int_{[0,T]_{\mathbb{T}}} |u^\Delta(t)|^2 \Delta t \right)^{(\alpha+1)/2} - C_8 \left(\int_{[0,T]_{\mathbb{T}}} |u^\Delta(t)|^2 \Delta t \right)^{1/2}
\end{aligned} \tag{4.4}$$

for all $u \in H^1_{\Delta,T}$. As $\|u\| \rightarrow \infty$ if and only if $(|\bar{u}|^2 + \int_0^T |u^\Delta(t)|^2 dt)^{1/2} \rightarrow \infty$, (4.4) and (ii) imply that

$$\varphi(u) \rightarrow +\infty \quad \text{as } \|u\| \rightarrow \infty. \tag{4.5}$$

By Lemma 3.8 and Theorem 3.4, φ has a minimum point on $H^1_{\Delta,T}$, which is a critical point of φ . From Theorem 3.3, problem (1.1) has at least one solution. \square

Example 4.2. Let $\mathbb{T} = \mathbb{R}, T = 2\pi, N = 3$. Consider the damped vibration problem on time-scale \mathbb{T}

$$\begin{aligned} \ddot{u}(t) + \cos t \dot{u}(t) &= \nabla F(t, u(t)), \quad \text{a.e. } t \in [0, 2\pi], \\ u(0) - u(2\pi) &= \dot{u}(0) - \dot{u}(2\pi) = 0, \end{aligned} \tag{4.6}$$

where $F(t, x) = (4/3 + t)|x|^{3/2}$.

Since, $F(t, x) = (4/3 + t)|x|^{3/2}, w(t) = \cos t, \nabla F(t, x) = (3/2)(4/3 + t)x|x|^{-1/2}, \alpha = 1/2$,

$$\begin{aligned} |\nabla F(t, x)| &\leq \frac{3}{2} \left(\frac{4}{3} + t \right) |x|^{1/2}, \\ |x|^{-2 \times 1/2} \int_0^{2\pi} e^{\sin t} \left(\frac{4}{3} + t \right) |x|^{3/2} dt &\geq |x|^{1/2} e^{-1} \int_0^{2\pi} \left(\frac{4}{3} + t \right) dt \\ &= \left(\frac{8}{3}\pi + 2\pi^2 \right) e^{-1} |x|^{1/2} \rightarrow +\infty \quad \text{as } |x| \rightarrow \infty, \end{aligned} \tag{4.7}$$

all conditions of Theorem 4.1 hold. According to Theorem 4.1, problem (4.6) has at least one solution. Moreover, 0 is not the solution of problem (4.6). Thus, problem (4.6) has at least one nontrivial solution.

Theorem 4.3. *Suppose that assumption (A) and the condition (i) of Theorem 4.1 hold. Assume that*

$$(iii) \quad |x|^{-2\alpha} \int_{[0,T]_{\mathbb{T}}} e_w(t, 0) F(\sigma(t), x) \Delta t \rightarrow -\infty \quad \text{as } |x| \rightarrow \infty.$$

Then problem (1.1) has at least one solution.

Firstly, we prove the following lemma.

Lemma 4.4. *Suppose that the conditions of Theorem 4.3 hold. Then φ satisfies P.S. condition.*

Proof. Let $\{u_n\} \subseteq H_{\Delta, T}^1$ be a P.S. sequence for φ , that is, $\{\varphi(u_n)\}$ is bounded and $\varphi'(u_n) \rightarrow 0$ as $n \rightarrow \infty$. It follows from (i), Theorem 2.23 and (4.2) that

$$\begin{aligned}
& \left| \int_{[0, T]_{\mathbb{T}}} e_w(t, 0) (F(\sigma(t), u_n^\sigma(t)) - F(\sigma(t), \bar{u}_n)) \Delta t \right| \\
& \leq \left| \int_{[0, T]_{\mathbb{T}}} e_w(t, 0) \left(\int_0^1 (\nabla F(\sigma(t), \bar{u}_n + s\tilde{u}_n^\sigma(t)), \tilde{u}_n^\sigma(t)) ds \right) \Delta t \right| \\
& \leq M_2 \int_{[0, T]_{\mathbb{T}}} \left(\int_0^1 f^\sigma(t) |\bar{u}_n + s\tilde{u}_n^\sigma(t)|^\alpha |\tilde{u}_n^\sigma(t)| ds \right) \Delta t + M_2 \int_{[0, T]_{\mathbb{T}}} \left(\int_0^1 g^\sigma(t) |\tilde{u}_n^\sigma(t)| ds \right) \Delta t \\
& \leq 2M_2 (|\bar{u}_n|^\alpha + \|\tilde{u}_n\|_\infty^\alpha) \|\tilde{u}_n\|_\infty \int_{[0, T]_{\mathbb{T}}} f^\sigma(t) \Delta t + M_2 \|\tilde{u}_n\|_\infty \int_{[0, T]_{\mathbb{T}}} g^\sigma(t) \Delta t \\
& \leq \frac{M_1}{4C_5} \|\tilde{u}_n\|_\infty^2 + \frac{4M_2^2 C_5}{M_1} |\bar{u}_n|^{2\alpha} \left(\int_{[0, T]_{\mathbb{T}}} f^\sigma(t) \Delta t \right)^2 \\
& \quad + 2M_2 \|\tilde{u}_n\|_\infty^{\alpha+1} \int_{[0, T]_{\mathbb{T}}} f^\sigma(t) \Delta t + M_2 \|\tilde{u}_n\|_\infty \int_{[0, T]_{\mathbb{T}}} g^\sigma(t) \Delta t \\
& \leq \frac{M_1}{4} \int_{[0, T]_{\mathbb{T}}} |u_n^\Delta(t)|^2 \Delta t + C_6 |\bar{u}_n|^{2\alpha} \\
& \quad + C_7 \left(\int_{[0, T]_{\mathbb{T}}} |u_n^\Delta(t)|^2 \Delta t \right)^{(\alpha+1)/2} + C_8 \left(\int_{[0, T]_{\mathbb{T}}} |u_n^\Delta(t)|^2 \Delta t \right)^{1/2}
\end{aligned} \tag{4.8}$$

for all n . By (4.8) and (i), one has

$$\begin{aligned}
\|\tilde{u}_n\| & \geq \langle \varphi'(u_n), \tilde{u}_n \rangle \\
& = \int_{[0, T]_{\mathbb{T}}} e_w(t, 0) |u_n^\Delta(t)|^2 \Delta t + \int_{[0, T]_{\mathbb{T}}} e_w(t, 0) \nabla F(\sigma(t), u_n^\sigma(t), \tilde{u}_n(t)) \Delta t \\
& \geq \frac{3M_1}{4} \int_{[0, T]_{\mathbb{T}}} |u_n^\Delta(t)|^2 \Delta t - C_6 |\bar{u}_n|^{2\alpha} - C_7 \left(\int_{[0, T]_{\mathbb{T}}} |u_n^\Delta(t)|^2 \Delta t \right)^{(\alpha+1)/2} \\
& \quad - C_8 \left(\int_{[0, T]_{\mathbb{T}}} |u_n^\Delta(t)|^2 \Delta t \right)^{1/2}
\end{aligned} \tag{4.9}$$

for all large n . It follows from (3.5) and (4.2) that

$$M_1 \int_{[0, T]_{\mathbb{T}}} |u_n^\Delta(t)|^2 \Delta t \leq \|\tilde{u}_n\|^2 \leq M_2 (1 + TC_5) \int_{[0, T]_{\mathbb{T}}} |u_n^\Delta(t)|^2 \Delta t. \tag{4.10}$$

The inequalities (4.9) and (4.10) imply that

$$C_9 |\bar{u}_n|^\alpha \geq \left(\int_{[0,T]_{\mathbb{T}}} |u_n^\Delta(t)|^2 \Delta t \right)^{1/2} - C_{10} \tag{4.11}$$

for all large n and some positive constants C_9 and C_{10} . Similar to the proof of Theorem 4.1, one has

$$\begin{aligned} & \left| \int_{[0,T]_{\mathbb{T}}} e_w(t,0) (F(\sigma(t), u_n^\sigma(t)) - F(\sigma(t), \bar{u}_n)) \Delta t \right| \\ & \leq \frac{M_1}{4} \int_{[0,T]_{\mathbb{T}}} |u_n^\Delta(t)|^2 dt + C_6 |\bar{u}_n|^{2\alpha} \\ & \quad + C_7 \left(\int_{[0,T]_{\mathbb{T}}} |u_n^\Delta(t)|^2 \Delta t \right)^{(\alpha+1)/2} + C_8 \left(\int_{[0,T]_{\mathbb{T}}} |u_n^\Delta(t)|^2 \Delta t \right)^{1/2} \end{aligned} \tag{4.12}$$

for all n . By the boundedness of $\{\varphi(u_n)\}$, (4.11) and (4.12), there exists constant C_{11} such that

$$\begin{aligned} C_{11} & \leq \varphi(u_n) \\ & = \frac{1}{2} \int_{[0,T]_{\mathbb{T}}} e_w(t,0) |u_n^\Delta(t)|^2 dt + \int_{[0,T]_{\mathbb{T}}} e_w(t,0) (F(\sigma(t), u_n^\sigma(t)) - F(\sigma(t), \bar{u}_n)) \Delta t \\ & \quad + \int_{[0,T]_{\mathbb{T}}} e_w(t,0) F(\sigma(t), \bar{u}_n) \Delta t \\ & \leq \frac{3}{4} M_2 \int_{[0,T]_{\mathbb{T}}} |u_n^\Delta(t)|^2 \Delta t + C_6 |\bar{u}_n|^{2\alpha} + \int_{[0,T]_{\mathbb{T}}} e_w(t,0) F(\sigma(t), \bar{u}_n) \Delta t \\ & \quad + C_7 \left(\int_{[0,T]_{\mathbb{T}}} |u_n^\Delta(t)|^2 \Delta t \right)^{(\alpha+1)/2} + C_8 \left(\int_{[0,T]_{\mathbb{T}}} |u_n^\Delta(t)|^2 \Delta t \right)^{1/2} \\ & \leq |\bar{u}_n|^{2\alpha} \left(|\bar{u}_n|^{-2\alpha} \int_{[0,T]_{\mathbb{T}}} e_w(t,0) F(\sigma(t), \bar{u}_n) \Delta t + C_{12} \right) \end{aligned} \tag{4.13}$$

for all large n and some constant C_{12} . It follows from (4.13) and (iii) that $\{|\bar{u}_n|\}$ is bounded. Hence $\{u_n\}$ is bounded in $H^1_{\Delta,T}$ by (4.10) and (4.11). Therefore, there exists a subsequence of $\{u_n\}$ (for simplicity denoted again by $\{u_n\}$) such that

$$u_n \rightharpoonup u \quad \text{in } H^1_{\Delta,T}. \tag{4.14}$$

By Theorem 2.25, one has

$$u_n \longrightarrow u \quad \text{in } C([0, T]_{\mathbb{T}}, \mathbb{R}^N). \tag{4.15}$$

On the other hand, one has

$$\begin{aligned} & \langle \varphi'(u_n) - \varphi'(u), u_n - u \rangle \\ &= \int_{[0,T]_{\mathbb{T}}} e_w(t,0) \left| u_n^\Delta(t) - u^\Delta(t) \right|^2 \Delta t \\ & \quad + \int_{[0,T]_{\mathbb{T}}} e_w(t,0) (\nabla F(\sigma(t), u_n^\sigma(t)) - \nabla F(\sigma(t), u^\sigma(t)), u_n^\sigma(t) - u^\sigma(t)) \Delta t. \end{aligned} \quad (4.16)$$

From (4.14), (4.15), (4.16), and (A), it follows that $u_n \rightarrow u$ in $H_{\Delta,T}^1$. Thus, φ satisfies P.S. condition. \square

Now, we prove Theorem 4.3.

Proof. Let W be the subspace of $H_{\Delta,T}^1$ given by

$$W = \left\{ u \in H_{\Delta,T}^1 : \int_{[0,T]_{\mathbb{T}}} u(t) \Delta t = 0 \right\}, \quad (4.17)$$

then, $H_{\Delta,T}^1 = \mathbb{R}^N \oplus W$. We show that

$$\varphi(u) \rightarrow +\infty \quad \text{as } u \in W, \|u\| \rightarrow \infty. \quad (4.18)$$

Indeed, for $u \in W$, then $\bar{u} = 0$, similar to the proof of Theorem 4.1, one has

$$\begin{aligned} & \left| \int_{[0,T]_{\mathbb{T}}} e_w(t,0) (F(\sigma(t), u^\sigma(t)) - F(\sigma(t), 0)) \Delta t \right| \\ & \leq \frac{M_1}{4} \int_{[0,T]_{\mathbb{T}}} |u^\Delta(t)|^2 \Delta t + C_7 \left(\int_{[0,T]_{\mathbb{T}}} |u^\Delta(t)|^2 \Delta t \right)^{(\alpha+1)/2} + C_8 \left(\int_{[0,T]_{\mathbb{T}}} |u^\Delta(t)|^2 \Delta t \right)^{1/2}. \end{aligned} \quad (4.19)$$

It follows from (4.19) that

$$\begin{aligned} \varphi(u) &= \frac{1}{2} \int_{[0,T]_{\mathbb{T}}} e_w(t,0) |u^\Delta(t)|^2 \Delta t + \int_{[0,T]_{\mathbb{T}}} e_w(t,0) (F(\sigma(t), u^\sigma(t)) - F(\sigma(t), 0)) \Delta t \\ & \quad + \int_{[0,T]_{\mathbb{T}}} e_w(t,0) F(\sigma(t), 0) \Delta t \\ & \geq \frac{M_1}{4} \int_{[0,T]_{\mathbb{T}}} |u^\Delta(t)|^2 \Delta t - C_7 \left(\int_{[0,T]_{\mathbb{T}}} |u^\Delta(t)|^2 \Delta t \right)^{(\alpha+1)/2} - C_8 \left(\int_{[0,T]_{\mathbb{T}}} |u^\Delta(t)|^2 \Delta t \right)^{1/2} \\ & \quad + \int_{[0,T]_{\mathbb{T}}} e_w(t,0) F(\sigma(t), 0) \Delta t \end{aligned} \quad (4.20)$$

for all $u \in W$. By Theorem 2.23 and Theorem 3.1, one has

$$\|u\| \rightarrow \infty \iff \|u^\Delta\|_{L^2} \rightarrow \infty \tag{4.21}$$

on W . Hence (4.18) follows from (4.20).

On the other hand, by (iii), one has

$$\begin{aligned} \varphi(u) &= \int_{[0,T]_{\mathbb{T}}} e_w(t,0)F(\sigma(t),u)\Delta t \\ &\leq |u|^{2\alpha} \left(|u|^{-2\alpha} \int_{[0,T]_{\mathbb{T}}} e_w(t,0)F(\sigma(t),u)\Delta t \right) \rightarrow -\infty \end{aligned} \tag{4.22}$$

as $u \in \mathbb{R}^N$ and $|u| \rightarrow \infty$. By Theorem 3.3, Lemmas 3.9 and 4.4, problem (1.1) has at least one solution. \square

Example 4.5. Let $\mathbb{T} = \mathbb{Z}, T = 20, N = 5$. Consider the damped vibration problem on time-scale \mathbb{T}

$$\begin{aligned} \Delta^2(t) + w(t)\Delta u(t+1) &= \nabla F(t+1, u(t+1)), \quad t \in [0, 19] \cap \mathbb{Z}, \\ u(0) - u(20) &= \Delta u(0) - \Delta u(20) = 0, \end{aligned} \tag{4.23}$$

where $F(t, x) = -|x|^{5/3} + ((1, 1, 2, 1, 0), x)$ and

$$w(t) = \begin{cases} -\frac{1}{2}, & t \in [0, 18] \cap \mathbb{Z}, \\ 2^{18} - 1, & t = 19. \end{cases} \tag{4.24}$$

Since, $F(t, x) = -|x|^{5/3} + ((1, 1, 2, 1, 0), x)$, $\nabla F(t, x) = -(5/3)x|x|^{-1/3} + (1, 1, 2, 1, 0)$, $\alpha = 2/3$, $e_w(t, 0) = \prod_{s=0}^{t-1} (1 + w(s))$, $e_w(20, 0) = 1$,

$$\begin{aligned} |\nabla F(t, x)| &\leq \frac{5}{3}|x|^{2/3} + \sqrt{7}, \\ |x|^{-2 \times (2/3)} \int_{[0,T]_{\mathbb{T}}} e_w(t,0)F(\sigma(t),x)\Delta t \\ &= |x|^{-4/3} \left(-|x|^{5/3} + ((1, 1, 2, 1, 0), x) \right) \int_{[0,T]_{\mathbb{T}}} e_w(t,0)\Delta t \\ &\leq -|x|^{1/3} \int_{[0,T]_{\mathbb{T}}} e_w(t,0)\Delta t + \sqrt{7}|x|^{-1/3} \int_{[0,T]_{\mathbb{T}}} e_w(t,0)\Delta t \rightarrow -\infty \quad \text{as } |x| \rightarrow \infty, \end{aligned} \tag{4.25}$$

all conditions of Theorem 4.3 hold. According to Theorem 4.3, problem (4.23) has at least one solution. Moreover, 0 is not the solution of problem (4.23). Thus, problem (4.23) has at least one nontrivial solution.

Theorem 4.6. *Suppose that assumption (A) and the following condition are satisfied.*

(iv) $F(t, \cdot)$ is convex for Δ -a.e. $t \in [0, T]_{\mathbb{T}}$ and that

$$\int_{[0, T]_{\mathbb{T}}} e_w(t, 0) F(\sigma(t), x) \Delta t \longrightarrow +\infty \quad \text{as } |x| \longrightarrow \infty. \quad (4.26)$$

Then problem (1.1) has at least one solution which minimizes the function φ .

Proof. By assumption, the function $G : \mathbb{R}^N \rightarrow \mathbb{R}$ defined by

$$G(x) = \int_{[0, T]_{\mathbb{T}}} e_w(t, 0) F(\sigma(t), x) \Delta t \quad (4.27)$$

has a minimum at some point \bar{x} for which

$$\int_{[0, T]_{\mathbb{T}}} e_w(t, 0) \nabla F(\sigma(t), \bar{x}) \Delta t = 0. \quad (4.28)$$

Let $\{u_k\}$ be a minimizing sequence for φ . From Lemma 3.10 and (4.28), one has

$$\begin{aligned} \varphi(u_k) &= \frac{1}{2} \int_{[0, T]_{\mathbb{T}}} e_w(t, 0) \left| u_k^\Delta(t) \right|^2 \Delta t + \int_{[0, T]_{\mathbb{T}}} e_w(t, 0) (F(\sigma(t), u_k^\sigma(t)) - F(\sigma(t), \bar{x})) \Delta t \\ &\quad + \int_{[0, T]_{\mathbb{T}}} e_w(t, 0) F(\sigma(t), \bar{x}) \Delta t \\ &\geq \frac{1}{2} \int_{[0, T]_{\mathbb{T}}} e_w(t, 0) \left| u_k^\Delta(t) \right|^2 \Delta t + \int_{[0, T]_{\mathbb{T}}} e_w(t, 0) F(\sigma(t), \bar{x}) \Delta t \\ &\quad + \int_{[0, T]_{\mathbb{T}}} e_w(t, 0) (\nabla F(\sigma(t), \bar{x}), u_k^\sigma(t) - \bar{x}) \Delta t \\ &= \frac{1}{2} \int_{[0, T]_{\mathbb{T}}} e_w(t, 0) \left| u_k^\Delta(t) \right|^2 \Delta t + \int_{[0, T]_{\mathbb{T}}} e_w(t, 0) F(\sigma(t), \bar{x}) \Delta t \\ &\quad + \int_{[0, T]_{\mathbb{T}}} e_w(t, 0) (\nabla F(\sigma(t), \bar{x}), \tilde{u}_k^\sigma(t)) \Delta t, \end{aligned} \quad (4.29)$$

where $\tilde{u}_k(t) = u_k(t) - \bar{u}_k$, $\bar{u}_k = (1/T) \int_{[0,T]_{\mathbb{T}}} u_k(t) \Delta t$. By (4.29), (A) and Theorem 2.23, we obtain

$$\begin{aligned} \varphi(u_k) &\geq \frac{1}{2} \int_{[0,T]_{\mathbb{T}}} e_w(t,0) \left| u_k^\Delta(t) \right|^2 \Delta t + \int_{[0,T]_{\mathbb{T}}} e_w(t,0) F(\sigma(t), \bar{x}) \Delta t \\ &\quad - M_2 \left(\int_{[0,T]_{\mathbb{T}}} |\nabla F(\sigma(t), \bar{x})| \Delta t \right) \|\tilde{u}_k\|_\infty \\ &\geq \frac{1}{2} M_1 \int_{[0,T]_{\mathbb{T}}} e_w(t,0) \left| u_k^\Delta(t) \right|^2 \Delta t - C_{13} - C_{14} \left(\int_{[0,T]_{\mathbb{T}}} \left| u_k^\Delta(t) \right|^2 \Delta t \right)^{1/2} \end{aligned} \tag{4.30}$$

for some positive constants C_{13} and C_{14} . Thus, by (4.30), there exists $C_{15} > 0$ such that

$$\int_{[0,T]_{\mathbb{T}}} \left| u_k^\Delta(t) \right|^2 \Delta t \leq C_{15}. \tag{4.31}$$

Theorem 2.23 and (4.31) imply that there exists $C_{16} > 0$ such that

$$\|\tilde{u}_k\|_\infty \leq C_{16}. \tag{4.32}$$

By (iv), one has

$$\begin{aligned} F\left(\sigma(t), \frac{\bar{u}_k}{2}\right) &= F\left(\sigma(t), \frac{u_k^\sigma(t) - \tilde{u}_k^\sigma(t)}{2}\right) \\ &\leq \frac{1}{2} F(\sigma(t), u_k^\sigma(t)) + \frac{1}{2} F(\sigma(t), -\tilde{u}_k^\sigma(t)) \end{aligned} \tag{4.33}$$

for Δ -a.e. $t \in [0, T]_{\mathbb{T}}$ and all $k \in \mathbb{N}$. It follows from (3.9) and (4.33) that

$$\begin{aligned} \varphi(u_k) &\geq \frac{1}{2} \int_{[0,T]_{\mathbb{T}}} e_w(t,0) \left| u_k^\Delta(t) \right|^2 \Delta t + 2 \int_{[0,T]_{\mathbb{T}}} e_w(t,0) F\left(\sigma(t), \frac{\bar{u}_k}{2}\right) \Delta t \\ &\quad - \int_{[0,T]_{\mathbb{T}}} e_w(t,0) F(\sigma(t), -\tilde{u}_k^\sigma(t)) \Delta t. \end{aligned} \tag{4.34}$$

Combining (4.32) and (4.34), there exists $C_{17} > 0$ such that

$$\varphi(u_k) \geq 2 \int_{[0,T]_{\mathbb{T}}} e_w(t,0) F\left(\sigma(t), \frac{\bar{u}_k}{2}\right) \Delta t - C_{17}. \tag{4.35}$$

Therefore, by (4.35) and (iv), $\{\bar{u}_k\}$ is bounded. Hence $\{u_k\}$ is bounded in $H_{\Delta,T}^1$ by Theorem 2.23 and (4.31). By Lemma 3.8 and Theorem 3.4, φ has a minimum point on $H_{\Delta,T}^1$, which is a critical point of φ . Hence, problem (1.1) has at least one solution which minimizes the function φ . \square

Example 4.7. Let $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$, $\mathbb{T} = \{2^k : k \in \mathbb{N}_0\}$, $T = 64$, $N = 1$. Consider the damped vibration problem on time-scale \mathbb{T}

$$\begin{aligned} u^{\Delta^2}(t) + w(t)u^{\Delta}(2t) &= \nabla F(2t, u(2t)), \quad t \in \{1, 2, 4, 8, 16, 32\}, \\ u(0) - u(64) &= u^{\Delta}(0) - u^{\Delta}(64) = 0, \end{aligned} \quad (4.36)$$

where $F(t, x) = x^2 + 2x$ and

$$w(t) = \begin{cases} -\frac{1}{2t}, & t \in \{1, 2, 4, 8, 16\}, \\ \frac{31}{32}, & t = 32. \end{cases} \quad (4.37)$$

Since, $F(t, x) = x^2 + 2x$, $e_w(t, 0) = \prod_{s \in \mathbb{T} \cap (0, t)} (1 + sw(s))$, $e_w(64, 0) = 1$, all conditions of Theorem 4.6 hold. According to Theorem 4.6, problem (4.36) has at least one solution. Moreover, 0 is not the solution of problem (4.36). Thus, problem (4.36) has at least one nontrivial solution.

5. Conclusion

In this paper, we present a new approach via variational methods and critical point theory to obtain the existence of solutions for a class of damped vibration problems on time-scales. Three existence results are obtained. Three examples are presented to illustrate the feasibility and effectiveness of our results.

Acknowledgment

This work is supported by the National Natural Sciences Foundation of People's Republic of China under Grant 10971183.

References

- [1] B. Aulbach and S. Hilger, "Linear dynamic processes with inhomogeneous time scale," in *Nonlinear Dynamics and Quantum Dynamical Systems (Gaussig, 1990)*, vol. 59 of *Math. Res.*, pp. 9–20, Akademie, Berlin, Germany, 1990.
- [2] V. Lakshmikantham, S. Sivasundaram, and B. Kaymakcalan, *Dynamic Systems on Measure Chains*, vol. 370 of *Mathematics and Its Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1996.
- [3] M. Bohner and A. Peterson, *Dynamic Equations on Time Scales: An Introduction with Applications*, Birkhäuser, Boston, Mass, USA, 2001.
- [4] R. P. Agarwal and D. O'Regan, *Infinite Interval Problems for Differential Difference and Integral Equations*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 2001.
- [5] M. Bohner and A. Peterson, Eds., *Advances in Dynamic Equations on Time Scales*, Birkhäuser, Boston, Mass, USA, 2003.
- [6] R. P. Agarwal, M. Bohner, and P. Řehák, "Half-linear dynamic equations," in *Nonlinear Analysis and Applications: to V. Lakshmikantham on His 80th Birthday. Vol. 1, 2*, pp. 1–57, Kluwer Academic Publishers, Dordrecht, The Netherlands, 2003.

- [7] R. P. Agarwal, M. Bohner, and W.-T. Li, *Nonoscillation and Oscillation: Theory for Functional Differential Equations*, vol. 267 of *Monographs and Textbooks in Pure and Applied Mathematics*, Marcel Dekker, New York, NY, USA, 2004.
- [8] R. P. Agarwal and M. Bohner, "Basic calculus on time scales and some of its applications," *Results in Mathematics*, vol. 35, no. 1-2, pp. 3–22, 1999.
- [9] D. R. Anderson, "Eigenvalue intervals for a second-order mixed-conditions problem on time scales," *International Journal of Differential Equations*, vol. 7, pp. 97–104, 2002.
- [10] G. Sh. Guseinov, "Integration on time scales," *Journal of Mathematical Analysis and Applications*, vol. 285, no. 1, pp. 107–127, 2003.
- [11] M. Bohner and G. Sh. Guseinov, "Improper integrals on time scales," *Dynamic Systems and Applications*, vol. 12, no. 1-2, pp. 45–65, 2003.
- [12] E. R. Kaufmann and Y. N. Raffoul, "Periodic solutions for a neutral nonlinear dynamical equation on a time scale," *Journal of Mathematical Analysis and Applications*, vol. 319, no. 1, pp. 315–325, 2006.
- [13] R. P. Agarwal, V. Otero-Espinar, K. Perera, and D. R. Vivero, "Basic properties of Sobolev's spaces on time scales," *Advances in Difference Equations*, vol. 2006, Article ID 38121, 14 pages, 2006.
- [14] H. Zhang and Y. Li, "Existence of positive periodic solutions for functional differential equations with impulse effects on time scales," *Communications in Nonlinear Science and Numerical Simulation*, vol. 14, no. 1, pp. 19–26, 2009.
- [15] V. Otero-Espinar and D. R. Vivero, "Existence and approximation of extremal solutions to first-order infinite systems of functional dynamic equations," *Journal of Mathematical Analysis and Applications*, vol. 339, no. 1, pp. 590–597, 2008.
- [16] R. P. Agarwal, V. Otero-Espinar, K. Perera, and D. R. Vivero, "Multiple positive solutions of singular Dirichlet problems on time scales via variational methods," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 67, no. 2, pp. 368–381, 2007.
- [17] H.-R. Sun and W.-T. Li, "Existence theory for positive solutions to one-dimensional p -Laplacian boundary value problems on time scales," *Journal of Differential Equations*, vol. 240, no. 2, pp. 217–248, 2007.
- [18] Z.-C. Hao, T.-J. Xiao, and J. Liang, "Existence of positive solutions for singular boundary value problem on time scales," *Journal of Mathematical Analysis and Applications*, vol. 325, no. 1, pp. 517–528, 2007.
- [19] L. Jiang and Z. Zhou, "Existence of weak solutions of two-point boundary value problems for second-order dynamic equations on time scales," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 69, no. 4, pp. 1376–1388, 2008.
- [20] R. P. Agarwal, M. Bohner, and P. J. Y. Wong, "Sturm-Liouville eigenvalue problems on time scales," *Applied Mathematics and Computation*, vol. 99, no. 2-3, pp. 153–166, 1999.
- [21] B. P. Rynne, " L^2 spaces and boundary value problems on time-scales," *Journal of Mathematical Analysis and Applications*, vol. 328, no. 2, pp. 1217–1236, 2007.
- [22] F. A. Davidson and B. P. Rynne, "Eigenfunction expansions in L^2 spaces for boundary value problems on time-scales," *Journal of Mathematical Analysis and Applications*, vol. 335, no. 2, pp. 1038–1051, 2007.
- [23] J. Zhou and Y. Li, "Sobolev's spaces on time scales and its applications to a class of second order Hamiltonian systems on time scales," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 73, no. 5, pp. 1375–1388, 2010.
- [24] J. Mawhin and M. Willem, *Critical Point Theory and Hamiltonian Systems*, vol. 74 of *Applied Mathematical Sciences*, Springer, New York, NY, USA, 1989.