

RESEARCH ARTICLE

Open Access

A study of nonlocal problems of impulsive integrodifferential equations with measure of noncompactness

Bashir Ahmad^{1*}, Kandasamy Malar² and Kulandhaivel Karthikeyan³

*Correspondence:

bashirahmad_qau@yahoo.com

¹Department of Mathematics,
Faculty of Science, King Abdulaziz
University, P.O. Box 80203, Jeddah,
21589, Saudi Arabia
Full list of author information is
available at the end of the article

Abstract

In this paper, we show the existence of mild solutions to a nonlocal problem of impulsive integrodifferential equations via a measure of noncompactness in a Banach space. Our work is based on a new fixed point theorem and it generalizes some existing results on the topic in the sense that we do not require the semigroup and nonlinearity involved in the problem to be compact.

Keywords: integrodifferential equation; impulse; semigroup; mild solution; fixed point; measure of noncompactness

1 Introduction

In this paper, we discuss the existence of solutions for the following nonlocal problem of integrodifferential equations:

$$\begin{aligned}\frac{du(t)}{dt} &= Au(t) + f(t, u(t), Gu(t)), \quad t \in [0, K], t \neq t_i, \\ u(0) &= u_0 + g(u), \\ \Delta u(t_i) &= I_i(u(t_i)), \quad i = 1, 2, 3, \dots, p,\end{aligned}\tag{1.1}$$

where A generates a C_0 -semigroup $T(t)$, $t \geq 0$, in a Banach space X , $f: [0, K] \times X \times X \rightarrow X$, $Gu(t) = \int_0^t H(t, s)u(s) ds$, $H \in C[D, R^+]$, $D = \{(t, s) \in R^2 : 0 \leq s \leq t \leq K\}$, $0 < t_1 < t_2 < t_3 < \dots < t_p < K$ with $t_{p+1} = K$, $g: X \rightarrow X$, $u_0 \in X$, and $\Delta u(t_i) = u(t_i^+) - u(t_i^-)$ with $u(t_i^+)$, $u(t_i^-)$ representing the right and left limit of u at t_i , respectively. Here $PC([0, K]; X) = \{u: [0, K] \rightarrow X \text{ is continuous at } t \neq t_i, \text{ left continuous at } t = t_i, \text{ and the right-hand limit } u(t_i^+) \text{ exists for } i = 1, 2, \dots, p\}$. Notice that the set $PC([0, K]; X)$ equipped with the norm $\|u\| = \sup\{\|u(t)\| : t \in [0, K]\}$ is a Banach space.

Integrodifferential equations arise in the mathematical modeling of several natural phenomena and various investigations led to the exploration of their different aspects. The theory of semigroups of bounded linear operators is closely related to the solution of differential and integrodifferential equations in Banach spaces. In recent years, this theory has been applied to a large class of nonlinear differential equations in Banach spaces. Based on the method of semigroups, the existence and uniqueness of mild, strong and classical solutions of semilinear evolution equations were discussed by Pazy [1]. For further details, see [2–5] and the references cited therein.

The theory of impulsive differential equations is an important branch of differential equations and has an extensive physical background. Impulsive differential equations help modeling many physical systems whose states are subject to abrupt changes at certain moments. Examples include population biology, the diffusion of chemicals, the spread of heat, the radiation of electromagnetic waves, *etc.* [6–9]. Dynamical systems with impulsive effects have been an object of intensive investigations [10–12]. The study of semilinear nonlocal initial problem was initiated by Byszewski [13, 14] and the importance of the problem lies in the fact that it is more general and yields better effect than the classical initial conditions. Therefore, it has been extensively studied under various conditions on the operator A and the nonlinearity f by several authors [15–17].

Byszewski and Lakshmikantham [18] showed the existence and uniqueness of mild solutions and classical solutions when f and g in (1.1) satisfy Lipschitz type conditions. Ntouyas and Tsamotas [19, 20] studied the case of compact-valued f and T . Zhu *et al.* [21] discussed the existence of mild solutions for abstract semilinear evolution equations in Banach spaces. In [11], the author discussed the existence and uniqueness of mild and classical solutions for the impulsive semilinear differential evolution equation. In [22], Agarwal *et al.* studied the existence and dimension of the set of mild solutions to semilinear fractional differential inclusions. Lizama and Pozo [23] investigated the existence of mild solutions for semilinear integrodifferential equation with nonlocal initial conditions by using Hausdorff measure of noncompactness via a fixed-point. In a recent paper [24], the authors studied the existence of mild solutions to an impulsive differential equation with nonlocal conditions by applying Darbo-Sadovskii's fixed point theorem. For some more recent results and details, see [25–37].

Motivated by [23], in this paper we aim to establish some existence results for mild solutions of (1.1) without demanding the compactness condition on T and f . In this scenario, our work extends and improves some results obtained in [38, 39]. In Section 2, we recall some definitions and facts about C_0 semigroup T and the measure of noncompactness, while Section 3 deals with the existence of mild solutions for (1.1).

2 Preliminary result

Let $L^1([0, K]; X)$ denote the space of X -valued Bochner functions on $[0, K]$ with the norm defined by $\|u\| = \int_0^K \|u(s)\| ds$. A C_0 -semigroup $T(t)$ is said to be compact if $T(t)$ is compact for any $t > 0$. If the semigroup $T(t)$ is compact, then $t \rightarrow T(t)u$ are equicontinuous at all $t > 0$ with respect to u in all bounded subset of X , that is, the semigroup $T(t)$ is equicontinuous.

In this paper, α denotes the Hausdorff measure of noncompactness on both X and $PC([0, K]; X)$. The following lemma describes some properties of the Hausdorff measure of noncompactness.

Lemma 2.1 [40] *Let $\beta_{\mathbb{Y}}$ and $\beta_{\mathbb{Z}}$ denote Hausdorff measures of noncompactness on the real Banach spaces \mathbb{Y} and \mathbb{Z} , respectively, and $B, C \subseteq \mathbb{Y}$ be bounded. Then*

1. B is pre-compact if and only if $\beta_{\mathbb{X}}(B) = 0$;
2. $\beta_{\mathbb{Y}}(B) = \beta_{\mathbb{Y}}(\overline{B}) = \beta_{\mathbb{Y}}(\text{conv } B)$, where \overline{B} and $\text{conv } B$ mean the closure and convex hull of B , respectively;
3. $\beta_{\mathbb{Y}}(B) \leq \beta_{\mathbb{Y}}(C)$, where $B \subseteq C$;
4. $\beta_{\mathbb{Y}}(B + C) \leq \beta_{\mathbb{Y}}(B) + \beta_{\mathbb{Y}}(C)$, where $B + C = \{x + y : x \in B, y \in C\}$;
5. $\beta_{\mathbb{Y}}(B \cup C) \leq \max\{\beta_{\mathbb{Y}}(B), \beta_{\mathbb{Y}}(C)\}$;

6. $\beta_{\mathbb{Y}}(\lambda B) \leq |\lambda| \beta_{\mathbb{Y}}(B)$ for any $\lambda \in \mathbb{R}$;
7. If the map $Q : D(Q) \subseteq \mathbb{Y} \rightarrow \mathbb{Z}$ is Lipschitz continuous with constant k , then $\beta_{\mathbb{Z}}(QB) \leq k \beta_{\mathbb{Y}}(B)$ for any bounded subset $B \subseteq D(Q)$;
8. $\beta_{\mathbb{Y}}(B) = \inf\{d_{\mathbb{Y}}(B, C); C \subseteq \mathbb{Y} \text{ is precompact}\} = \inf\{d_{\mathbb{Y}}(B, C); C \subseteq \mathbb{Y} \text{ is finite valued}\}$, where $d_{\mathbb{Y}}(B, C)$ means the nonsymmetric (or symmetric) Hausdorff distance between B and C in \mathbb{Y} ;
9. If $\{W_n\}_{n=1}^{+\infty}$ is a decreasing sequence of bounded closed nonempty subsets of \mathbb{Y} and $\lim_{n \rightarrow \infty} \beta_{\mathbb{Y}}(W_n) = 0$, then $\bigcap_{n=1}^{+\infty} W_n$ is nonempty and compact in \mathbb{Y} .

The map $Q : W \subseteq \mathbb{Y} \rightarrow \mathbb{Y}$ is said to be a $\beta_{\mathbb{Y}}$ -contraction if there exists a constant $0 < k < 1$ such that $\beta_{\mathbb{Y}}(Q(B)) \leq k \beta_{\mathbb{Y}}(B)$ for any bounded closed subset $B \subseteq W$, where \mathbb{Y} is a Banach space.

In the sequel, we need the following known results.

Lemma 2.2 [40] *If $W \subseteq PC([0, K]; X)$ is bounded, then $\alpha(W(t)) \leq \alpha(W)$ for all $t \in [0, K]$, where $W(t) = \{u(t) : u \in W\} \subseteq X$. Furthermore, if W is equicontinuous on $[0, K]$, then $\alpha(W(t))$ is continuous on $[0, K]$, and $\alpha(W) = \sup\{\alpha(W(t)) : t \in [0, K]\}$.*

Lemma 2.3 [41] *If $\{u_n\}_{n=1}^{\infty} \subset L^1(0, K; X)$ is uniformly integrable, then $\alpha(\{u_n(t)\}_{n=1}^{\infty})$ is measurable and*

$$\alpha\left(\left\{\int_0^t u_n(s) ds\right\}_{n=1}^{\infty}\right) \leq 2 \int_0^t \alpha\{u_n(s)\}_{n=1}^{\infty} ds.$$

Lemma 2.4 [39] *If the semigroup $T(t)$ is equicontinuous and there exists $\eta \in L^1(0, K; R^+)$, then the set*

$$\left\{t \rightarrow \int_0^t T(t-s)u(s) ds; u \in L^1(0, K; R^+), \|u(s)\| \leq \eta(s), \text{ for a.e } s \in [0, K]\right\}$$

is equicontinuous on $[0, K]$.

Lemma 2.5 [42] *If W is bounded, then for each $\varepsilon > 0$, there is a sequence $\{u_n\}_{n=1}^{\infty} \subseteq W$ such that $\alpha(W) \leq 2\alpha(\{u_n\}_{n=1}^{\infty}) + \varepsilon$.*

Lemma 2.6 [43] *Suppose that $0 < \varepsilon < 1$, $h > 0$ and let*

$$S = \varepsilon^n + C_n^1 \varepsilon^{n-1} h + C_n^2 \varepsilon^{n-2} \frac{h^2}{2!} + \dots + \frac{h^n}{n!}, \quad n \in N.$$

Then $S = o(1/n^s)$ ($n \rightarrow +\infty$), where $s > 1$ is an arbitrary real number and C_n^1, C_n^2, \dots are binomial coefficients [44].

Lemma 2.7 ([45] Fixed point theorem) *Let Q be a closed and convex subset of a real Banach space X , let $A : Q \rightarrow Q$ be a continuous operator and $A(Q)$ be bounded. For each bounded subset $B \subset Q$, set*

$$A^1(B) = A(B), \quad A^n(B) = A(\bar{co}(A^{n-1}(B))), \quad n = 2, 3, \dots$$

If there exist a constant $0 \leq k \leq 1$ and a positive integer n_0 such that for each bounded subset $B \subset Q$, $\alpha(A^{n_0}(B)) \leq k\alpha(B)$, then A has a fixed point in Q .

Definition 2.8 A function $u : [0, K] \rightarrow X$ is called a mild solution of system (1.1) if $u \in PC([0, K] : X)$ and satisfies the following equation:

$$u(t) = \begin{cases} T(t)[u_0 - g(u)] + \int_0^t T(t-s)f(s, u(s), Gu(s)) ds, & t \in [0, t_1]; \\ T(t - t_1)(u(t_1^-) + I_1(u(t_1^-))) + \int_{t_1}^t T(t-s)f(s, u(s), Gu(s)) ds, & t \in (t_1, t_2]; \\ \vdots \\ T(t - t_p)(u(t_p^-) + I_p(u(t_p^-))) + \int_{t_p}^t T(t-s)f(s, u(s), Gu(s)) ds, & t \in (t_p, K]. \end{cases}$$

3 Existence result

In this section, we show the existence of solutions for problem (1.1) by applying Lemma 2.7. For some real constants r and w , we define

$$W = \{u \in PC([0, K]; X), \|u(t)\| \leq r, \|Gu(t)\| \leq w, \forall t \in [0, K]\}. \tag{3.1}$$

For the forthcoming analysis, we need the following assumptions:

- (A₁) The c_0 semigroup $T(t)$ generated by A is equicontinuous and $N = \sup\{\|T(t)\|; t \in [0, K]\}$;
- (A₂) $g : X \rightarrow X$ is such that there exist positive constants c and d such that $\|g(u)\| \leq c\|u\| + d$, for all $u \in PC([0, K]; X)$;
- (A₃) $f : [0, K] \times X \times X \rightarrow X$ is of Caratheodory type, that is, $f(\cdot, u, Gu)$ is measurable for all $u \in X$, and $f(t, \cdot, \cdot)$ is continuous for a.e. $t \in [0, K]$;
- (A₄) there exist a function $m \in L^1(0, K; R^+)$ and a nondecreasing continuous function $\Omega : R^+ \rightarrow R^+$ such that $\|f(t, u, (Gu))\| \leq m(t)\Omega(\|u\|, \|Gu\|)$ for all $u \in X$ a.e. $t \in [0, K]$;
- (A₅) there exist $L, L_1 \in L^1(0, K; R^+)$ such that for any bounded sets $D_1, D_2 \subset X$,

$$\alpha(f(t, D_1, D_2)) \leq L(t)\alpha(D_1) + L_1(t)\alpha(D_2)$$

for a.e. $t \in [0, K]$;

- (A₆) The functions $I_k : X \rightarrow X, k = 1, 2, \dots, p$, are completely continuous and uniformly bounded, and $\max_{1 \leq k \leq p, u \in W} \|I_k(u)\| = \Lambda$;
- (A₇) there exists a positive constant γ such that

$$N[\|u_0\| + (cr + d) + \Lambda] + N\Omega(r, w) \int_0^K m(s) ds \leq \gamma,$$

where N, c, d, Ω, Λ are given by assumptions (A₁), (A₂), (A₄), (A₆).

In passing, we remark that

- (i) If A is the generator of an analytic semigroup $T(t)$ or a differentiable semigroup $T(t)$, then $T(t)$ is an equicontinuous c_0 -semigroup [1].
- (ii) If $\|f(t, u_1, u_2) - f(t, v_1, v_2)\| \leq L\|u_1 - v_1\| + L_1\|u_2 - v_2\|, t \in [0, K], u_1, u_2, v_1, v_2 \in X$, then $\alpha(f(t, D_1, D_2)) \leq L(t)\alpha(D_1) + L_1(t)\alpha(D_2)$ for any bounded sets $D_1, D_2 \subset X$ and a.e. $t \in [0, K]$.

Theorem 3.1 Assume that conditions (A₁)-(A₇) hold. Then there exists at least one mild solution for problem (1.1).

Proof Let us define an operator $Q : PC([0, K]; X) \rightarrow PC([0, K]; X)$ by

$$Q u(t) = \begin{cases} T(t)[u_0 - g(u)] + \int_0^t T(t-s)f(s, u(s), Gu(s)) ds, & t \in [0, t_1]; \\ T(t - t_1)(u(t_1^-) + I_1(u(t_1^-))) \\ \quad + \int_{t_1}^t T(t-s)f(s, u(s), Gu(s)) ds, & t \in (t_1, t_2]; \\ \vdots \\ T(t - t_p)(u(t_p^-) + I_p(u(t_p^-))) \\ \quad + \int_{t_p}^t T(t-s)f(s, u(s), Gu(s)) ds, & t \in (t_p, K] \end{cases} \quad (3.2)$$

for all $u \in PC([0, K]; X)$ and show that the operator Q satisfies the hypothesis of Lemma 2.7. The proof consists of several steps.

(i) Q is continuous. Let (u_n) be a sequence in $PC([0, K]; X)$ such that $u_n \rightarrow u$ in $PC([0, K]; X)$. Then, in view of (A_3) , it follows that $f(s, u_n(s), Gu_n(s)) \rightarrow f(s, u(s), Gu(s))$ as $n \rightarrow \infty$. Now, for small $\varepsilon > 0$ and $n \rightarrow \infty$, we have

$$\begin{aligned} \|Qu_n(t) - Qu(t)\| &\leq N \int_0^{t_1} \|f(s, u_n(s), Gu_n(s)) - f(s, u(s), Gu(s))\| ds \\ &\leq \varepsilon N, \quad \forall t \in [0, t_1], \end{aligned} \quad (3.3)$$

$$\begin{aligned} \|Qu_n(t) - Qu(t)\| &\leq N[\|u_n(t_i^-) - u(t_i^-)\| \\ &\quad + \|I_i(u_n(t_i^-)) - I_i(u(t_i^-))\|] + \varepsilon N, \quad \forall t \in (t_i, t_{i+1}], \end{aligned} \quad (3.4)$$

for $i = 1, 2, \dots, p$. By assumption (A_6) together with (3.3)-(3.4), we obtain

$$\lim_{n \rightarrow \infty} \|Qu_n - Qu\|_{PC} = 0.$$

Thus, $W \subseteq PC([0, K]; X)$ is bounded and convex (W is defined by (3.1)).

For any $u \in W$, $t \in [0, t_1]$, we have

$$\begin{aligned} \|(Qu)(t)\| &\leq \|T(t)[u_0 - g(u)]\| + \int_0^{t_1} \|T(t-s)f(s, u(s), Gu(s))\| ds \\ &\leq N[\|u_0\| + (c\|u\| + d)] + N\Omega(r, w) \int_0^{t_1} m(s) ds \\ &\leq N[\|u_0\| + (cr + d)] + N\Omega(r, w) \int_0^{t_1} m(s) ds. \end{aligned} \quad (3.5)$$

Similarly, for any $u \in W$ and $i = 1, 2, \dots, p$, we get

$$\|(Qu)(t)\| \leq N[\|u_0\| + (cr + d) + \Lambda] + N\Omega(r, w) \int_0^K m(s) ds, \quad t \in (t_i, t_{i+1}]. \quad (3.6)$$

Using (3.5)-(3.6) and (A_6) - (A_7) , we obtain

$$\|(Qu)(t)\| \leq N[\|u_0\| + (cr + d) + \Lambda] + N\Omega(r, w) \int_0^K m(s) ds \leq \gamma, \quad t \in [0, K],$$

which implies that $Q : W \rightarrow W$ is a bounded operator.

(ii) $Q(W)$ is equicontinuous, where W is defined by (3.1). For all $s_1, s_2 \in [0, t_1]$ and for each $Q \in W(u)$, we have by Lemma 2.4 that

$$\begin{aligned}
 & \| (Qu)(s_2) - (Qu)(s_1) \| \\
 & \leq \| T(s_2) - T(s_1) \| \| u_0 - g(u) \| + \left\| \int_0^{s_2} T(s_2 - s) f(s, u(s), Gu(s)) ds \right. \\
 & \quad \left. - \int_0^{s_1} T(s_1 - s) f(s, u(s), Gu(s)) ds \right\| \\
 & \leq \| T(s_2) - T(s_1) \| \| u_0 - g(u) \| + \int_0^{s_1} \| T(s_2 - s) - T(s_1 - s) \| m(s) \Omega(r, w) ds \\
 & \quad + \int_{s_1}^{s_2} \| T(s_2 - s) \| m(s) \Omega(r, w) ds \\
 & \leq \| T(s_2) - T(s_1) \| [\| u_0 \| + (cr + d)] \\
 & \quad + \int_0^{s_1} \| T(s_2 - s) - T(s_1 - s) \| m(s) \Omega(r, w) ds \\
 & \quad + \int_{s_1}^{s_2} \| T(s_2 - s) \| m(s) \Omega(r, w) ds. \tag{3.7}
 \end{aligned}$$

Similarly, for all $s_1, s_2 \in (t_i, t_{i+1}]$, with $s_1 < s_2$, $i = 1, 2, \dots, p$, we get

$$\begin{aligned}
 & \| (Qu)(s_2) - (Qu)(s_1) \| \\
 & \leq \| T(s_2) - T(s_1) \| [\| u_0 \| + (cr + d) + \Lambda] \\
 & \quad + \int_0^{s_1} \| T(s_2 - s) - T(s_1 - s) \| m(s) \Omega(r, w) ds \\
 & \quad + \int_{s_1}^{s_2} \| T(s_2 - s) \| m(s) \Omega(r, w) ds. \tag{3.8}
 \end{aligned}$$

Thus, from inequalities (3.7) and (3.8), we obtain

$$\lim_{s_1 \rightarrow s_2} \| (Qu)(s_2) - (Qu)(s_1) \| = 0.$$

So, $Q(W)$ is equicontinuous.

Let $B_0 = \overline{\text{co}}(Q(W))$. For any $B \subset B_0$ and $\varepsilon > 0$, we know from Lemma 2.5 that there is a sequence $\{u_n\}_{n=1}^\infty \subset B$ such that

$$\begin{aligned}
 \alpha(Q^1 B(t)) &= \alpha(QB(t)) \\
 &\leq 2\alpha \left(\int_0^t T(t-s) f(s, \{u_n(s)\}_{n=1}^\infty, (G\{u_n(s)\}_{n=1}^\infty)) ds + \varepsilon \right) \\
 &\leq 4 \int_0^t \alpha(T(t-s) f(s, \{u_n(s)\}_{n=1}^\infty, (G\{u_n(s)\}_{n=1}^\infty))) ds + \varepsilon \\
 &\leq 4N \int_0^t (L(s) \alpha \{u_n(s)\}_{n=1}^\infty + kL_1(s) \alpha \{u_n(s)\}_{n=1}^\infty) ds + \varepsilon \\
 &\leq 4N \left(\alpha \{u_n(s)\}_{n=1}^\infty \int_0^t L(s) ds + k \alpha \{u_n(s)\}_{n=1}^\infty \int_0^t L_1(s) ds \right) + \varepsilon
 \end{aligned}$$

$$\begin{aligned} &\leq 4N \left(\alpha_{PC}(B) \int_0^t L(s) ds + k\alpha_{PC}(B) \int_0^t L_1(s) ds \right) + \varepsilon \\ &\leq 4N\alpha_{PC}(B) \int_0^t (L(s) + kL_1(s)) ds + \varepsilon, \quad t \in [0, t_1], \end{aligned}$$

where we have used Lemma 2.3. Similarly, we have

$$\alpha(Q^1B(t)) \leq 4N\alpha_{PC}(B) \int_0^t (L(s) + kL_1(s)) ds + \Lambda_1\alpha_{PC} + \varepsilon, \quad t \in (t_i, t_{i+1}], i = 1, 2, \dots, p.$$

Using the fact that there is a continuous function $\phi : [0, K] \rightarrow R^+$ with $\max\{|\phi(t)| : t \in [0, K]\} = M$ satisfying the relation $\int_0^K |L(s) + kL_1(s) - \phi(s)| ds < \delta$ for any $\delta > 0$ ($\delta < \frac{1}{N}$), the above inequality takes the form

$$\begin{aligned} \alpha(Q^1B(t))l &\leq 4N \left(\int_0^t |L(s) + kL_1(s) - \phi(s)| ds + \int_0^t |\phi(s)| ds \right) \alpha_{PC}(B) + \Lambda_1\alpha_{PC} + \varepsilon \\ &\leq (a + bt)\alpha_{PC}(B) + \Lambda_1\alpha_{PC} + \varepsilon, \end{aligned}$$

where $a = 4N\delta$, $b = 4NM$ and $\Lambda_1 = 2N \sum_{i=1}^p \beta_i$. Again, by Lemma 2.5, for any $\varepsilon > 0$, there is a sequence $\{v_n\}_{n=1}^\infty \subset \overline{\text{co}}(Q^1B)$ such that

$$\begin{aligned} \alpha(Q^2B(t)) &= \alpha(Q(\overline{\text{co}}(Q^1B(t)))) \\ &\leq 2\alpha \left(\int_0^t T(t-s)f(s, \{v_n(s)\}_{n=1}^\infty, (G\{v_n(s)\}_{n=1}^\infty)) ds + \varepsilon \right) \\ &\leq 4 \int_0^t \alpha(T(t-s)f(s, \{v_n(s)\}_{n=1}^\infty, (G\{v_n(s)\}_{n=1}^\infty)) ds + \varepsilon \\ &\leq 4N \int_0^t (L(s)\alpha\{v_n(s)\}_{n=1}^\infty + kL_1(s)\alpha\{v_n(s)\}_{n=1}^\infty) ds + \varepsilon \\ &\leq 4N \int_0^t (L(s) + kL_1(s))(Q^1B(s)) ds + \varepsilon \\ &\leq 4N \int_0^t \{ |L(s) + kL_1(s) - \phi(s)| + |\phi(s)| \} [(a + bs) + \Lambda_1] \alpha_{PC}(B) ds + \varepsilon. \end{aligned}$$

Similarly, for $t \in (t_i, t_{i+1}]$, $i = 1, 2, \dots, p$, we can obtain

$$\begin{aligned} \alpha(Q^2B(t)) &\leq 4N \int_0^t \{ |L(s) + kL_1(s) - \phi(s)| + |\phi(s)| \} [(a + bs) + \Lambda_1] \alpha_{PC}(B) ds \\ &\quad + \Lambda_2\alpha_{PC}(B) + \varepsilon \\ &\leq 4N \int_0^t |L(s) + kL_1(s) - \phi(s)| ds [(a + bt) + \Lambda_1] \alpha_{PC}(B) \\ &\quad + 4N \int_0^t [M[(a + bs) + \Lambda_1] ds] \alpha_{PC}(B) + \Lambda_2\alpha_{PC}(B) + \varepsilon \\ &\leq \left(a^2 + 2abt + \frac{(bt)^2}{2!} \right) \alpha_{PC}(B) + [(a + bt)\Lambda_1] \alpha_{PC}(B) + \Lambda_2\alpha_{PC}(B) + \varepsilon. \end{aligned}$$

Hence, by mathematical induction, for any positive integer n and $t \in [0, K]$, we obtain

$$\begin{aligned} \alpha_{PC}(Q^n B(t)) &\leq \left(a^n + C_n^1 a^{n-1}(bt) + C_n^2 a^{n-2} \frac{(bt)^2}{2!} + \dots + \frac{(bt)^n}{n!} \right) \alpha_{PC}(B) \\ &\quad + \left(a^{n-1} + C_{n-1}^1 a^{n-2}(bt) + C_{n-1}^2 a^{n-3} \frac{(bt)^2}{2!} + \dots + \frac{(bt)^{n-1}}{(n-1)!} \right) \Lambda_{n-1} \alpha_{PC}(B) \\ &\quad + \Lambda_n \alpha_{PC}(B). \end{aligned}$$

By Lemma 2.2, we have

$$\begin{aligned} \alpha(Q^n B) &\leq \left(a^n + C_n^1 a^{n-1} b + C_n^2 a^{n-2} \frac{(b)^2}{2!} + \dots + \frac{(b)^n}{n!} \right) \alpha_{PC}(B) \\ &\quad + \left(a^{n-1} + C_{n-1}^1 a^{n-2}(b) + C_{n-1}^2 a^{n-3} \frac{(b)^2}{2!} + \dots + \frac{(b)^{n-1}}{(n-1)!} \right) \Lambda_{n-1} \alpha_{PC}(B) \\ &\quad + \Lambda_n \alpha_{PC}(B). \end{aligned}$$

From Lemma 2.6, there exists a positive integer n_0 such that

$$\begin{aligned} a^{n_0} + C_{n_0}^1 a^{n_0-1} b + C_{n_0}^2 a^{n_0-2} \frac{(b)^2}{2!} + \dots + \frac{(b)^{n_0}}{n_0!} &= R < 1, \\ a^{n_0-1} + C_{n_0-1}^1 a^{n_0-2}(b) + C_{n_0-1}^2 a^{n_0-3} \frac{(b)^2}{2!} + \dots + \frac{(b)^{n_0-1}}{(n_0-1)!} &= T < 1, \\ \Lambda_{n_0-1} = l \quad \text{and} \quad \Lambda_{n_0} = q. \end{aligned}$$

Then

$$\begin{aligned} \alpha_{PC}(Q^{n_0} B) &\leq R \alpha_{PC}(B) + T l \alpha_{PC} + q \alpha_{PC} \\ &\leq (R + Tl + q) \alpha_{PC}. \end{aligned}$$

Thus, it follows by Lemma 2.7 that Q has at least one fixed point in B_0 , that is, the nonlocal integrodifferential equation (1.1) has at least one mild solution in B_0 . This completes the proof. \square

Remark 3.2 In [38], the author discussed the nonlocal initial value problem by taking f to be compact in (1.1). From the above theorem, we notice that the key condition in [38] is no more required. So, Theorem 3.1 generalizes the related results in [38]. Furthermore, we extend the problem addressed in [23] to the impulse case with the nonlinearity of a more general form $f(t, u(t), Gu(t))$.

Theorem 3.3 *If assumptions (A_1) - (A_6) are satisfied, then there is at least one mild solution for (1.1) provided that*

$$\int_0^K m(s) ds < \lim_{T \rightarrow \infty} \frac{T - N[\|u_0\| + (cT + d) + \Lambda]}{N\Omega(T, w)}.$$

Proof We do not provide the proof of this theorem as it is similar to that of Theorem 3.1. \square

For our final result, we introduce the following condition:

(A₂') Let $g : PC([0, K]; X) \rightarrow X$ be continuous and compact. Then there exists a positive constant $M > 0$ such that $\|g(u)\| \leq M$ for every $u \in PC([0, K]; X)$.

Theorem 3.4 *Suppose that conditions (A₁)-(A₂') and (A₃)-(A₆) hold. Then there exists at least one mild solution for (1.1) if there exists a constant r such that*

$$N[\|u_0\| + M + \Lambda] + N\Omega(r, w) \int_0^K m(s) ds \leq \gamma.$$

Proof We omit the proof as it is similar to that of Theorem 3.1. This completes the proof. □

Example Consider a nonlocal problem of integrodifferential equations given by

$$\begin{aligned} \frac{\partial u(t, w)}{\partial t} &= \frac{\partial^2 u(t, w)}{\partial w^2} u(t, w) \\ &+ \sin^2 u(t, w) + \int_0^K \frac{u(s, w)}{\sqrt{(1+t)(2+s)^3}} ds, \quad t \in [0, K], \\ u(0) &= u_0 + \int_0^K \sqrt{3+s^2} \log(2 + |u(s, w)|) ds, \\ \Delta u(t) &= \int_0^K \frac{(1 + |\cos^2 u(s)|)}{\sqrt{t^2 + s^2 + 7}} ds, \quad i \geq 1, \end{aligned} \tag{3.9}$$

with $0 < t_1 < t_2 < \dots < t_p < K$, $0 < s_1 < s_2 < \dots < s_q < K$. Let us take $X = L^2([0, K], R)$ and define the operator A by $A(t)y = y''$ with the domain $D(A) = \{y \in X: y, y' \text{ are absolutely continuous, } y'' \in X, y(0) = 0, y(K) = 0\}$. The assumptions of Theorem 3.1 clearly hold for a large positive constant γ . Hence the conclusion of Theorem 3.1 applies to problem (3.9).

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

Each of the authors, BA, KM and KK, contributed to each part of this study equally and read and approved the final version of the manuscript.

Author details

¹Department of Mathematics, Faculty of Science, King Abdulaziz University, P.O. Box 80203, Jeddah, 21589, Saudi Arabia. ²Department of Mathematics, Erode Arts and Science College, Erode, Tamil Nadu 638 009, India. ³Department of Mathematics, K.S.R. College of Technology, Tiruchengode, Tamil Nadu 637 215, India.

Acknowledgements

The authors thank the reviewers for their useful comments that led to the improvement of the original manuscript. The research of B. Ahmad was partially supported by Deanship of Scientific Research (DSR), King Abdulaziz University, Jeddah, Saudi Arabia.

Received: 29 March 2013 Accepted: 20 June 2013 Published: 9 July 2013

References

1. Pazy, A: Semigroups of Linear Operators and Applications to Partial Differential Equations. Springer, New York (1983)
2. Dong, Q, Li, G: Existence of solutions for semilinear differential equations with nonlocal conditions in Banach spaces. Electron. J. Qual. Theory Differ. Equ. **2009**, 47 (2009)
3. Kato, T: Quasilinear equations of evolution with applications to partial differential equations. Lect. Notes Math. **448**, 25-70 (1975)

4. Kato, T: Abstract evolution equation linear and quasilinear, revisited. *Lect. Notes Math.* **1540**, 103-125 (1993)
5. Ntouyas, SK, Tsamotas, PC: Global existence for semilinear integrodifferential equations with delay and nonlocal conditions. *Appl. Anal.* **64**, 99-105 (1997)
6. Anguraj, A, Karthikeyan, K: Existence of solutions for impulsive neutral functional differential equations with nonlocal conditions. *Nonlinear Anal.* **70**, 2717-2721 (2009)
7. Guo, DJ, Lakshmikantham, V, Liu, XZ: *Nonlinear Integral Equations in Abstract Spaces*. Kluwer Academic, Dordrecht (1996)
8. Lakshmikantham, V, Bainov, DD, Simeonov, PS: *Theory of Impulsive Differential Equations*. World Scientific, Singapore (1989)
9. Samoilenko, AM, Perestyuk, NA: *Impulsive Differential Equations*. World Scientific, Singapore (1995)
10. Fan, Z, Li, G: Existence results for semilinear differential equations with nonlocal and impulsive conditions. *J. Funct. Anal.* **258**, 1709-1727 (2010)
11. Liu, JH: Nonlinear impulsive evolution equations. *Dyn. Contin. Discrete Impuls. Syst.* **6**, 77-85 (1999)
12. Migorski, S, Ochal, A: Nonlinear impulsive evolution inclusions of second order. *Dyn. Syst. Appl.* **16**, 679-687 (1997)
13. Byszewski, L: Theorems about the existence and uniqueness of solutions of a semilinear evolution nonlocal Cauchy problem. *J. Math. Anal. Appl.* **162**, 494-505 (1991)
14. Byszewski, L, Akca, H: Existence of solutions of a semilinear functional differential evolution nonlocal problem. *Nonlinear Anal.* **34**, 65-72 (1998)
15. Aizicovici, S, McKibben, M: Existence results for a class of abstract nonlocal Cauchy problems. *Nonlinear Anal.* **39**, 649-668 (2000)
16. Ji, SC, Wen, S: Nonlocal Cauchy problem for impulsive differential equations in Banach spaces. *Int. J. Nonlinear Sci.* **10**(1), 88-95 (2010)
17. Liang, J, Liu, JH, Xiao, TJ: Nonlocal Cauchy problem governed by compact operator families. *Nonlinear Anal.* **57**, 183-189 (2004)
18. Byszewski, L, Lakshmikantham, V: Theorems about the existence and uniqueness of solutions of nonlocal Cauchy problem in a Banach space. *Appl. Anal.* **40**, 11-19 (1990)
19. Ntouyas, S, Tsamotas, P: Global existence for semilinear evolution equations with nonlocal conditions. *J. Math. Anal. Appl.* **210**, 679-687 (1997)
20. Ntouyas, SK, Tsamotas, PC: Global existence for second order semilinear ordinary and delay integrodifferential equations with nonlocal conditions. *Appl. Anal.* **67**(3-4), 245-257 (1997)
21. Zhu, T, Song, C, Li, G: Existence of mild solutions for abstract semilinear evolution equations in Banach spaces. *Nonlinear Anal.* **75**, 177-181 (2012)
22. Agarwal, RP, Ahmad, B, Alsaedi, A, Shahzad, N: Existence and dimension of the set of mild solutions to semilinear fractional differential inclusions. *Adv. Differ. Equ.* **2012**, 74 (2012)
23. Lizama, C, Pozo, JC: Existence of mild solutions for a semilinear integrodifferential equation with nonlocal initial conditions. *Abstr. Appl. Anal.* **2012**, Article ID 647103 (2012)
24. Shaochun, J, Gang, L: A unified approach to nonlocal impulsive differential equations with the measure of noncompactness. *Adv. Differ. Equ.* **2012**, 182 (2012)
25. Xue, X: Nonlocal nonlinear differential equations with a measure of noncompactness in Banach spaces. *Nonlinear Anal.* **70**, 2593-2601 (2009)
26. Agarwal, RP, Benchohra, M, Seba, D: On the application of measure of noncompactness to the existence of solutions for fractional differential equations. *Results Math.* **55**, 221-230 (2009)
27. Dong, Q, Li, G: Existence of solutions for semilinear differential equations with nonlocal conditions in Banach spaces. *Electron. J. Qual. Theory Differ. Equ.* **2009**, 47 (2009)
28. Diagana, T, Mophou, GM, N'Guérékata, GM: On the existence of mild solutions to some semilinear fractional integro-differential equations. *Electron. J. Qual. Theory Differ. Equ.* **2010**, 58 (2010)
29. Diagana, T: Existence of pseudo-almost automorphic mild solutions to some nonautonomous partial evolution equations. *Adv. Differ. Equ.* **2011**, Article ID 895079 (2011)
30. Ji, S, Li, G: Existence results for impulsive differential inclusions with nonlocal conditions. *Comput. Math. Appl.* **62**, 1908-1915 (2011)
31. Herzallah, MAE, Baleanu, D: Existence of a periodic mild solution for a nonlinear fractional differential equation. *Comput. Math. Appl.* **64**(10), 3059-3064 (2012)
32. Liang, J, Liu, JH, Xiao, TJ: Nonlocal impulsive problems for nonlinear differential equations in Banach spaces. *Math. Comput. Model.* **49**, 798-804 (2009)
33. Zhu, L, Dong, Q, Li, G: Impulsive differential equations with nonlocal conditions in general Banach spaces. *Adv. Differ. Equ.* **2012**, 10 (2012)
34. Baleanu, D, Mustafa, OG, Agarwal, RP: Asymptotically linear solutions for some linear fractional differential equations. *Abstr. Appl. Anal.* **2010**, Article ID 865139 (2010)
35. Baleanu, D, Mustafa, OG, Agarwal, RP: Asymptotic integration of $(1 + \alpha)$ -order fractional differential equations. *Comput. Math. Appl.* **62**, 1492-1500 (2011)
36. Debbouche, A, Baleanu, D, Agarwal, RP: Nonlocal nonlinear integrodifferential equations of fractional orders. *Bound. Value Probl.* **2012**, 78 (2012)
37. Baleanu, D, Diethelm, DK, Scalas, E, Trujillo, JJ: *Fractional Calculus Models and Numerical Methods*. Series on Complexity, Nonlinearity and Chaos. World Scientific, Singapore (2012)
38. Fan, Z, Dong, Q, Li, G: Semilinear differential equations with nonlocal conditions in Banach spaces. *Int. J. Nonlinear Sci.* **2**, 131-139 (2006)
39. Xue, X: Semilinear nonlocal differential equations with measure of noncompactness in Banach spaces. *J. Nanjing Univ. Math. Big.* **24**, 264-276 (2007)
40. Banas, J, Goebel, K: *Measure of Noncompactness in Banach Space*. Lecture Notes in Pure and Applied Mathematics, vol. 60. Marcel Dekker, New York (1980)
41. Mönch, H: Boundary value problems for nonlinear ordinary differential equations of second order in Banach spaces. *Nonlinear Anal.* **4**, 985-999 (1980)

42. Bothe, D: Multivalued perturbation of m -accretive differential inclusions. *Isr. J. Math.* **108**, 109-138 (1998)
43. Zhang, X, Liu, LS, Wu, CX: Global solutions of nonlinear second order impulsive integrodifferential equations of mixed type in Banach spaces. *Nonlinear Anal.* **67**, 2335-2349 (2007)
44. Lou, B: Fixed points for operators in a space of continuous functions and applications. *Proc. Am. Math. Soc.* **127**, 2259-2264 (1999)
45. Liu, LS, Guo, F, Wu, CX, Wu, YH: Existence theorems of global solutions for nonlinear Volterra type integral equations in Banach spaces. *J. Math. Anal. Appl.* **309**, 638-649 (2005)

doi:10.1186/1687-1847-2013-205

Cite this article as: Ahmad et al.: A study of nonlocal problems of impulsive integrodifferential equations with measure of noncompactness. *Advances in Difference Equations* 2013 2013:205.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
