

RESEARCH

Open Access



An oscillation criterion in 4th-order neutral differential equations with a continuously distributed delay

George E. Chatzarakis¹, Elmetwally M. Elabbasy² and Omar Bazighifan^{3*}

*Correspondence:

o.bazighifan@gmail.com

³Department of Mathematics, Faculty of Science, Hadhramout University, Hadhramout, Yemen Full list of author information is available at the end of the article

Abstract

In this paper, a class of 4th-order neutral delay differential equations with continuously distributed delay is studied. We establish a new oscillation criterion using the Riccati transformation. An example illustrating the results is also given.

MSC: 34K10; 34K11

Keywords: 4th-order; Neutral differential equations; Oscillatory solutions

1 Introduction

In this work, we consider a 4th-order neutral differential equation with a continuously distributed delay of the form

$$[r(t)([x(t) + p(t)x(\tau(t))]''')^\alpha]' + \int_a^b q(t, \xi)f(x(g(t, \xi))) d\xi = 0. \quad (1)$$

We assume that the following conditions hold:

(H₁) α is a quotient of odd positive integers;

(H₂) $p, q, \tau, g \in C([t_0, \infty), \mathbb{R})$, $r(t)$, and $q(t, \xi)$ are positive, $0 \leq p(t) \leq p < 1$, $r(t) \in C^1([t_0, +\infty))$, $r'(t) \geq 0$, $\tau(t) \leq t$, $g(t, \xi) \leq t$, $\lim_{t \rightarrow \infty} \tau(t) = \infty$, $\lim_{t \rightarrow \infty} g(t, \xi) = \infty$, $q(t, \xi)$ is not zero on any half line $[t_\lambda, \infty) \times [a, b]$, $t_\lambda \geq t_0$, for $t \geq t_0$, $\xi \in [a, b]$, $g(t, \xi)$ is nondecreasing with respect to ξ .

(H₃) There exists a constant $k > 0$ such that $f(u)/u^\gamma \geq k$ for $u \neq 0$.

We define the corresponding function $z(t)$ of a solution $x(t)$ of (1) by $z(t) = x(t) + p(t)x(\tau(t))$, we mean a non-trivial real function $x(t) \in C([t_x, \infty))$, $t_x \geq t_0$, satisfying (1) on $[t_x, \infty)$ and, moreover, having the properties: $z(t)$, $z'(t)$, $z''(t)$ and $r(t)[z'''(t)]^\alpha$ are continuously differentiable for all $t \in [t_x, \infty)$. We consider only those solutions $x(t)$ of (1) which satisfy $\sup\{|x(t)| : t \geq T\} > 0$ for any $T \geq t_x$. A solution of (1) is called oscillatory if it has arbitrary large zeros, otherwise it is called nonoscillatory.

The oscillations of higher- and fourth-order differential equations have been studied by several authors, and several techniques have been proposed for obtaining oscillatory criteria for higher- and fourth-order differential equations. For treatments on this subject, we refer the reader to the texts [2, 5, 16–18, 21] and the articles [1, 3–15, 19–26]. In what

follows, we review some results that have provided the background and motivation for the present work.

Cesarano and Bazighifan [8], Moaaz et al. [21], and Zhang et al. [26] studied the oscillation of the fourth-order nonlinear differential equation with a continuously distributed delay

$$[r(t)(x'''(t))^\alpha]' + \int_a^b q(t, \xi)f(x(g(t, \xi))) d\xi = 0. \tag{2}$$

Li et al. [19] studied the oscillatory behavior of the fourth-order nonlinear differential equation

$$[r(t)z(t)]^{(4)} + q(t)x(\tau(t)) = 0. \tag{3}$$

Parhi and Tripathy [23] have considered the fourth-order neutral differential equations of the form

$$[r(t)(y(t) + p(t)y(t - \tau))'''] + q(t)G(y(t - \sigma)) = 0 \tag{4}$$

and

$$[r(t)(y(t) + p(t)y(t - \tau))'''] + q(t)G(y(t - \sigma)) = f(t) \tag{5}$$

and established the oscillation and asymptotic behavior of the above equations under the conditions

$$\int_{t_0}^\infty \frac{1}{r^{\frac{1}{\alpha}}(t)} dt < \infty \tag{6}$$

and

$$\int_{t_0}^\infty \frac{1}{r^{\frac{1}{\alpha}}(t)} dt = \infty, \tag{7}$$

respectively.

Our aim in the present paper is to use the Riccati method to establish new conditions for the oscillation of all solutions of (1) under condition (7).

The proof of our main results is essentially based on the following lemmas.

Lemma 1.1 *Let $\beta \geq 1$ be a ratio of two numbers, where U and V are constants. Then*

$$Uy - Vy^{\frac{\beta+1}{\beta}} \leq \frac{\beta^\beta}{(\beta + 1)^{\beta+1}} \frac{U^{\beta+1}}{V^\beta}, \quad V > 0. \tag{8}$$

Lemma 1.2 *If the function z satisfies $z^{(i)} > 0, i = 0, 1, \dots, n$, and $z^{(n+1)} < 0$, then*

$$\frac{z(t)}{t^n/n!} \geq \frac{z'(t)}{t^{n-1}/(n-1)!}. \tag{9}$$

Lemma 1.3 *Let $h \in C^n([t_0, \infty), (0, \infty))$. Assume that $h^{(n)}(t)$ is of a fixed sign and not identically zero on $[t_0, \infty)$ and that there exists $t_1 \geq t_0$ such that $h^{(n-1)}(t)h^{(n)}(t) \leq 0$ for all $t \geq t_1$. If $\lim_{t \rightarrow \infty} h(t) \neq 0$, then for every $\lambda \in (0, 1)$ there exists $t_\lambda \geq t_0$ such that*

$$h(t) \geq \frac{\lambda}{(n-1)!} t^{n-1} |h^{(n-1)}(t)| \quad \text{for all } t \geq t_\lambda. \tag{10}$$

2 Main results

In this section, we shall establish some oscillation criteria for equation (1).

For convenience, we denote

$$\begin{aligned} \eta(t) &= \int_{t_0}^\infty r^{-1/\alpha}(s) ds, \\ \rho'_+(t) &:= \max\{0, \rho'(t)\}, \\ \vartheta'_+(t) &:= \max\{0, \vartheta'(t)\}, \\ Q(t) &= \int_a^b q(t, \xi) d\xi \end{aligned}$$

and

$$Q^*(v) = \int_v^\infty kQ(s)(1-p)^\alpha (g(s, a)/s)^{3\alpha} ds. \tag{11}$$

Theorem 2.1 *Assume that (7) holds. If there exist positive functions $\rho, \vartheta \in C([t_0, \infty))$ such that*

$$\int_{t_0}^\infty \left(\Psi(s) - \frac{2^\alpha}{(\alpha+1)^{\alpha+1}} \frac{r(s)(\rho'(s))^{\alpha+1}}{\mu^\alpha s^{2\alpha} \rho^\alpha(s)} \right) ds = \infty \tag{12}$$

for some $\mu \in (0, 1)$, where

$$\Psi(t) = k\rho(t)Q(t)(1-p)^\alpha (g(t, a)/t)^{3\alpha}, \tag{13}$$

and either

$$\int_{t_0}^\infty kQ(s)(1-p)^\alpha (g(s, a)/s)^\alpha ds = \infty \tag{14}$$

or

$$\int_{t_0}^\infty (Q^*(v))^{1/\alpha} r^{-1/\alpha}(v) dv = \infty, \tag{15}$$

or

$$\int_{t_0}^\infty \left[\vartheta(t) \int_t^\infty (Q^*(v))^{1/\alpha} r^{-1/\alpha}(v) dv - \frac{\vartheta_+^2(t)}{4\vartheta(t)} \right] dt = \infty, \tag{16}$$

then all solutions of (1) are oscillatory.

Proof Let x be a nonoscillatory solution of equation (1) defined in the interval $[t_0, \infty)$. Without loss of generality, we can assume that $x(t)$ is eventually positive. It follows from (1) that there are two possible cases for $t \geq t_1$, where $t_1 \geq t_0$ is sufficiently large:

- (C₁) $z'(t) > 0, z''(t) > 0, z'''(t) > 0, (r(t)(z'''(t))^\alpha) < 0,$
- (C₂) $z'(t) > 0, z''(t) < 0, z'''(t) > 0, (r(t)(z'''(t))^\alpha) < 0.$

Assume that Case (C₁) holds. Since $\tau(t) \leq t$ and $z'(t) > 0$, we get

$$\begin{aligned} z(t) &= x(t) + p(t)x(\tau(t)), \\ x(t) &= z(t) - p(t)x(\tau(t)) \\ &\geq z(t) - p(t)z(\tau(t)) \\ &= (1 - p(t))z(t). \end{aligned}$$

From equation (1), we see that

$$(r(t)(z'''(t))^\alpha)' = - \int_a^b q(t, \xi)f(x(g(t, \xi))) d\xi \tag{17}$$

so that

$$\begin{aligned} (r(t)(z'''(t))^\alpha)' &= - \int_a^b kq(t, \xi)x^\alpha(g(t, \xi)) d\xi \\ &\leq - \int_a^b kq(t, \xi)(1 - p(g(t, \xi)))^\alpha z^\alpha(g(t, \xi)) d\xi. \end{aligned}$$

Since $g(t, \xi)$ is nondecreasing with respect to ξ and $z'(t) > 0$, we have

$$z(g(t, a)) \leq z(g(t, \xi)) \leq z(g(t, b)). \tag{18}$$

Thus

$$(r(t)(z'''(t))^\alpha)' \leq -(1 - p)^\alpha \int_a^b kq(t, \xi)z^\alpha(g(t, \xi)) d\xi \tag{19}$$

$$\leq -k(1 - p)^\alpha z^\alpha(g(t, a)) \int_a^b q(t, \xi) d\xi \tag{20}$$

$$= -kQ(t)(1 - p)^\alpha z^\alpha(g(t, a)). \tag{21}$$

Now, we define a generalized Riccati substitution by

$$\omega(t) := \rho(t) \frac{r(t)(z'''(t))^\alpha}{z^\alpha(t)}. \tag{22}$$

Then $\omega(t) > 0$. Differentiating and using (19), we obtain

$$\omega'(t) \leq \frac{\rho'(t)}{\rho(t)}\omega(t) - k\rho(t)Q(t)(1 - p)^\alpha \frac{z^\alpha(g(t, a))}{z^\alpha(t)} \tag{23}$$

$$- \alpha\rho(t) \frac{r(t)(z'''(t))^\alpha}{z^{\alpha+1}(t)} z'(t). \tag{24}$$

From Lemma 1.2, we have that $z(t) \geq \frac{t}{3}z'(t)$, and hence

$$\frac{z(g(t, a))}{z(t)} \geq \frac{g^3(t, a)}{t^3}. \tag{25}$$

Since $r'(t) > 0$ and $(r(t)(z'''(t))^\alpha)' \leq 0$, we get $z^{(4)}(t) < 0$. It follows from Lemma 1.3 that

$$z'(t) \geq \frac{\mu}{2}t^2z'''(t) \tag{26}$$

for all $\mu \in (0, 1)$ and every sufficiently large t . Thus, by (24), (25), and (26), we get

$$\begin{aligned} \omega'(t) \leq & \frac{\rho'(t)}{\rho(t)}\omega(t) - k\rho(t)Q(t)(1-p)^\alpha \left(\frac{g(t, a)}{t}\right)^{3\alpha} \\ & - \alpha\mu \frac{t^2}{2r^{1/\alpha}(t)\rho^{1/\alpha}(t)}\omega^{\frac{\alpha+1}{\alpha}}(t). \end{aligned}$$

Using Lemma 1.1 with $U = \frac{\rho'(t)}{\rho(t)}$, $V = \frac{\alpha\mu t^2}{2r^{1/\alpha}(t)\rho^{1/\alpha}(t)}$ and $y = \omega$, we get

$$\omega'(t) \leq -\Psi(t) + \frac{2^\alpha}{(\alpha + 1)^{\alpha+1}} \frac{r(t)(\rho'(t))^{\alpha+1}}{\mu^\alpha t^{2\alpha} \rho^\alpha(t)}. \tag{27}$$

This implies that

$$\int_{t_1}^t \left(\Psi(s) - r(s)(\rho'(s)/\alpha + 1)^{\alpha+1} \left(\frac{2}{\mu s^2 \rho(s)}\right)^\alpha \right) ds \leq \omega(t_1) \tag{28}$$

for some $\mu \in (0, 1)$ which contradicts (12).

Assume that Case (C₂) holds. Integrating (1) from t_1 to t , we obtain

$$-r(t_1)(z'''(t_1))^\alpha \leq - \int_{t_1}^t kQ(t)x^\alpha(g(s, \xi)) ds. \tag{29}$$

From $z'(t) > 0$, $x(t) \geq (1 - p(t))z(t)$ and $g(s, \xi) \leq t$, it follows that

$$\int_{t_1}^t kQ(s)(1 - p)^\alpha z^\alpha(g(s, a)) ds \leq r(t_1)(z'''(t_1))^\alpha. \tag{30}$$

From (25), we get

$$\int_{t_1}^t kQ(s)(1 - p)^\alpha (g(s, a)/s)^{3\alpha} ds \leq r(t_1) \left(\frac{z'''(t_1)}{z(t_1)}\right)^\alpha, \tag{31}$$

which contradicts (12). Integrating (1) from t to ∞ , we obtain

$$-r(t)(z'''(t))^\alpha \leq - \int_t^\infty kQ(t)x^\alpha(g(s, \xi)) ds. \tag{32}$$

By virtue of $z'(t) > 0$, $x(t) \geq (1 - p(t))z(t)$, $g(s, \xi) \leq t$, and (25), we obtain

$$-(z'''(t)) + \frac{z(t)}{r(t)^{1/\alpha}} \left[\int_t^\infty kQ(s)(1 - p)^\alpha (g(s, a)/s)^{3\alpha} ds \right]^{1/\alpha} \leq 0. \tag{33}$$

Integrating (33) from t_1 to t , we get

$$\int_{t_1}^t \frac{1}{r^{1/\alpha}(v)} \left(\int_v^\infty kQ(s)(1-p)^\alpha (g(s,a)/s)^{3\alpha} ds \right)^{1/\alpha} dv \leq -\frac{z''(t_1)}{z(t_1)}. \tag{34}$$

This yields

$$\int_{t_1}^t (Q^*(v))^{\frac{1}{\alpha}} r^{\frac{-1}{\alpha}}(v) dv \leq -\frac{z''(t_1)}{z(t_1)}, \tag{35}$$

which contradicts (15). Integrating (33) from t to ∞ , we get

$$\int_t^\infty (Q^*(v))^{\frac{1}{\alpha}} r^{\frac{-1}{\alpha}}(v) dv \leq -\frac{z''(t)}{z(t)}, \tag{36}$$

so that

$$z''(t) + z(t) \int_t^\infty (Q^*(v))^{\frac{1}{\alpha}} r^{\frac{-1}{\alpha}}(v) dv \leq 0. \tag{37}$$

Now, we define the Riccati substitution

$$\psi(t) := \vartheta(t) \frac{z'(t)}{z(t)}, \tag{38}$$

then $\psi(t) > 0$ for $t \geq t_1$ and

$$\psi'(t) := \vartheta'(t) \frac{z'(t)}{z(t)} + \vartheta(t) \frac{z''(t)z(t) - (z'(t))^2}{z^2(t)}. \tag{39}$$

From (37) and (38), it follows that

$$\psi'(t) \leq -\vartheta(t) \int_t^\infty (Q^*(v))^{\frac{1}{\alpha}} r^{\frac{-1}{\alpha}}(v) dv + \frac{(\vartheta'(t))_+}{\vartheta(t)} \psi(t) - \frac{1}{\vartheta(t)} \psi^2(t). \tag{40}$$

Hence, we have

$$\psi'(t) \leq -\vartheta(t) \int_t^\infty (Q^*(v))^{\frac{1}{\alpha}} r^{\frac{-1}{\alpha}}(v) dv + \frac{((\vartheta'(t))_+)^2}{4\vartheta(t)}. \tag{41}$$

Integrating (41) from t_1 to s , we get

$$\int_{t_1}^s \left(\vartheta(t) \int_t^\infty (Q^*(v))^{\frac{1}{\alpha}} r^{\frac{-1}{\alpha}}(v) dv + \frac{((\vartheta'(t))_+)^2}{4\vartheta(t)} \right) dt \leq \psi(t_1) \tag{42}$$

for all large s , which contradicts (16).

The proof of the theorem is complete. □

Let $\rho(t) = t^3$ and $\vartheta(t) = t$. As a consequence of Theorem 2.1, we obtain the following oscillation criterion.

Corollary Assume that (7) holds and for some constant $\lambda_0 \in (0, 1)$

$$\int_{t_0}^{\infty} \left(\Psi^*(t) - \left(\frac{3}{\alpha + 1} \right)^{\alpha+1} \left(\frac{2}{\lambda_0} \right)^{\alpha} t^{2-3\alpha} r(t) \right) ds = \infty, \tag{43}$$

where

$$\Psi^*(t) = kt^3 Q(t)(1 - p)^{\alpha} (g(t, a)/t)^{3\alpha}. \tag{44}$$

If either (14) or (15) is satisfied, or

$$\int_{t_0}^{\infty} \left[t \int_t^{\infty} (Q^*(v))^{\frac{1}{\alpha}} r^{\frac{1}{\alpha}}(v) dv - \frac{1}{4t} \right] dt = \infty, \tag{45}$$

then all solutions of (1) are oscillatory.

3 Example

In this section, we give the following example to illustrate our main results.

Example Consider the differential equation

$$\left(\left[x(t) + \frac{1}{2} x\left(\frac{t}{3}\right) \right]''' \right)' + \int_0^1 (v/t^4) \xi x\left(\frac{t-\xi}{2}\right) d\xi = 0, \tag{46}$$

where $v > 0$ is a constant. Let

$$\begin{aligned} \alpha &= 1, & r(t) &= 1, & p(t) &= \frac{1}{2}, & \tau(t) &= \frac{t}{3}, \\ g(t, a) &= \frac{t}{2}, & q(t, \xi) &= (v/t^4)\xi, & f(x) &= x, \end{aligned} \tag{47}$$

we get

$$\begin{aligned} \eta(s) &= \int_{t_0}^{\infty} ds = \infty, \\ Q(t) &= \int_0^1 q(t, \xi) d\xi = \frac{v}{2t^4} \\ Q^*(t) &= \int_{t_0}^{\infty} kQ(s)(1 - p)^{\alpha} (g(s, a)/s)^{3\alpha} ds \\ &= \int_{t_0}^{\infty} \frac{v}{32s^4} ds = \frac{v}{96t^3}. \end{aligned}$$

If we now set $k = 1$, then

$$\int_{t_0}^{\infty} \left(kt^3 Q(t)(1 - p)^{\alpha} (g(t, a)/t)^{3\alpha} - \left(\frac{3}{\alpha + 1} \right)^{\alpha+1} \left(\frac{2}{\lambda_0} \right)^{\alpha} t^{2-3\alpha} r(t) \right) ds \tag{48}$$

$$= \left(\frac{v}{32} - \frac{9}{2\lambda_0} \right) \int_{t_0}^{\infty} \frac{1}{t} dt = \infty, \quad \text{if } v > \frac{144}{\lambda_0} \text{ for some constant } \lambda_0 \in (0, 1), \tag{49}$$

and

$$\begin{aligned}
 & \int_{t_0}^{\infty} \left[t \int_t^{\infty} (Q^*(v))^{\frac{1}{\alpha}} r^{\frac{-1}{\alpha}}(v) dv - \frac{1}{4t} \right] dt \\
 &= \int_{t_0}^{\infty} \left[t \int_t^{\infty} \frac{v}{96t^3} dt - \frac{1}{4t} \right] dt \\
 &= \int_{t_0}^{\infty} \left[t \frac{v}{192t^2} - \frac{1}{4t} \right] dt \\
 &= \left(\frac{v}{192} - \frac{1}{4} \right) \int_{t_0}^{\infty} \frac{1}{t} dt \\
 &= \infty, \quad \text{if } v > 48.
 \end{aligned}$$

Thus, by Corollary, every solution of equation (46) is oscillatory.

4 Conclusion

The results of this paper are presented in a form which is essentially new and of high degree of generality. In this paper, using a Riccati transformation technique, we offer some new sufficient conditions which ensure that any solution of Eq. (1) oscillates under the condition $\int_{t_0}^{\infty} \frac{1}{r^{\frac{1}{\alpha}}(t)} dt = \infty$. Further, we can consider the case of $z(t) = x(t) - p(t)x(\tau(t))$, and we can try to get some oscillation criteria of Eq. (1) in the future work.

Acknowledgements

The authors express their debt of gratitude to the editors and the anonymous referee for accurate reading of the manuscript and beneficial comments.

Funding

The authors received no direct funding for this work.

Competing interests

The authors declare that there are no competing interests between them.

Authors' contributions

The authors declare that they have read and approved the final manuscript.

Author details

¹Department of Electrical and Electronic Engineering Educators, School of Pedagogical and Technological Education (ASPETE), Athens, Greece. ²Department of Mathematics, Faculty of Science, Mansoura University, Mansoura, Egypt.

³Department of Mathematics, Faculty of Science, Hadhramout University, Hadhramout, Yemen.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 17 January 2019 Accepted: 1 August 2019 Published online: 09 August 2019

References

1. Agarwal, R., Grace, S., Manojlovic, J.: Oscillation criteria for certain fourth order nonlinear functional differential equations. *Math. Comput. Model.* **44**, 163–187 (2006)
2. Agarwal, R., Grace, S., O'Regan, D.: *Oscillation Theory for Difference and Functional Differential Equations*. Kluwer Academic, Dordrecht (2000)
3. Agarwal, R., Grace, S., O'Regan, D.: Oscillation criteria for certain nth order differential equations with deviating arguments. *J. Math. Anal. Appl.* **262**, 601–622 (2001)
4. Baculikova, B., Dzurina, J., Graef, J.R.: On the oscillation of higher-order delay differential equations. *Math. Slovaca* **187**, 387–400 (2012)
5. Bazighifan, O.: Oscillatory behavior of higher-order delay differential equations. *Gen. Lett. Math.* **2**, 105–110 (2017)
6. Bazighifan, O., Cesarano, C.: Some new oscillation criteria for second-order neutral differential equations with delayed arguments. *Mathematics* **7**, 1–8 (2019)
7. Bazighifan, O., Elabbasy, E.M., Moaaz, O.: Oscillation of higher-order differential equations with distributed delay. *J. Inequal. Appl.* **2019**, 55 (2019)

8. Cesarano, C., Bazighifan, O.: Oscillation of fourth-order functional differential equations with distributed delay. *Axioms* **7**, 1–9 (2019)
9. Cesarano, C., Bazighifan, O.: Qualitative behavior of solutions of second order differential equations. *Symmetry* **11**, 1–8 (2019)
10. Cesarano, C., Pinelas, S., Al-Showaikh, F., Bazighifan, O.: Asymptotic properties of solutions of fourth-order delay differential equations. *Symmetry* **11**, 1–10 (2019)
11. Elabbasy, E.M., Hassan, T.S., Moaaz, O.: Oscillation behavior of second order nonlinear neutral differential equations with deviating arguments. *Opuscula Mathematica* **32**, 719–730 (2012)
12. Grace, S., Graef, J., Tunc, E.: Oscillatory behavior of a third order neutral dynamic equations with distributed delays. *Electron. J. Qual. Theory Differ. Equ.* **2016**, 14 (2016)
13. Grace, S., Graef, J., Tunc, E.: Oscillatory behavior of second order damped neutral differential equations with distributed deviating arguments. *Miskolc Math. Notes* **18**, 759–769 (2017)
14. Grace, S., Lalli, B.: Oscillation theorems for nth order nonlinear differential equations with deviating arguments. *Proc. Am. Math. Soc.* **90**, 65–70 (1984)
15. Graef, J., Tunc, E.: Oscillation of fourth-order nonlinear dynamic equations on time scales. *Panam. Math. J.* **25**, 16–34 (2015)
16. Györi, I., Ladas, G.: *Oscillation Theory of Delay Differential Equations with Applications*. Clarendon, Oxford (1991)
17. Kiguradze, I., Chanturia, T.: *Asymptotic Properties of Solutions of Nonautonomous Ordinary Differential Equations*. Kluwer Academic, Dordrecht (1993)
18. Ladde, G., Lakshmikantham, V., Zhang, B.: *Oscillation Theory of Differential Equations with Deviating Arguments*. Dekker, New York (1987)
19. Li, T., Baculikova, B., Dzurina, J., Zhang, C.: Oscillation of fourth order neutral differential equations with p-Laplacian like operators. *Bound. Value Probl.* **2014**, 56 (2014)
20. Moaaz, O., Chalishajar, D., Bazighifan, O.: Some qualitative behavior of solutions of general class difference equations. *Mathematics* **7**, 1–12 (2019)
21. Moaaz, O., Elabbasy, E.M., Bazighifan, O.: On the asymptotic behavior of fourth-order functional differential equations. *Adv. Differ. Equ.* **2017**, 261 (2017)
22. Moaaz, O., Elabbasy, E.M., Muhib, A.: Oscillation criteria for even-order neutral differential equations with distributed deviating arguments. *Adv. Differ. Equ.* **2019**, 297 (2019)
23. Parhi, N., Tripathy, A.: On oscillatory fourth order linear neutral differential equations-I. *Math. Slovaca* **54**, 389–410 (2004)
24. Philos, C.: On the existence of nonoscillatory solutions tending to zero at ∞ for differential equations with positive delay. *Arch. Math. (Basel)* **36**, 168–178 (1981)
25. Tunc, E., Graef, J.: Oscillation results for second order neutral dynamic equations with distributed deviating arguments. *Dyn. Syst. Appl.* **3**, 289–303 (2014)
26. Zhang, C., Li, T., Saker, S.: Oscillation of fourth-order delay differential equations. *J. Math. Sci.* **201**, 296–308 (2014)

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)
