

RESEARCH

Open Access



A detailed study on a new $(2 + 1)$ -dimensional mKdV equation involving the Caputo–Fabrizio time-fractional derivative

K. Hosseini^{1*}, M. Ilie¹, M. Mirzazadeh² and D. Baleanu^{3,4}

*Correspondence:

kamyar_hosseini@yahoo.com

¹Department of Mathematics, Rasht Branch, Islamic Azad University, Rasht, Iran

Full list of author information is available at the end of the article

Abstract

The present article aims to present a comprehensive study on a nonlinear time-fractional model involving the Caputo–Fabrizio (CF) derivative. More explicitly, a new $(2 + 1)$ -dimensional mKdV (2D-mKdV) equation involving the Caputo–Fabrizio time-fractional derivative is considered and an analytic approximation for it is retrieved through a systematic technique, called the homotopy analysis transform (HAT) method. Furthermore, after proving the Lipschitz condition for the kernel $\psi(x, y, t; u)$, the fixed-point theorem is formally utilized to demonstrate the existence and uniqueness of the solution of the new 2D-mKdV equation involving the CF time-fractional derivative. A detailed study finally is carried out to examine the effect of the Caputo–Fabrizio operator on the dynamics of the obtained analytic approximation.

Keywords: $(2 + 1)$ -dimensional mKdV equation; Caputo–Fabrizio time-fractional derivative; Homotopy analysis transform method; Analytic approximation; Fixed-point theorem; Existence and uniqueness of the solution

1 Introduction

The classical KdV equation is a nonlinear partial differential equation to model waves on shallow water surfaces that was established by Korteweg and de Vries in 1895. This exactly solvable model has been the topic of many research works. Nowadays, unique applications of the classical KdV equation have been suggested by many scholars as it can be used to describe long internal waves in a density-stratified ocean, ion-acoustic waves in a plasma, and acoustic waves on a crystal lattice. The mathematical form of the classical KdV equation is given by [1]

$$u_t + \alpha_1 u_{xxx} + \alpha_2 uu_x = 0.$$

© The Author(s) 2020. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

There are different variations of the classical KdV equation, some of those reported in the literature are [2, 3]:

$$\begin{aligned}
 u_t + \alpha_1 u_{xxx} + \alpha_2 u^2 u_x &= 0, & (\text{modified KdV}), \\
 u_t + \alpha_1 u_{xxx} + \alpha_2 u^n u_x &= 0, & (\text{generalized KdV}),
 \end{aligned}$$

and the (2 + 1)-dimensional KdV equation

$$\begin{cases} u_t - u_{xxx} + 3(uv)_x = 0, \\ u_x - v_y = 0. \end{cases}$$

Many efforts have been devoted to studying the KdV-type equations, for example, Wazwaz [2] used different reliable methods to obtain solitons and periodic solutions of the KdV, modified KdV, and generalized KdV equations, and Wang [3] derived lump solutions of the (2 + 1)-dimensional KdV equation by means of an ansatz based on the quadratic functions. Recently, the fractional forms of the KdV-type equations have been explored using a series of systematic methods in [4, 5].

Our aim of this paper is studying a new (2 + 1)-dimensional mKdV equation involving the Caputo–Fabrizio time-fractional derivative as follows:

$${}_0^{\text{CF}}D_t^\alpha u = 6u^2 u_x - 6u^2 u_y + u_{xxx} - u_{yyy} - 3u_{xxy} + 3u_{xyy}, \quad 0 < \alpha \leq 1, \tag{1}$$

through a systematic technique called the homotopy analysis transform method [6–11]. The classical form of the new 2D-mKdV Eq. (1) was first proposed by Wang and Kara [12] in 2019 using the extended Lax pair. Wang and Kara in [12] extracted a group of solitary wave solutions of the new 2D-mKdV equation (its classical form) through the Lie symmetry method.

Recently, nonlinear ODEs/PDEs involving the Caputo–Fabrizio fractional derivative have gained significant attention owing to their potential to describe many complicated physical phenomena. In this respect, Shah et al. [13] analyzed a nonlinear model of dengue fever disease with the CF fractional derivative using the Laplace Adomian decomposition method. Owolabi and Atangana [14] explored a series of nonlinear fractional parabolic differential equations involving the CF derivative thought a numerical scheme. In another work performed by Arshad et al. [15], the CD4⁺ T-cells model of HIV infection with the CF fractional derivative was studied using an effective numerical scheme. Shaikh et al. [16] employed the iterative Laplace transform method to analyze a group of fractional reaction-diffusion equations involving the CF derivative. More articles can be found in [17–45].

The rest of this paper is as follows: In Sect. 2, the Caputo–Fabrizio fractional operators are reviewed in detail. In Sect. 3, the Lipschitz condition for the kernel $\psi(x, y, t; u)$ is proved, then the fixed-point theorem is formally applied to show the existence and uniqueness of the solution of the new 2D-mKdV equation involving the CF time-fractional derivative. In Sect. 4, an analytic approximation for the new 2D-mKdV equation involving the Caputo–Fabrizio time-fractional derivative is acquired using the HAT method. The results of this paper are summarized in the last section.

2 Basic definitions and features

This section presents the basic definitions and features of the Caputo–Fabrizio fractional operators. In this respect, first the Caputo–Fabrizio fractional derivative and integral are defined, then the Laplace transform of the Caputo–Fabrizio fractional derivative is given.

Definition 1 Suppose that $u(t) \in H^1(a, b)$, $b > a$ and $\alpha \in (0, 1]$. Then, the Caputo–Fabrizio fractional derivative of $u(t)$ of order α is given by [17]

$${}^{\text{CF}}D_t^\alpha u(t) = \frac{M(\alpha)}{1 - \alpha} \int_a^t u'(\varepsilon) e^{-\frac{\alpha}{1-\alpha}(t-\varepsilon)} d\varepsilon,$$

where $M(\alpha)$ is a normalized function satisfying $M(0) = M(1) = 1$.

Definition 2 The Caputo–Fabrizio fractional integral of $u(t)$ of order α ($\alpha \in (0, 1]$) is given by [18]

$${}_0^{\text{CF}}I_t^\alpha u(t) = \frac{2(1 - \alpha)}{(2 - \alpha)M(\alpha)} u(t) + \frac{2\alpha}{(2 - \alpha)M(\alpha)} \int_0^t u(\varepsilon) d\varepsilon, \quad t \geq 0.$$

Definition 3 The Laplace transform of ${}^{\text{CF}}D_t^\alpha u(t)$ is given as [17]

$$\mathcal{L}[{}^{\text{CF}}D_t^\alpha u(t)](s) = \frac{s\mathcal{L}[u(t)](s) - u(0)}{s + \alpha(1 - s)},$$

and in the general case, we have

$$\mathcal{L}[{}^{\text{CF}}D_t^{(\alpha+n)} u(t)](s) = \frac{s^{n+1}\mathcal{L}[u(t)](s) - s^n u(0) - s^{n-1} u'(0) - \dots - u^{(n)}(0)}{s + \alpha(1 - s)}.$$

Theorem 1 *The following Lipschitz condition holds for the Caputo–Fabrizio fractional derivative given in Definition 1:*

$$\| {}^{\text{CF}}D_t^\alpha u(t) - {}^{\text{CF}}D_t^\alpha v(t) \| \leq \lambda \| u(t) - v(t) \|.$$

Proof In a similar manner as in [23], it is easy to show that

$$\begin{aligned} \| {}^{\text{CF}}D_t^\alpha u(t) - {}^{\text{CF}}D_t^\alpha v(t) \| &= \left\| \frac{M(\alpha)}{1 - \alpha} \int_a^t u'(\varepsilon) e^{-\frac{\alpha}{1-\alpha}(t-\varepsilon)} d\varepsilon - \frac{M(\alpha)}{1 - \alpha} \int_a^t v'(\varepsilon) e^{-\frac{\alpha}{1-\alpha}(t-\varepsilon)} d\varepsilon \right\| \\ &= \left\| \frac{M(\alpha)}{1 - \alpha} \left(\int_a^t u'(\varepsilon) e^{-\frac{\alpha}{1-\alpha}(t-\varepsilon)} d\varepsilon - \int_a^t v'(\varepsilon) e^{-\frac{\alpha}{1-\alpha}(t-\varepsilon)} d\varepsilon \right) \right\| \\ &= \left\| \frac{M(\alpha)}{1 - \alpha} e^{-\frac{\alpha}{1-\alpha}(t-\varepsilon)} \left(\int_a^t (u'(\varepsilon) - v'(\varepsilon)) d\varepsilon \right) \right\| \\ &\leq \frac{M(\alpha)}{1 - \alpha} \mu e^{-\frac{\alpha}{1-\alpha}(t-\alpha)} \| u(t) - v(t) \| \\ &= \lambda \| u(t) - v(t) \|. \quad \square \end{aligned}$$

3 The model and the existence and uniqueness of its solution

To start, let us consider

$$\psi(x, y, t; u) = 6u^2u_x - 6u^2u_y + u_{xxx} - u_{yyy} - 3u_{xxy} + 3u_{xyy}.$$

This suggests that Eq. (1) can be rewritten as

$${}_0^{CF}D_t^\alpha u(x, y, t) = \psi(x, y, t; u). \tag{2}$$

Applying the CF fractional integral to both sides of Eq. (2) results in

$$u(x, y, t) - u(x, y, 0) = \frac{2(1 - \alpha)}{(2 - \alpha)M(\alpha)} \psi(x, y, t; u) + \frac{2\alpha}{(2 - \alpha)M(\alpha)} \int_0^t \psi(x, y, \varepsilon; u) d\varepsilon. \tag{3}$$

In order to show that the kernel $\psi(x, y, t; u)$ satisfies the Lipschitz condition, we first consider bounded functions $u(x, y, t)$ and $v(x, y, t)$. Using the triangle property of norms, one can find

$$\begin{aligned} & \| \psi(x, y, t; u) - \psi(x, y, t; v) \| \\ &= \| (6u^2u_x - 6v^2v_x) - (6u^2u_y - 6v^2v_y) + (u_{xxx} - v_{xxx}) - (u_{yyy} - v_{yyy}) \\ &\quad - (3u_{xxy} - 3v_{xxy}) + (3u_{xyy} - 3v_{xyy}) \| \\ &= \left\| 2 \frac{\partial}{\partial x} (u^3 - v^3) - 2 \frac{\partial}{\partial y} (u^3 - v^3) + \frac{\partial^3}{\partial x^3} (u - v) - \frac{\partial^3}{\partial y^3} (u - v) \right. \\ &\quad \left. - 3 \frac{\partial^3}{\partial y \partial x^2} (u - v) + 3 \frac{\partial^3}{\partial y^2 \partial x} (u - v) \right\| \\ &\leq 2 \left\| \frac{\partial}{\partial x} (u^3 - v^3) \right\| + 2 \left\| \frac{\partial}{\partial y} (u^3 - v^3) \right\| + \left\| \frac{\partial^3}{\partial x^3} (u - v) \right\| + \left\| \frac{\partial^3}{\partial y^3} (u - v) \right\| \\ &\quad + 3 \left\| \frac{\partial^3}{\partial y \partial x^2} (u - v) \right\| + 3 \left\| \frac{\partial^3}{\partial y^2 \partial x} (u - v) \right\| \\ &\leq 2A \|u^3 - v^3\| + 2B \|u^3 - v^3\| + C \|u - v\| + D \|u - v\| + 3E \|u - v\| + 3F \|u - v\| \\ &\leq (2A + 2B)(\mu^2 + \mu v + v^2) \|u - v\| + C \|u - v\| + D \|u - v\| + 3E \|u - v\| + 3F \|u - v\| \\ &= ((2A + 2B)(\mu^2 + \mu v + v^2) + C + D + 3E + 3F) \|u - v\|, \quad \|u\| \leq \mu, \|v\| \leq v. \end{aligned}$$

Therefore

$$\| \psi(x, y, t; u) - \psi(x, y, t; v) \| \leq \lambda \|u - v\|,$$

which

$$\lambda = ((2A + 2B)(\mu^2 + \mu v + v^2) + C + D + 3E + 3F).$$

This confirms that the Lipschitz condition is satisfied for the kernel $\psi(x, y, t; u)$.

Now, based on the Eq. (3) and the fixed-point theorem, an iterative scheme is established as follows:

$$u_{n+1}(x, y, t) = \frac{2(1 - \alpha)}{(2 - \alpha)M(\alpha)} \psi(x, y, t; u_n) + \frac{2\alpha}{(2 - \alpha)M(\alpha)} \int_0^t \psi(x, y, \varepsilon; u_n) d\varepsilon,$$

where

$$u_0(x, y, t) = u(x, y, 0).$$

It is clear that

$$\begin{aligned} e_n(x, y, t) &= u_n(x, y, t) - u_{n-1}(x, y, t) \\ &= \frac{2(1 - \alpha)}{(2 - \alpha)M(\alpha)} (\psi(x, y, t; u_{n-1}) - \psi(x, y, t; u_{n-2})) \\ &\quad + \frac{2\alpha}{(2 - \alpha)M(\alpha)} \int_0^t (\psi(x, y, \varepsilon; u_{n-1}) - \psi(x, y, \varepsilon; u_{n-2})) d\varepsilon, \end{aligned}$$

and

$$u_n(x, y, t) = \sum_{i=0}^n e_i(x, y, t). \tag{4}$$

Theorem 2 *If the function $u(x, y, t)$ is bounded, then*

$$\|e_n(x, y, t)\| \leq \left(\frac{2(1 - \alpha)}{(2 - \alpha)M(\alpha)} \lambda + \frac{2\alpha}{(2 - \alpha)M(\alpha)} \lambda t \right)^n \|u(x, y, 0)\|.$$

Proof by induction Suppose that $n = 1$. Then, one can write

$$\begin{aligned} \|e_1(x, y, t)\| &= \|u_1(x, y, t) - u_0(x, y, t)\| \\ &\leq \frac{2(1 - \alpha)}{(2 - \alpha)M(\alpha)} \|\psi(x, y, t; u_0) - \psi(x, y, t; u_{-1})\| \\ &\quad + \frac{2\alpha}{(2 - \alpha)M(\alpha)} \int_0^t \|\psi(x, y, \varepsilon; u_0) - \psi(x, y, \varepsilon; u_{-1})\| d\varepsilon \\ &\leq \frac{2(1 - \alpha)}{(2 - \alpha)M(\alpha)} \lambda \|u_0 - u_{-1}\| + \frac{2\alpha}{(2 - \alpha)M(\alpha)} \int_0^t \lambda \|u_0 - u_{-1}\| d\varepsilon \\ &= \frac{2(1 - \alpha)}{(2 - \alpha)M(\alpha)} \lambda \|u(x, y, 0)\| + \frac{2\alpha}{(2 - \alpha)M(\alpha)} \int_0^t \lambda \|u(x, y, 0)\| d\varepsilon \\ &= \frac{2(1 - \alpha)}{(2 - \alpha)M(\alpha)} \lambda \|u(x, y, 0)\| + \frac{2\alpha}{(2 - \alpha)M(\alpha)} \lambda \|u(x, y, 0)\| \int_0^t d\varepsilon \\ &= \frac{2(1 - \alpha)}{(2 - \alpha)M(\alpha)} \lambda \|u(x, y, 0)\| + \frac{2\alpha}{(2 - \alpha)M(\alpha)} \lambda \|u(x, y, 0)\| t \\ &= \left(\frac{2(1 - \alpha)}{(2 - \alpha)M(\alpha)} \lambda + \frac{2\alpha}{(2 - \alpha)M(\alpha)} \lambda t \right) \|u(x, y, 0)\|. \end{aligned}$$

Now, if the relation holds for $n - 1$, namely

$$\|e_{n-1}(x, y, t)\| \leq \left(\frac{2(1-\alpha)}{(2-\alpha)M(\alpha)}\lambda + \frac{2\alpha}{(2-\alpha)M(\alpha)}\lambda t \right)^{n-1} \|u(x, y, 0)\|,$$

then, we will prove that

$$\|e_n(x, y, t)\| \leq \left(\frac{2(1-\alpha)}{(2-\alpha)M(\alpha)}\lambda + \frac{2\alpha}{(2-\alpha)M(\alpha)}\lambda t \right)^n \|u(x, y, 0)\|.$$

To show this, we proceed as follows:

$$\begin{aligned} \|e_n(x, y, t)\| &= \|u_n(x, y, t) - u_{n-1}(x, y, t)\| \\ &\leq \frac{2(1-\alpha)}{(2-\alpha)M(\alpha)} \|\psi(x, y, t; u_{n-1}) - \psi(x, y, t; u_{n-2})\| \\ &\quad + \frac{2\alpha}{(2-\alpha)M(\alpha)} \int_0^t \|\psi(x, y, \varepsilon; u_{n-1}) - \psi(x, y, \varepsilon; u_{n-2})\| d\varepsilon \\ &\leq \frac{2(1-\alpha)}{(2-\alpha)M(\alpha)} \lambda \|u_{n-1} - u_{n-2}\| + \frac{2\alpha}{(2-\alpha)M(\alpha)} \int_0^t \lambda \|u_{n-1} - u_{n-2}\| d\varepsilon \\ &= \frac{2(1-\alpha)}{(2-\alpha)M(\alpha)} \lambda \|e_{n-1}(x, y, t)\| + \frac{2\alpha}{(2-\alpha)M(\alpha)} \int_0^t \lambda \|e_{n-1}(x, y, \varepsilon)\| d\varepsilon \\ &= \left(\frac{2(1-\alpha)}{(2-\alpha)M(\alpha)}\lambda + \frac{2\alpha}{(2-\alpha)M(\alpha)}\lambda t \right)^{n-1} \|u(x, y, 0)\| \\ &\quad \times \left(\frac{2(1-\alpha)}{(2-\alpha)M(\alpha)}\lambda + \frac{2\alpha}{(2-\alpha)M(\alpha)}\lambda \int_0^t d\varepsilon \right) \\ &= \left(\frac{2(1-\alpha)}{(2-\alpha)M(\alpha)}\lambda + \frac{2\alpha}{(2-\alpha)M(\alpha)}\lambda t \right)^{n-1} \\ &\quad \times \left(\frac{2(1-\alpha)}{(2-\alpha)M(\alpha)}\lambda + \frac{2\alpha}{(2-\alpha)M(\alpha)}\lambda t \right) \|u(x, y, 0)\| \\ &= \left(\frac{2(1-\alpha)}{(2-\alpha)M(\alpha)}\lambda + \frac{2\alpha}{(2-\alpha)M(\alpha)}\lambda t \right)^n \|u(x, y, 0)\|. \quad \square \end{aligned}$$

Theorem 3 *If at $t = t_0$ we have*

$$0 \leq \frac{2(1-\alpha)}{(2-\alpha)M(\alpha)}\lambda + \frac{2\alpha}{(2-\alpha)M(\alpha)}\lambda t_0 < 1,$$

then the solution of the new 2D-mKdV equation involving the CF time-fractional derivative exists.

Proof Based on Eq. (4), one can write

$$\begin{aligned} \|u_n(x, y, t)\| &\leq \sum_{i=0}^n \|e_i(x, y, t)\| \\ &\leq \sum_{i=0}^n \left(\frac{2(1-\alpha)}{(2-\alpha)M(\alpha)}\lambda + \frac{2\alpha}{(2-\alpha)M(\alpha)}\lambda t \right)^i \|u(x, y, 0)\|. \end{aligned}$$

For $t = t_0$, one obtains

$$\|u_n(x, y, t)\| \leq \|u(x, y, 0)\| \sum_{i=0}^n \left(\frac{2(1-\alpha)}{(2-\alpha)M(\alpha)}\lambda + \frac{2\alpha}{(2-\alpha)M(\alpha)}\lambda t_0 \right)^i.$$

Consequently,

$$\lim_{n \rightarrow \infty} \|u_n(x, y, t)\| \leq \|u(x, y, 0)\| \sum_{i=0}^{\infty} \left(\frac{2(1-\alpha)}{(2-\alpha)M(\alpha)}\lambda + \frac{2\alpha}{(2-\alpha)M(\alpha)}\lambda t_0 \right)^i.$$

Since

$$0 \leq \frac{2(1-\alpha)}{(2-\alpha)M(\alpha)}\lambda + \frac{2\alpha}{(2-\alpha)M(\alpha)}\lambda t_0 < 1,$$

the above series is convergent, and therefore, $u_n(x, y, t)$ exists and is bounded for any n .

Besides, by assuming

$$R_n(x, y, t) = u(x, y, t) - u_n(x, y, t),$$

one can easily prove that

$$\|R_n(x, y, t)\| \leq \left(\frac{2(1-\alpha)}{(2-\alpha)M(\alpha)}\lambda + \frac{2\alpha}{(2-\alpha)M(\alpha)}\lambda t_0 \right)^{n+1} \mu,$$

and so

$$\lim_{n \rightarrow \infty} \|R_n(x, y, t)\| \leq \lim_{n \rightarrow \infty} \left(\frac{2(1-\alpha)}{(2-\alpha)M(\alpha)}\lambda + \frac{2\alpha}{(2-\alpha)M(\alpha)}\lambda t_0 \right)^{n+1} \mu.$$

But

$$0 \leq \frac{2(1-\alpha)}{(2-\alpha)M(\alpha)}\lambda + \frac{2\alpha}{(2-\alpha)M(\alpha)}\lambda t_0 < 1,$$

thus

$$\lim_{n \rightarrow \infty} \|R_n(x, y, t)\| = 0 \quad \text{or} \quad \lim_{n \rightarrow \infty} u_n(x, y, t) = u(x, y, t). \quad \square$$

Theorem 4 *If at $t = t_0$ we have*

$$0 \leq \frac{2(1-\alpha)}{(2-\alpha)M(\alpha)}\lambda + \frac{2\alpha}{(2-\alpha)M(\alpha)}\lambda t_0 < 1,$$

then the solution of the new 2D- m KdV equation involving the CF time-fractional derivative is unique.

Proof by contradiction To start, let us consider two solutions, say $u(x, y, t)$ and $v(x, y, t)$, for the model (1). One can write

$$u(x, y, t) - v(x, y, t) = \frac{2(1 - \alpha)}{(2 - \alpha)M(\alpha)} (\psi(x, y, t; u) - \psi(x, y, t; v)) + \frac{2\alpha}{(2 - \alpha)M(\alpha)} \int_0^t (\psi(x, y, \varepsilon; u) - \psi(x, y, \varepsilon; v)) d\varepsilon.$$

Consequently,

$$\begin{aligned} \|u(x, y, t) - v(x, y, t)\| &\leq \frac{2(1 - \alpha)}{(2 - \alpha)M(\alpha)} \|\psi(x, y, t; u) - \psi(x, y, t; v)\| \\ &\quad + \frac{2\alpha}{(2 - \alpha)M(\alpha)} \int_0^t \|\psi(x, y, \varepsilon; u) - \psi(x, y, \varepsilon; v)\| d\varepsilon \\ &\leq \frac{2(1 - \alpha)}{(2 - \alpha)M(\alpha)} \lambda \|u - v\| + \frac{2\alpha}{(2 - \alpha)M(\alpha)} \int_0^t \lambda \|u - v\| d\varepsilon \\ &= \left(\frac{2(1 - \alpha)}{(2 - \alpha)M(\alpha)} \lambda + \frac{2\alpha}{(2 - \alpha)M(\alpha)} \lambda \int_0^t d\varepsilon \right) \|u - v\| \\ &= \left(\frac{2(1 - \alpha)}{(2 - \alpha)M(\alpha)} \lambda + \frac{2\alpha}{(2 - \alpha)M(\alpha)} \lambda t \right) \|u - v\|. \end{aligned}$$

But

$$0 \leq \frac{2(1 - \alpha)}{(2 - \alpha)M(\alpha)} \lambda + \frac{2\alpha}{(2 - \alpha)M(\alpha)} \lambda t_0 < 1,$$

therefore

$$\|u(x, y, t) - v(x, y, t)\| = 0,$$

and so, the solution of the new 2D-mKdV equation involving the CF time-fractional derivative is unique. □

4 The new CF time-fractional 2D-mKdV equation and its analytical solutions

In the present section, first soliton solutions of the classical form of the model are extracted using an ansatz method, then the HAT method is used to acquire an analytic approximation for the new CF time-fractional 2D-mKdV equation.

4.1 Soliton solutions of the classical form of the model

To find soliton solutions, a test function is considered as follows:

$$u(x, y, t) = A_0 + A_1 \operatorname{sech}(\alpha_1 x + \alpha_2 y + \alpha_3 t).$$

By substituting the above function into the classical form of the model, we obtain a non-linear algebraic system as follows:

$$6A_0^2\alpha_1 - 6A_0^2\alpha_2 + \alpha_1^3 - 3\alpha_1^2\alpha_2 + 3\alpha_2^2\alpha_1 - \alpha_2^3 - \alpha_3 = 0,$$

$$12A_0A_1\alpha_1 - 12A_0A_1\alpha_2 = 0,$$

$$6A_1^2\alpha_1 - 6A_1^2\alpha_2 - 6\alpha_1^3 + 18\alpha_1^2\alpha_2 - 18\alpha_2^2\alpha_1 + 6\alpha_2^3 = 0,$$

whose solution yields

$$A_0 = 0, A_1 = \mp\alpha_1 \pm \alpha_2, \alpha_3 = \alpha_1^3 - 3\alpha_1^2\alpha_2 + 3\alpha_2^2\alpha_1 - \alpha_2^3.$$

Now, the following soliton solutions to the classical form of the model (1) can be constructed:

$$u_{1,2}(x, y, t) = (\mp\alpha_1 \pm \alpha_2) \operatorname{sech}(\alpha_1x + \alpha_2y + (\alpha_1^3 - 3\alpha_1^2\alpha_2 + 3\alpha_2^2\alpha_1 - \alpha_2^3)t).$$

For $\alpha_1 = 1$ and $\alpha_2 = -1$, the above solitons are reduced to

$$u_{1,2}(x, y, t) = \mp 2 \operatorname{sech}(x - y + 8t) = \mp 4 \frac{e^{x-y+8t}}{1 + e^{2(x-y+8t)}}.$$

4.2 The model and its analytic approximation

To obtain an analytic approximation, we apply the Laplace transform to both sides of Eq. (1). Such an operation results in

$$\begin{aligned} \mathcal{L}[u(x, y, t)] - \frac{u(x, y, 0)}{s} - \left(\frac{s + \alpha(1 - s)}{s}\right) \mathcal{L}[6u^2(x, y, t)u_x(x, y, t) \\ - 6u^2(x, y, t)u_y(x, y, t) + u_{xxx}(x, y, t) - u_{yyy}(x, y, t) \\ - 3u_{xxy}(x, y, t) + 3u_{xyy}(x, y, t)] = 0. \end{aligned} \tag{5}$$

Based on Eq. (5), the nonlinear operator can be defined as

$$\begin{aligned} \Omega[\phi(x, y, t; p)] &= \mathcal{L}[\phi(x, y, t; p)] - \frac{u_0(x, y, t)}{s} \\ &\quad - \left(\frac{s + \alpha(1 - s)}{s}\right) \mathcal{L}[6\phi^2(x, y, t; p)\phi_x(x, y, t; p) \\ &\quad - 6\phi^2(x, y, t; p)\phi_y(x, y, t; p) + \phi_{xxx}(x, y, t; p) - \phi_{yyy}(x, y, t; p) \\ &\quad - 3\phi_{xxy}(x, y, t; p) + 3\phi_{xyy}(x, y, t; p)] \\ &= 0. \end{aligned}$$

Now, the following m th order deformation equation is considered:

$$\mathcal{L}[u_m(x, y, t) - \chi_m u_{m-1}(x, y, t)] = h\mathcal{R}_m(\bar{u}_{m-1}),$$

where

$$\mathcal{R}_m(\bar{u}_{m-1}) = \frac{1}{(m - 1)!} \frac{\partial^{m-1} \Omega[\phi(x, y, t; p)]}{\partial p^{m-1}} \Big|_{p=0},$$

and

$$\chi_m = \begin{cases} 0 & m \leq 1, \\ 1 & m > 1. \end{cases}$$

It is worth mentioning that by selecting

$$u(x, y, 0) = -4 \frac{e^{x-y}}{1 + e^{2(x-y)}}$$

and solving the resulting equations, one obtains

$$\begin{aligned} u_0(x, y, t) &= -4 \frac{e^{x-y}}{1 + e^{2(x-y)}}, \\ u_1(x, y, t) &= -32 \frac{he^{x-y}(\alpha e^{2(x-y)}t - \alpha e^{2(x-y)} + e^{2(x-y)} - \alpha t + \alpha - 1)}{(e^{2(x-y)} + 1)^2}, \\ &\vdots \end{aligned}$$

Therefore, the series solution of the new CF time-fractional 2D-mKdV equation is derived as

$$u(x, y, t) = -4 \frac{e^{x-y}}{1 + e^{2(x-y)}} - 32 \frac{he^{x-y}(\alpha e^{2(x-y)}t - \alpha e^{2(x-y)} + e^{2(x-y)} - \alpha t + \alpha - 1)}{(e^{2(x-y)} + 1)^2} + \dots$$

It is noteworthy that for $\alpha = 1$ and $h = -1$, the above series converges to the following exact solution:

$$u(x, y, t) = -4 \frac{e^{x-y+8t}}{1 + e^{2(x-y+8t)}}.$$

Figure 1 presents the 3-order approximation of the new CF time-fractional 2D-mKdV equation for $\alpha = 1, 0.99$, and 0.98 against the exact solution. From this figure, a full agreement between the 3-order approximation (when $\alpha = 1$) and the exact solution is obviously observed. The absolute error of the 3-order approximation (when $\alpha = 1$) and the exact solution has been presented in Table 1. The results confirm the efficiency of the HAT method




Figure 1 The 3th order approximation against the exact solution (a) $y = 0.5, t = 0.01, h = -1$, and $\alpha = 1, 0.99$, and 0.98 ; (b) $x = 0.5, t = 0.01, h = -1$, and $\alpha = 1, 0.99$, and 0.98

Table 1 The absolute error of the 3-order approximation and the exact solution

t	Absolute error when $y = 0.5, t = 0.01, h = -1, \text{ and } \alpha = 1$	Absolute error when $x = 0.5, t = 0.01, h = -1, \text{ and } \alpha = 1$
0	9.921881×10^{-7}	2.607245×10^{-7}
0.5	1.702236×10^{-5}	1.702236×10^{-5}
1	2.607245×10^{-7}	9.921881×10^{-7}
1.5	6.913822×10^{-6}	7.076190×10^{-6}
2	2.534162×10^{-6}	2.783063×10^{-6}
2.5	2.305783×10^{-7}	3.045477×10^{-7}
3	2.741436×10^{-7}	2.651408×10^{-7}
3.5	2.706178×10^{-7}	2.751423×10^{-7}
4	1.884674×10^{-7}	1.936185×10^{-7}
4.5	1.198279×10^{-7}	1.235044×10^{-7}
5	7.391757×10^{-8}	7.627198×10^{-8}




Figure 2 (a) The exact solution for $t = 0.01$ against (b, c, d) the 3-order approximation when (b) $t = 0.01, h = -1, \text{ and } \alpha = 1$; (c) $t = 0.01, h = -1, \text{ and } \alpha = 0.99$; (d) $t = 0.01, h = -1, \text{ and } \alpha = 0.98$

in deriving an analytic approximation with high accuracy. Finally, Fig. 2 shows the 3D plots of the exact solution and the 3-order approximation for $\alpha = 1, 0.99, \text{ and } 0.98$.

It is believed that the analytic approximation given by the HAT method can precisely predict the dynamics of the dark soliton solution of the new 2D-mKdV equation involving the CF time-fractional derivative.

5 Conclusion

A thorough study on a nonlinear model involving the Caputo–Fabrizio time-fractional derivative was carried out successfully in the current paper. In this respect, a new 2D-mKdV equation designed with the CF time-fractional derivative was considered, and an analytic approximation for it was formally derived using a systematic approach, named the homotopy analysis transform method. The existence and uniqueness of the solution of the new 2D-mKdV equation involving the Caputo–Fabrizio time-fractional derivative

were studied by proving the Lipschitz condition for the kernel $\psi(x, y, t; u)$ and applying the fixed-point theorem. A detailed study was finally accomplished to investigate the effect of the Caputo–Fabrizio operator on the dynamics of the obtained analytic approximation. The results presented herein confirm the efficiency of the HAT method in deriving an analytic approximation with high accuracy for nonlinear models involving the Caputo–Fabrizio time-fractional derivative. Our future work is to obtain an analytic approximation of the new 2D-mKdV equation with the Atangana–Baleanu time-fractional derivative and study the effect of the Atangana–Baleanu operator on the dynamics of the approximate solution.

Acknowledgements

Not applicable.

Funding

There is no funding for this work.

Availability of data and materials

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors jointly worked on the results and they read and approved the final manuscript.

Author details

¹Department of Mathematics, Rasht Branch, Islamic Azad University, Rasht, Iran. ²Department of Engineering Sciences, Faculty of Technology and Engineering, East of Guilan, University of Guilan, P.C. 44891-63157, Rudsar-Vajargah, Iran.

³Department of Mathematics, Faculty of Arts and Sciences, Cankaya University, Ankara 06530, Turkey. ⁴Institute of Space Sciences, Magurele-Bucharest, R 76900, Romania.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 2 April 2020 Accepted: 21 June 2020 Published online: 06 July 2020

References

1. https://en.wikipedia.org/wiki/Korteweg-de_Vries_equation
2. Wazwaz, A.M.: New sets of solitary wave solutions to the KdV, mKdV, and the generalized KdV equations. *Commun. Nonlinear Sci. Numer. Simul.* **13**, 331–339 (2008)
3. Wang, C.: Spatiotemporal deformation of lump solution to (2 + 1)-dimensional KdV equation. *Nonlinear Dyn.* **84**, 697–702 (2016)
4. Sontakke, B.R., Shaikh, A.: The new iterative method for approximate solutions of time fractional KdV, $K(2, 2)$, Burgers and cubic Boussinesq equations. *Asian Res. J. Math.* **1**, 1–10 (2016)
5. Sontakke, B.R., Shaikh, A., Nisar, K.S.: Approximate solutions of a generalized Hirota–Satsuma coupled KdV and a coupled mKdV systems with time fractional derivatives. *Malaysian J. Math. Sci.* **12**, 175–196 (2018)
6. Saad, K.M., AL-Shareef, E.H.F., Alomari, A.K., Baleanu, D., Gómez-Aguilar, J.F.: On exact solutions for time-fractional Korteweg–de Vries and Korteweg–de Vries–Burger's equations using homotopy analysis transform method. *Chin. J. Phys.* **63**, 149–162 (2020)
7. Bhattar, S., Mathur, A., Kumar, D., Nisar, K.S., Singh, J.: Fractional modified Kawahara equation with Mittag-Leffler law. *Chaos Solitons Fractals* **131**, 109508 (2020)
8. Kumar, D., Singh, J., Kumar, S., Sushil: Numerical computation of Klein–Gordon equations arising in quantum field theory by using homotopy analysis transform method. *Alex. Eng. J.* **53**, 469–474 (2014)
9. Yépez-Martínez, H., Gómez-Aguilar, J.F.: A new modified definition of Caputo–Fabrizio fractional-order derivative and their applications to the multi step homotopy analysis method (MHAM). *J. Comput. Appl. Math.* **346**, 247–260 (2019)
10. Baleanu, D., Mohammadi, H., Rezapour, S.: Analysis of the model of HIV-1 infection of CD4⁺ T-cell with a new approach of fractional derivative. *Adv. Differ. Equ.* **2020**, 71 (2020)
11. Baleanu, D., Jajarmi, A., Mohammadi, H., Rezapour, S.: A new study on the mathematical modelling of human liver with Caputo–Fabrizio fractional derivative. *Chaos Solitons Fractals* **134**, 109705 (2020)
12. Wang, G., Kara, A.H.: A (2 + 1)-dimensional KdV equation and mKdV equation: symmetries, group invariant solutions and conservation laws. *Phys. Lett. A* **383**, 728–731 (2019)
13. Shah, K., Jarad, F., Abdeljawad, T.: On a nonlinear fractional order model of Dengue fever disease under Caputo–Fabrizio derivative. *Alex. Eng. J.* (2020). <https://doi.org/10.1016/j.aej.2020.02.022>
14. Owolabi, K.M., Atangana, A.: Numerical approximation of nonlinear fractional parabolic differential equations with Caputo–Fabrizio derivative in Riemann–Liouville sense. *Chaos Solitons Fractals* **99**, 171–179 (2017)

15. Arshad, S., Defterli, O., Baleanu, D.: A second order accurate approximation for fractional derivatives with singular and non-singular kernel applied to a HIV model. *Appl. Math. Comput.* **374**, 125061 (2020)
16. Shaikh, A., Tassaddiq, A., Nisar, K.S., Baleanu, D.: Analysis of differential equations involving Caputo–Fabrizio fractional operator and its applications to reaction-diffusion equations. *Adv. Differ. Equ.* **2019**, 178 (2019)
17. Caputo, M., Fabrizio, M.: A new definition of fractional derivative without singular kernel. *Prog. Fract. Differ. Appl.* **1**, 73–85 (2015)
18. Losada, J., Nieto, J.J.: Properties of a new fractional derivative without singular kernel. *Prog. Fract. Differ. Appl.* **1**, 87–92 (2015)
19. Alizadeh, S., Baleanu, D., Rezapour, S.: Analyzing transient response of the parallel RCL circuit by using the Caputo–Fabrizio fractional derivative. *Adv. Differ. Equ.* **2020**, 55 (2020)
20. Algahtani, O.J.J.: Comparing the Atangana–Baleanu and Caputo–Fabrizio derivative with fractional order: Allen Cahn model. *Chaos Solitons Fractals* **89**, 552–559 (2016)
21. Ali, F., Ali, F., Sheikh, N.A., Khan, I., Nisar, K.S.: Caputo–Fabrizio fractional derivatives modeling of transient MHD Brinkman nanoliquid: applications in food technology. *Chaos Solitons Fractals* **131**, 109489 (2020)
22. Shaikh, A.S., Nisar, K.S.: Transmission dynamics of fractional order Typhoid fever model using Caputo–Fabrizio operator. *Chaos Solitons Fractals* **128**, 355–365 (2019)
23. Atangana, A., Baleanu, D.: New fractional derivative with nonlocal and non-singular kernel, theory and application to heat transfer model. *Therm. Sci.* **20**, 763–769 (2016)
24. Inc, M., Yusuf, A., Aliyu, A.I., Baleanu, D.: Investigation of the logarithmic-KdV equation involving Mittag-Leffler type kernel with Atangana–Baleanu derivative. *Physica A* **506**, 520–531 (2018)
25. Korpınar, Z., Inc, M., Bayram, M.: Theory and application for the system of fractional Burger equations with Mittag-Leffler kernel. *Appl. Math. Comput.* **367**, 124781 (2020)
26. Morales-Delgado, V.F., Gómez-Aguilar, J.F., Saad, K.M., Khan, M.A., Agarwal, P.: Analytic solution for oxygen diffusion from capillary to tissues involving external force effects: a fractional calculus approach. *Physica A* **523**, 48–65 (2019)
27. Saad, K.M., Gómez-Aguilar, J.F.: Analysis of reaction-diffusion system via a new fractional derivative with non-singular kernel. *Physica A* **509**, 703–716 (2018)
28. Veerasha, P., Prakasha, D.G.: Solution for fractional generalized Zakharov equations with Mittag-Leffler function. *Results Eng.* **5**, 100085 (2020)
29. Gao, W., Veerasha, P., Prakasha, D.G., Baskonus, H.M., Yel, G.: A powerful approach for fractional Drinfeld–Sokolov–Wilson equation with Mittag-Leffler law. *Alex. Eng. J.* **58**, 1301–1311 (2019)
30. Owolabi, K.M., Atangana, A.: Computational study of multi-species fractional reaction-diffusion system with ABC operator. *Chaos Solitons Fractals* **128**, 280–289 (2019)
31. Heydari, M.H., Atangana, A., Avazzadeh, Z., Mahmoudi, M.R.: An operational matrix method for nonlinear variable-order time fractional reaction-diffusion equation involving Mittag-Leffler kernel. *Eur. Phys. J. Plus* **135**, 237 (2020)
32. Jajarmi, A., Baleanu, D.: A new fractional analysis on the interaction of HIV with CD4 + T-cells. *Chaos Solitons Fractals* **113**, 221–229 (2018)
33. Yusuf, A., Inc, M., Aliyu, A.I., Baleanu, D.: Efficiency of the new fractional derivative with nonsingular Mittag-Leffler kernel to some nonlinear partial differential equations. *Chaos Solitons Fractals* **116**, 220–226 (2018)
34. Liao, S.: *Beyond Perturbation: Introduction to the Homotopy Analysis Method*. CRC Press, Boca Raton (2004)
35. Samadani, F., Moradweysi, P., Ansari, R., Hosseini, K., Darvizeh, A.: Application of homotopy analysis method for the pull-in and nonlinear vibration analysis of nanobeams using a nonlocal Euler–Bernoulli beam model. *Z. Naturforsch. A* **72**, 1093–1104 (2017)
36. Zabihi, A., Ansari, R., Hosseini, K., Samadani, F., Torabi, J.: Nonlinear pull-in instability of rectangular nanoplates based on the positive and negative second-order strain gradient theories with various edge supports. *Z. Naturforsch. A, J. Phys. Sci.* **75**, 317–331 (2020)
37. Torabi, J., Ansari, R., Zabihi, A., Hosseini, K.: Dynamic and pull-in instability analyses of functionally graded nanoplates via nonlocal strain gradient theory. *Mech. Based Des. Struct. Mach.* (2020). <https://doi.org/10.1080/15397734.2020.1721298>
38. Hosseini, S.M.J., Ansari, R., Torabi, J., Zabihi, A., Hosseini, K.: Nonlocal strain gradient pull-in study of nanobeams considering various boundary conditions. *Iran. J. Sci. Technol., Trans. Mech. Eng.* (2020). <https://doi.org/10.1007/s40997-020-00365-6>
39. Sontakke, B.R., Shaikh, A.: Approximate solutions of time fractional Kawahara and modified Kawahara equations by fractional complex transform. *Commun. Numer. Anal.* **2016**, 218–229 (2016)
40. Sontakke, B.R., Shaikh, A., Nisar, K.S.: Existence and uniqueness of integrable solutions of fractional order initial value equations. *J. Math. Model.* **6**, 137–148 (2018)
41. Nisar, K.S.: Generalized Mittag-Leffler type function: fractional integrations and application to fractional kinetic equations. *Front. Phys.* **8**, 33 (2020)
42. Khan, O., Khan, N., Baleanu, D., Nisar, K.S.: Computable solution of fractional kinetic equations using Mathieu-type series. *Adv. Differ. Equ.* **2019**, 234 (2019)
43. Khan, O., Khan, N., Nisar, K.S., Saif, M., Baleanu, D.: Fractional calculus of a product of multivariable Srivastava polynomial and multi-index Bessel function in the kernel F_3 . *AIMS Math.* **5**, 1462–1475 (2020)
44. Jumani, T.A., Mustafa, M.W., Hussain, Z., Rasid, M.M., Saeed, M.S., Memon, M.M., Khan, I., Nisar, K.S.: Jaya optimization algorithm for transient response and stability enhancement of a fractional-order PID based automatic voltage regulator system. *Alex. Eng. J.* (2020). <https://doi.org/10.1016/j.aej.2020.03.005>
45. Yusuf, A., Inc, M., Aliyu, A.I., Baleanu, D.: Conservation laws, soliton-like and stability analysis for the time fractional dispersive long-wave equation. *Adv. Differ. Equ.* **2018**, 319 (2018)