

RESEARCH

Open Access



Hyers–Ulam stability of non-autonomous and nonsingular delay difference equations

Gul Rahmat¹, Atta Ullah¹, Aziz Ur Rahman¹, Muhammad Sarwar^{2*}, Thabet Abdeljawad^{3,4,5*} and Aiman Mukheimer³

*Correspondence:

sarwarswati@gmail.com;

tabdeljawad@psu.edu.sa

²Department of Mathematics,
University of Malakand, Chakdara,
Dir(L), Pakistan

³Department of Mathematics and
Sciences, Prince Sultan University,
P.O. Box 66833, Riyadh 11586, Saudi
Arabia

Full list of author information is
available at the end of the article

Abstract

In this paper, we study the uniqueness and existence of the solution of a non-autonomous and nonsingular delay difference equation using the well-known principle of contraction from fixed point theory. Furthermore, we study the Hyers–Ulam stability of the given system on a bounded discrete interval and then on an unbounded interval. An example is also given at the end to illustrate the theoretical work.

MSC: 34A08; 35R11

Keywords: Hyer–Ulam; Stability; Difference equations

1 Introduction

In most of the fields of science like mathematics, statistics, economics, and engineering, people take the number of samples as a discrete form, not in a continuous form. In real life, some model such as [1–4] are used to analyze the real life problems from mechanics and biomathematics by means of partial differential equations, but some models like cobweb [5] and national income model [6] can be described by difference equations. People study difference equations because of their many applications in population models and information transmission. S.A. Kuruklis [7] and J.S. Yu [8] studied the asymptotic behavior of the variable type delay difference equation. W. Kosmala [9] provided a good insight and discussed the behavior of solution of the difference equation of the type $U_{k+1} = (A + U_{k-1})/(BU_k + U_{k-1})$. Zheng-Fan Liu [10] designed the exponential behavior of switch discrete-time delay system. Marwen Kermani [11] discussed the stability techniques about the switched nonlinear time-delay difference equations. Yuanyuan Liu [12] described the stability techniques of a higher-order difference system. The stability of higher-order rational difference systems was studied by A. Khaliq [13].

A difference system has a lot of qualitative properties, among them stability is a very useful property. It is a vital part for a system to work appropriately. There are many types of stability, but today's general interest that leads people to want to know more is about Hyers–Ulam stability. Ulam [14], in 1940, initially studied the theory of Hyers–Ulam stability. When he lectured a seminar, he put out some problems related to the group ho-

© The Author(s) 2021. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

homomorphisms stability. A year later, Hyers [15] answered brightly to the problem by considering that groups are Banach spaces, specified by Hyers–Ulam stability. Rassias [16], in 1978, gave an outstanding general approach of the Hyers–Ulam stability, specified by Hyers–Ulam–Rassias stability. In particular, he also extended the same concept to Cauchy difference equation. This idea was then extended to differential equations by Obloza [17]. Later on, this stability of difference equations was proved by Jung [18] and Khan *et al.* [19].

Delay systems can have a lot of uses in the characterization of the evolution process in automatic engine, physiology system, and control theory. Khusainov and Shuklin [20] solved the linear autonomous delay-time system with commutable matrices. Diblik and Khusainov [21] gave the description about the solutions of a discrete delayed system using the idea [20]. Then Wang *et al.* [22] studied relative controllability and exponential stability of nonsingular systems. The consequences about Ulam stability of nonsingular delay differential equation having first order were shown by A. Zada *et al.* [23]. Recently, some results on Ulam type stability of a first-order nonlinear impulsive time varying delay dynamic system was discussed by S.O. Shah *et al.* in [24, 25].

In this study, we discuss the Hyers–Ulam stability of nonsingular delay difference system of the form:

$$\begin{cases} \mathbb{A}\mathbf{G}_{n+1} = \mathbf{M}\mathbf{G}_n + \mathcal{N}\mathbf{G}_{n-k} + \mathbb{F}(n, \mathbf{G}_{n-k}), & n \geq 0, k \geq 0, \\ \mathbf{G}_n = \phi_n, & -k \leq n \leq 0, \end{cases} \quad (1.1)$$

where the constant matrices \mathbb{A} , \mathbf{M} , and \mathcal{N} are commutable having order $n \times n$. The matrix A is invertible and $\phi \in \mathcal{B}(\mathbb{Z}_+, X)$, where $\mathcal{B}(\mathbb{Z}_+, X)$ is the space of bounded sequences, also $\mathbb{F} \in \mathcal{CS}(\mathbb{Z}_+ \times X, X)$, the space of all convergent sequences, where $I = \{-k, -k+1, \dots, 0\}$, $\mathbb{Z}_+ = \{0, 1, 2, \dots\}$ and $X = \mathcal{R}^n$. Such work in the continuous case is given in [26].

2 Preliminaries

In this portion, we discuss some definitions and basic concepts which are useful for establishing the main work. We will use the notation \mathcal{R} , \mathcal{R}^+ , \mathcal{Z}_+ for the real numbers, non-negative real numbers, all nonnegative integers, and the space of all n -tuples of \mathcal{R} is denoted by \mathcal{R}^n . The set $J = \{0, 1, \dots, k\}$ is the subset of \mathcal{Z} and $X = \mathcal{R}^n$, the space of all bounded and convergent sequences from J to X is represented by $\mathcal{CS}(J, X)$ with the norm

$$\|\mathbf{G}\|_{cs} = \left\{ \sup_{n \in J} \|\mathbf{G}(n)\| \text{ for all } \mathbf{G} \in \mathcal{CS}(J, X) \right\}.$$

Besides, we define $\mathcal{C}^1(J, X) = \{\mathbf{G} \in \mathcal{C}(J, X); \mathbf{G}' \in \mathcal{C}^1(J, X)\}$.

Lemma 2.1 *The nonsingular delay difference system*

$$\begin{cases} \mathbf{A}\mathbf{G}_{n+1} = \mathbf{M}\mathbf{G}_n + \mathbf{N}\mathbf{G}_{n-k} + \mathbb{F}(n, \mathbf{G}_{n-k}), & n \geq 0, k \geq 0, \\ \mathbf{G}_n = \Phi_n, & -k \leq n \leq 0, \end{cases}$$

has the solution

$$\mathbf{G}_n = M^n A^{-n} \Phi_0 + M^{n-1} A^{-n} \sum_{i=0}^k M^{-i} A^i (N \Phi_{i-k} + \mathbb{F}(i, \Phi_{i-k}))$$

$$+ M^{n-1} A^{-n} \sum_{i=k+1}^n M^{-i} A^i (N \mathbf{G}_{i-k} + \mathbb{F}(i, \mathbf{G}_{i-k})),$$

where $MN = NM$, $NA = AN$, and $MA = AM$.

The proof can be easily obtained by successively putting the values of $n \in \{-k, -k+1, \dots\}$.

Definition 2.1 The solution of system (1.1) will be exponentially stable if there exist positive real numbers λ_1 and λ_2 such that

$$\|\mathbf{G}_n\| \leq \lambda_1 e^{-\lambda_2 n}, \quad \forall n \geq 0.$$

Definition 2.2 For a positive real number ϵ , the sequence $\boldsymbol{\psi}_n$ is said to be an ϵ -approximate solution of (1.1) if

$$\begin{cases} \|A\boldsymbol{\psi}_{n+1} - M\boldsymbol{\psi}_n - N\boldsymbol{\psi}_{n-k} - \mathbb{F}(n, \boldsymbol{\psi}_{n-k})\| \leq \epsilon, & n \geq 0, k \geq 0, \\ \|\boldsymbol{\psi}_n - \phi_n\| \leq \epsilon, & -k \leq n \leq 0. \end{cases} \quad (2.1)$$

Definition 2.3 System (1.1) will be Hyers–Ulam stable if, for every ϵ -approximate solution $\boldsymbol{\psi}_n$ of system (1.1), there will be an exact solution \mathbf{Y}_n of (1.1) and a nonnegative real number \mathbf{K} such that

$$\|\mathbf{Y}_n - \boldsymbol{\psi}_n\| \leq \mathbf{K}\epsilon, \quad n \in J.$$

Definition 2.4 A function $\|\cdot\|_\beta : \mathbb{V} \rightarrow [0, \infty)$ is called β -norm, with $0 < \beta \leq 1$, where \mathbb{V} is a vector space over field \mathbf{K} , if the function satisfies the following properties:

- (1) $\|\mathcal{H}\|_\beta = 0$ if and only if $\mathcal{G} = 0$;
- (2) $\|\kappa \mathcal{H}\|_\beta = |\kappa|^\beta \|\mathcal{H}\|_\beta$ for each $\kappa \in \mathbf{K}$ and $\mathcal{H} \in \mathbb{V}$;
- (3) $\|\mathcal{H} + \mathcal{H}_1\|_\beta \leq \|\mathcal{H}\|_\beta + \|\mathcal{H}_1\|_\beta$ for all $\mathcal{H}, \mathcal{H}_1 \in \mathbb{V}$.

And $(\mathbb{V}, \|\cdot\|_\beta)$ is said to be a β -norm space.

Lemma 2.2 If z_n and g_n are nonnegative sequences and $a \geq 0$, which satisfies the inequality

$$\|z_n\| \leq a + \sum_{i=0}^n \|g_i\| \|z_i\|, \quad n \geq 0,$$

then

$$\|z_n\| \leq a \exp \left(\sum_{i=0}^n \|z_i\| \right).$$

Remark 2.1 It is clear from (2.1) that $\mathbf{Y} \in C^1(J, X)$ satisfies (2.1) if and only if there exists $f \in \mathcal{CS}(J, X)$ satisfying

$$\begin{cases} \|f_n\| \leq \epsilon, & n \in J, \\ \mathbb{A}\mathcal{Y}_{n+1} = \mathbf{M}\mathcal{Y}_n + \mathcal{N}\mathcal{Y}_{n-k} + \mathbb{F}(n, \mathcal{Y}_{n-k}) + f_n, & n \in \mathbb{Z}_+, \\ \mathcal{Y}_n = \phi_n, & -k \leq n \leq 0. \end{cases}$$

3 Existence result

To describe the existence result of the system given by (1.1), we have the following assumptions which will be needed:

Λ_1 : The linear system $A\mathbf{G}_{n+1} = M\mathbf{G}_n + N\mathbf{G}_{n-k}$ is well modeled.

Λ_2 : The continuous function $\mathcal{H} : J \times X \rightarrow X$ satisfies the Caratheodory condition

$$\|\mathcal{H}(n, f) - \mathcal{H}(n, f')\| \leq \mathcal{K}\|f - f'\|, \quad \mathcal{K} \geq 0,$$

for every $f, f' \in X$.

Λ_3 : $\mathcal{M}^{n-1}\mathcal{A}^{-n}(N + \mathcal{K})L < 1$.

Theorem 1 *If assumptions Λ_1 – Λ_3 hold, then system (1.1) has the unique solution $\mathbf{G} \in \mathcal{B}(J, X)$.*

Proof Define $T : \mathcal{B}(J, X) \rightarrow \mathcal{B}(J, X)$ by

$$\begin{aligned} (T\mathbf{G})_n &= \mathbf{M}^n \mathbb{A}^{-n} \phi_0 + \mathbf{M}^{n-1} \mathbb{A}^{-n} \sum_{i=0}^k \mathbf{M}^{-i} \mathbb{A}^i (\mathcal{N} \phi_{j-k} + \mathbb{F}(i, \phi_{i-k})) \\ &\quad + \mathbf{M}^{n-1} \mathbb{A}^{-n} \sum_{i=k+1}^n \mathbf{M}^{-i} \mathbb{A}^i (\mathcal{N} \mathbf{G}_{i-k} + \mathbb{F}(i, \mathbf{G}_{i-k})). \end{aligned}$$

Now, for any $\mathbf{G}, \mathbf{G}' \in \mathcal{B}(J, X)$, we have

$$\begin{aligned} \|(T\mathbf{G})_n - (T\mathbf{G}')_n\| &= \left\| \mathbf{M}^n \mathbb{A}^{-n} \phi_0 + \mathbf{M}^{n-1} \mathbb{A}^{-n} \sum_{i=0}^k \mathbf{M}^{-i} \mathbb{A}^i (\mathcal{N} \phi_{i-k} + \mathbb{F}(i, \phi_{i-k})) \right. \\ &\quad + \mathbf{M}^{n-1} \mathbb{A}^{-n} \sum_{i=k+1}^n \mathbf{M}^{-i} \mathbb{A}^i (\mathcal{N} \mathbf{G}_{i-k} + \mathbb{F}(i, \mathbf{G}_{i-k})) \\ &\quad - \mathbf{M}^{n-1} \mathbb{A}^{-n} \phi_0 - \mathbf{M}^{n-1} \mathbb{A}^{-n} \sum_{i=k+1}^n \mathbf{M}^{-i} \mathbb{A}^i (\mathcal{N} \phi_{i-k} + \mathbb{F}(i, \phi_{i-k})) \\ &\quad \left. - \mathbf{M}^{n-1} \mathbb{A}^{-n} \sum_{i=k+1}^n \mathbf{M}^{-i} \mathbb{A}^i (\mathcal{N} \mathbf{G}'_{i-k} + \mathbb{F}(i, \mathbf{G}'_{i-k})) \right\|. \end{aligned}$$

This implies that

$$\begin{aligned} \|(T\mathbf{G})_n - (T\mathbf{G}')_n\| &\leq \|\mathbf{M}^{n-1}\| \|\mathbb{A}^{-n}\| \sum_{i=k+1}^n \|\mathbf{M}^{-i}\| \|\mathbb{A}^i\| (\|\mathcal{N} \mathbf{G}_{i-k} - \mathcal{N} \mathbf{G}'_{i-k}\| \\ &\quad + \|\mathbb{F}(i, \mathbf{G}_{i-k}) - \mathbb{F}(i, \mathbf{G}'_{i-k})\|) \\ &\leq \|\mathbf{M}^{n-1}\| \|\mathbb{A}^{-n}\| \sum_{i=k+1}^n \mathbf{M}^{-i} \mathbb{A}^i (\|\mathcal{N} \mathbf{G}_{i-k} - \mathcal{N} \mathbf{G}'_{i-k}\| \\ &\quad + \mathcal{K} \|\mathbf{G}_{i-k} - \mathbf{G}'_{i-k}\|) \\ &= \|\mathbf{M}^{n-1}\| \|\mathbb{A}^{-n}\| (\mathcal{N} + \mathcal{K}) \sum_{i=k+1}^n \mathbf{M}^{-i} \mathbb{A}^i \|\mathbf{G}_{i-k} - \mathbf{G}'_{i-k}\| \\ &\leq \|\mathbf{M}^{n-1}\| \|\mathcal{A}^{-n}\| (\mathcal{N} + \mathcal{K}) L \|\mathbf{G} - \mathbf{G}'\|_{\mathcal{B}}. \end{aligned}$$

Thus, T is a contraction if $\|\mathbf{M}^n\| \|\mathbb{A}^{-n}\|(\mathcal{N} + \mathcal{K})L < 1$, so (by BCP) it has a unique fixed point and will be the solution of system (1.1). \square

4 Hyers–Ulam stability on bounded discrete interval

To describe the Hyers–Ulam stability of system (1.1) over a bounded discrete interval, we have to put some assumptions:

Λ_1 : The linear system $A\mathbf{G}_{n+1} = M\mathbf{G}_n + N\mathbf{G}_{n-k}$ is well posed.

Λ_2 : The map $F: J \times X \rightarrow X$ satisfies the Caratheodory condition

$$\|F(n, \vartheta) - F(n, \vartheta')\| \leq K \|\vartheta - \vartheta'\|$$

for some $K \geq 0$ and for all $\vartheta, \vartheta' \in \mathcal{B}(J, X)$.

Λ_3 : There exists nondecreasing $\varphi_n \in \mathcal{B}(J, X)$ with a constant η such that

$$\sum_{r=1}^{n-k} \phi_r \leq \eta \varphi_n \quad \text{for each } n \in J.$$

Theorem 2 *If Λ_1 , Λ_2 , and Λ_3 along with (2.1) and Remark 2.1 hold, then system (1.1) is Hyers–Ulam stable.*

Proof The solution of delay difference equation

$$\begin{cases} A\mathbf{G}_{n+1} = M\mathbf{G}_n + N\mathbf{G}_{n-k} + \mathbb{F}(n, \mathbf{G}_{n-k}), & n \geq 0, k \geq 0, \\ \mathbf{G}_n = \Psi_n, & -k \leq n \leq 0, \end{cases}$$

is

$$\begin{aligned} \mathbf{G}_n &= M^n A^{-n} \Psi_0 + M^{n-1} A^{-n} \sum_{i=0}^k M^{-i} A^i (N \Psi_{i-k} + \mathbb{F}(i, \Psi_{i-k})) \\ &\quad + M^{n-1} A^{-n} \sum_{i=k+1}^n M^{-i} A^i (N \mathbf{G}_{i-k} + \mathbb{F}(i, \mathbf{G}_{i-k})). \end{aligned}$$

From Remark 2.1 the solution of

$$\begin{cases} A\mathbb{Y}_{n+1} = M\mathbb{Y}_n + N\mathbb{Y}_{n-k} + \mathbb{F}(n, \mathbb{Y}_{n-k}) + \mathbf{f}_n, & n \geq 0, k \geq 0, \\ \mathbb{Y}_n = \Psi_n, & -k \leq n \leq 0, \end{cases}$$

is

$$\begin{aligned} \mathbb{Y}_n &= M^n A^{-n} \Psi_0 + M^{n-1} A^{-n} \sum_{i=0}^k M^{-i} A^i (N \Psi_{i-k} + \mathbb{F}(i, \Psi_{i-k})) \\ &\quad + M^{n-1} A^{-n} \sum_{i=k+1}^n M^{-i} A^i (N \mathbb{Y}_{i-k} + \mathbb{F}(i, \mathbb{Y}_{i-k}) + \mathbf{f}_{i-k}). \end{aligned}$$

Now, we have

$$\begin{aligned}
 \|\mathbb{Y}_n - \mathbf{G}_n\| &= \left\| M^n A^{-n} \Psi_0 + M^{n-1} A^{-n} \sum_{i=0}^k M^{-i} A^i (N \Psi_{i-k} + \mathbb{F}(i, \Psi_{i-k})) \right. \\
 &\quad \left. + M^{n-1} A^{-n} \sum_{i=k+1}^n M^{-i} A^i (N \mathbb{Y}_{i-k} + \mathbb{F}(i, \mathbb{Y}_{i-k}) + \mathbf{f}_{i-k}) \right. \\
 &\quad \left. - M^n A^{-n} \Psi_0 - M^{n-1} A^{-n} \sum_{i=0}^k M^{-i} A^i (N \Psi_{i-k} + \mathbb{F}(i, \Psi_{i-k})) \right. \\
 &\quad \left. - M^{n-1} A^{-n} \sum_{i=k+1}^n M^{-i} A^i (N \mathbf{G}_{i-k} + \mathbb{F}(i, \mathbf{G}_{i-k})) \right\|, \\
 \|\mathbb{Y}_n - \mathbf{G}_n\| &= \left\| M^{n-1} A^{-n} \sum_{i=k+1}^n M^{-i} A^i (N \mathbb{Y}_{i-k} + \mathbb{F}(i, \mathbb{Y}_{i-k}) + \mathbf{f}_{i-k}) \right. \\
 &\quad \left. - M^{n-1} A^{-n} \sum_{i=k+1}^n M^{-i} A^i (N \mathbf{G}_{i-k} + \mathbb{F}(i, \mathbf{G}_{i-k})) \right\|, \\
 \|\mathbb{Y}_n - \mathbf{G}_n\| &\leq \|M\|^{n-1} \|A\|^{-n} \sum_{i=k+1}^n \|M\|^{-i} \|A\|^i (\|N \mathbb{Y}_{i-k} - N \mathbf{G}_{i-k}\| \\
 &\quad + \|\mathbb{F}(i, \mathbb{Y}_{i-k}) - \mathbb{F}(i, \mathbf{G}_{i-k})\| + \|\mathbf{f}_{i-k}\|), \\
 \|\mathbb{Y}_n - \mathbf{G}_n\| &\leq \|M\|^{n-1} \|A\|^{-n} \sum_{i=k+1}^n \|M\|^{-i} \|A\|^i (\|N \mathbb{Y}_{i-k} - N \mathbf{G}_{i-k}\| \\
 &\quad + K \|\mathbb{Y}_{i-k} - \mathbf{G}_{i-k}\| + \|\mathbf{f}_{i-k}\|) \\
 &= \|M\|^{n-1} \|A\|^{-n} \sum_{j=k+1}^n \|M\|^{-i} \|A\|^i (\|(N + K) \mathbb{Y}_{i-k} \\
 &\quad - (N + K) \mathbf{G}_{i-k}\| + \|\mathbf{f}_{i-k}\|) \\
 &\leq \|M\|^{n-1} \|A\|^{-n} \sum_{i=k+1}^n \|M\|^{-i} \|A\|^i \|\mathbf{f}_{i-k}\| \\
 &\leq \|M\|^{n-1} \|A\|^{-n} \sum_{i=k+1}^n \|M\|^{-i} \|A\|^i \epsilon \phi_{i-k} \\
 &= \epsilon \|M\|^{n-1} \|A\|^{-n} \sum_{r=1}^{n-k} \|M\|^{-k-r} \|A\|^{k+r} \phi_r \\
 &= \epsilon L^4 \sum_{r=1}^{n-k} \phi_r \\
 &\leq \epsilon L^4 \eta_\varphi \varphi_n \\
 &= K \epsilon.
 \end{aligned}$$

Therefore, system (1.1) is Hyers–Ulam stable over a bounded discrete interval. \square

5 Hyers–Ulam stability on unbounded discrete interval

To discuss the Hyers–Ulam stability over an unbounded discrete interval, we have the following assumptions:

A_1 : The operator family $\|L^4\| \leq Ne^{-\nu n}$, $n \geq 0$, $\nu \geq 0$, $N \geq 1$.

A_2 : The linear system $\mathbf{A}\mathbf{G}_{n+1} = \mathbf{M}\mathbf{G}_n + \mathbf{N}\mathbf{G}_{n-k}$ is well posed.

A_3 : The continuous function $\mathbb{H}: \mathbb{Z}_+ \times X \rightarrow X$ satisfies the Caratheodory condition

$$\|\mathbb{H}(n, \omega) - \mathbb{H}(n, \omega')\| \leq K\|\omega - \omega'\|, \quad K \geq 0,$$

for every $n \in \mathbb{Z}_+$, $\omega, \omega' \in X$.

A_4 : Also, assume that

$$\sum_{r=1}^{n-1} \phi_r \leq \eta \varphi_n$$

for each $n \in \mathbb{Z}_+$, $\eta \geq 0$ and φ_n is a convergent sequence.

Theorem 3 *If A_1 – A_4 along with (2.1) and Remark 2.1 are satisfied, then system (1.1) is Hyers–Ulam stable over an unbounded interval.*

Proof Since the exact solution of the non-autonomous and nonsingular delay difference equation

$$\begin{cases} \mathbf{A}U_{n+1} = \mathcal{M}U_n + \mathbb{N}U_{n-k} + \mathbf{F}(n, U_{n-k}), & n \geq 0, k \geq 0, \\ U_n = \psi_n, & -k \leq n \leq 0, \end{cases}$$

is

$$\begin{aligned} U_n &= \mathcal{M}^n \mathbf{A}^{-n} \psi_0 + \mathcal{M}^{n-1} \mathbf{A}^{-n} \sum_{i=0}^k \mathcal{M}^{-i} \mathbf{A}^i (\mathbb{N} \psi_{i-k} + \mathbf{F}(i, \psi_{i-k})) \\ &\quad + \mathcal{M}^{n-1} \mathbf{A}^{-n} \sum_{i=k+1}^n \mathcal{M}^{-i} \mathbf{A}^i (\mathbb{N} U_{i-k} + \mathbf{F}(i, U_{i-k})). \end{aligned}$$

Letting Y be the approximate solution of the above system, then clearly for a sequence f_n with $\|f_n\| \leq \epsilon$ we have

$$\begin{cases} \mathbf{A}Y_{n+1} = \mathcal{M}Y_n + \mathbb{N}Y_{n-k} + \mathbf{F}(n, Y_{n-k}) + f_n, & n \geq 0, k \geq 0, \\ Y_n = \psi_n, & -k \leq n \leq 0, \end{cases}$$

and

$$\begin{aligned} Y_n &= \mathcal{M}^n \mathbf{A}^{-n} \psi_0 + \mathcal{M}^{n-1} \mathbf{A}^{-n} \sum_{i=0}^k \mathcal{M}^{-i} \mathbf{A}^i (\mathbb{N} \psi_{i-k} + \mathbf{F}(i, \psi_{i-k})) \\ &\quad + \mathcal{M}^{n-1} \mathbf{A}^{-n} \sum_{i=k+1}^n \mathcal{M}^{-i} \mathbf{A}^i (\mathbb{N} Y_{i-k} + \mathbf{F}(i, Y_{i-k}) + f_{i-k}). \end{aligned}$$

Now, consider

$$\begin{aligned}\|Y_n - U_n\| &= \left\| \mathcal{M}^n \mathbf{A}^{-n} \psi_0 + \mathcal{M}^{n-1} \mathbf{A}^{-n} \sum_{i=0}^k \mathcal{M}^{-i} \mathbf{A}^i (\mathbb{N} \psi_{i-k} + \mathbf{F}(i, \psi_{i-k})) \right. \\ &\quad + \mathcal{M}^{n-1} \mathbf{A}^{-n} \sum_{i=k+1}^n \mathcal{M}^{-i} \mathbf{A}^i (\mathbb{N} Y_{i-k} + \mathbf{F}(i, Y_{i-k}) + \mathbf{f}_{i-k}) \\ &\quad - \mathcal{M}^n \mathbf{A}^{-n} \psi_0 - \mathcal{M}^{n-1} \mathbf{A}^{-n} \sum_{i=0}^k \mathcal{M}^{-i} \mathbf{A}^i (\mathbb{N} \psi_{i-k} + \mathbf{F}(i, \psi_{i-k})) \\ &\quad \left. - \mathcal{M}^{n-1} \mathbf{A}^{-n} \sum_{i=k+1}^n \mathcal{M}^{-i} \mathbf{A}^i (\mathbb{N} U_{i-k} + \mathbf{F}(i, U_{i-k})) \right\|.\end{aligned}$$

That is,

$$\begin{aligned}\|Y_n - U_n\| &= \|\mathcal{M}\|^{n-1} \|\mathbf{A}\|^{-n} \left\| \sum_{i=k+1}^n \mathcal{M}^{-i} \mathbf{A}^i (\mathbb{N} Y_{i-k} + \mathbf{F}(i, Y_{i-k}) - \mathbb{N} U_{i-k} - \mathbf{F}(i, U_{i-k})) \right. \\ &\quad \left. + \sum_{i=k+1}^n \mathcal{M}^{-i} \mathbf{A}^i \mathbf{f}_{i-k} \right\| \\ &\leq \|\mathcal{M}\|^{n-1} \|\mathbf{A}\|^{-n} \sum_{i=k+1}^n \|\mathcal{M}\|^{-i} \|\mathbf{A}\|^i (\|\mathbb{N} Y_{i-k} - \mathbb{N} U_{i-k}\| \\ &\quad + \|\mathbf{F}(i, Y_{i-k}) - \mathbf{F}(i, U_{i-k})\| + \|\mathbf{f}_{i-k}\|) \\ &\leq \|\mathcal{M}\|^{n-1} \|\mathbf{A}\|^{-n} \sum_{i=k+1}^n \|\mathcal{M}\|^{-i} \|\mathbf{A}\|^i (\|\mathbb{N} Y_{i-k} - \mathbb{N} U_{i-k}\| + K \|Y_{i-k} - U_{i-k}\| \\ &\quad + \|\mathbf{f}_{i-k}\|) \\ &= \|\mathcal{M}\|^{n-1} \|\mathbf{A}\|^{-n} \sum_{i=k+1}^n \|\mathcal{M}\|^{-i} \|\mathbf{A}\|^i (\|(\mathbb{N} + K) Y_{i-k} \\ &\quad - (\mathbb{N} + K) U_{i-k}\| + \|\mathbf{f}_{i-k}\|) \\ &= \|\mathcal{M}\|^{n-1} \|\mathbf{A}\|^{-n} \sum_{i=k+1}^n \|\mathcal{M}\|^{-i} \|\mathbf{A}\|^i \|\mathbf{f}_{i-k}\|.\end{aligned}$$

That is,

$$\begin{aligned}\|Y_n - U_n\| &\leq \|\mathcal{M}\|^{n-1} \|\mathbf{A}\|^{-n} \sum_{i=k+1}^n \|\mathcal{M}\|^{-i} \|\mathbf{A}\|^i \epsilon \phi_{i-k} \\ &= \epsilon \|\mathcal{M}\|^{n-1} \|\mathbf{A}\|^{-n} \sum_{r=1}^{n-k} \|\mathcal{M}\|^{-k-r} \|\mathbf{A}\|^{k+r} \phi_r \\ &= \epsilon L^4 \sum_{r=1}^{n-k} \phi_r \\ &\leq \epsilon L^4 \eta \varphi_n \\ &\leq N e^{-\nu n} \eta K \epsilon\end{aligned}$$

$$\begin{aligned} &\leq N\eta K\epsilon \\ &= K\epsilon, \end{aligned}$$

where $K = N\eta K$. Thus, system (1.1) is Hyers–Ulam stable over an unbounded discrete interval. \square

6 β -Hyers–Ulam stability

To describe β -Hyers–Ulam stability over an unbounded interval, we needed some assumptions:

A_0 : The operator family $\|L^k\| \leq \mathbb{N}e^{kn}$, $n \geq 0$, $k \leq 0$, $N \geq 1$.

A_1 : The linear system $A\mathbf{G}_{n+1} = M\mathbf{G}_n + N\mathbf{G}_{n-k}$ is well posed.

A_2 : The continuous function $\mathbf{H}: \mathbb{Z}_+ \times X \rightarrow X$ satisfies the Caratheodory condition

$$\|\mathbf{H}(n, \rho) - \mathbf{H}(n, \rho')\| \leq K_n \|\rho - \rho'\|, \quad K \geq 0,$$

for every $n \in \mathbb{Z}_+$, $\rho, \rho' \in X$.

A_3 : Also, assume that

$$\sum_{i=k+1}^n e^{kn} (N + K_n) \leq \eta_\varphi \varphi_n, \quad k \leq 0,$$

for each $n \in \mathbb{Z}_+$, $\eta_\varphi \geq 0$, and φ_n is a convergent sequence.

By considering inequality (2.1) and the above mentioned assumptions, we are able to prove the following theorem.

Theorem 4 *If A_0 – A_3 are satisfied, then system (6.1) is β -Hyers–Ulam stable over an unbounded interval.*

Proof The only one solution of nonsingular delay difference equation

$$\begin{cases} \mathcal{A}\mathcal{U}_{n+1} = \mathcal{M}\mathcal{U}_n + \mathcal{N}\mathcal{U}_{n-k} + F(n, \mathcal{U}_{n-k}), & n \geq 0, k \geq 0, \\ \mathcal{U}_n = \Phi_n, & -k \leq n \leq 0, \end{cases} \quad (6.1)$$

is

$$\begin{aligned} \mathcal{U}_n &= \mathcal{M}^n \mathcal{A}^{-n} \Phi_0 + \mathcal{M}^{n-1} \mathcal{A}^{-n} \sum_{l=0}^k \mathcal{M}^{-l} \mathcal{A}^l (\mathcal{N} \Phi_{l-k} + F(l, \Phi_{l-k})) \\ &\quad + \mathcal{M}^{n-1} \mathcal{A}^{-n} \sum_{l=k+1}^n \mathcal{M}^{-l} \mathcal{A}^l (\mathcal{N} \mathcal{U}_{l-k} + F(l, \mathcal{U}_{l-k})). \end{aligned}$$

Let \mathcal{Y} satisfy (2.1), then for every $n \in \mathbb{Z}_+$ we have

$$\begin{aligned} &\left\| \mathcal{Y}_n - \mathcal{M}^n \mathcal{A}^{-n} \Phi_0 - \mathcal{M}^{n-1} \mathcal{A}^{-n} \sum_{l=0}^k \mathcal{M}^{-l} \mathcal{A}^l (\mathcal{N} \Phi_{l-k} + F(l, \Phi_{l-k})) \right. \\ &\quad \left. - \mathcal{M}^{n-1} \mathcal{A}^{-n} \sum_{l=k+1}^n \mathcal{M}^{-l} \mathcal{A}^l (\mathcal{N} \mathcal{Y}_{l-k} + F(l, \mathcal{Y}_{l-k})) \right\| \end{aligned}$$

$$\begin{aligned}
 &= \left\| \mathcal{M}^{n-1} \mathcal{A}^{-n} \sum_{l=k+1}^n \mathcal{M}^{-l} \mathcal{A}^l f_{l-k} \right\| \\
 &= \left\| \mathcal{M}^{n-1} \right\| \left\| \mathcal{A}^{-n} \right\| \sum_{l=k+1}^n \left\| \mathcal{M}^{-l} \right\| \left\| \mathcal{A}^l \right\| \|f_{l-k}\| \\
 &= \sum_{l=k+1}^n L^4 \epsilon \phi_n \\
 &= \sum_{l=k+1}^n \mathbb{N} e^{kn} \epsilon \phi_n.
 \end{aligned}$$

Now,

$$\begin{aligned}
 \|\mathcal{Y}_n - \mathcal{U}_n\|^\beta &= \left\| \mathcal{Y}_n - \mathcal{M}^n \mathcal{A}^{-n} \Phi_0 - \mathcal{M}^{n-1} \mathcal{A}^{-n} \sum_{i=0}^k \mathcal{M}^{-l} \mathcal{A}^l (\mathcal{N} \Phi_{l-k} + \mathbb{F}(l, \Phi_{l-k})) \right. \\
 &\quad \left. - \mathcal{M}^{n-1} \mathcal{A}^{-n} \sum_{l=k+1}^n \mathcal{M}^{-l} \mathcal{A}^l (\mathcal{N} \mathcal{U}_{l-k} + \mathbb{F}(l, \mathcal{U}_{l-k})) \right\|^\beta, \\
 \|\mathcal{Y}_n - \mathcal{U}_n\|^\beta &= \left\| \mathcal{Y}_n - \mathcal{M}^n \mathcal{A}^{-n} \Phi_0 - \mathcal{M}^{n-1} \mathcal{A}^{-n} \sum_{l=0}^k \mathcal{M}^{-l} \mathcal{A}^l (\mathcal{N} \Phi_{l-k} + \mathbb{F}(l, \Phi_{l-k})) \right. \\
 &\quad - \mathcal{M}^{n-1} \mathcal{A}^{-n} \sum_{l=k+1}^n \mathcal{M}^{-l} \mathcal{A}^l (\mathcal{N} \mathcal{U}_{l-k} + \mathbb{F}(l, \mathcal{U}_{l-k})) \\
 &\quad + \mathcal{M}^{n-1} \mathcal{A}^{-n} \sum_{l=k+1}^n \mathcal{M}^{-l} \mathcal{A}^l (\mathcal{N} \mathcal{Y}_{l-k} + \mathbb{F}(l, \mathcal{Y}_{l-k})) \\
 &\quad \left. - \mathcal{M}^{n-1} \mathcal{A}^{-n} \sum_{l=k+1}^n \mathcal{M}^{-l} \mathcal{A}^l (\mathcal{N} \mathcal{Y}_{l-k} + \mathbb{F}(l, \mathcal{Y}_{l-k})) \right\|^\beta \\
 &= \left\| \mathcal{Y}_n - \mathcal{M}^n \mathcal{A}^{-n} \Phi_0 - \mathcal{M}^{n-1} \mathcal{A}^{-n} \sum_{l=0}^k \mathcal{M}^{-l} \mathcal{A}^l (\mathcal{N} \Phi_{l-k} + \mathbb{F}(l, \Phi_{l-k})) \right. \\
 &\quad - \mathcal{M}^{n-1} \mathcal{A}^{-n} \sum_{l=k+1}^n \mathcal{M}^{-l} \mathcal{A}^l (\mathcal{N} \mathcal{Y}_{l-k} + \mathbb{F}(l, \mathcal{Y}_{l-k})) \\
 &\quad + \mathcal{M}^{n-1} \mathcal{A}^{-n} \sum_{l=k+1}^n \mathcal{M}^{-l} \mathcal{A}^l (\mathcal{N} \mathcal{Y}_{l-k} + \mathbb{F}(l, \mathcal{Y}_{l-k})) \\
 &\quad \left. - \mathcal{M}^{n-1} \mathcal{A}^{-n} \sum_{l=k+1}^n \mathcal{M}^{-l} \mathcal{A}^l (\mathcal{N} \mathcal{U}_{l-k} + \mathbb{F}(l, \mathcal{U}_{l-k})) \right\|^\beta, \\
 \|\mathcal{Y}_n - \mathcal{U}_n\|^\beta &\leq \left\| \mathcal{Y}_n - \mathcal{M}^n \mathcal{A}^{-n} \Phi_0 - \mathcal{M}^{n-1} \mathcal{A}^{-n} \sum_{l=0}^k \mathcal{M}^{-l} \mathcal{A}^l (\mathcal{N} \Phi_{l-k} + \mathbb{F}(l, \Phi_{l-k})) \right. \\
 &\quad \left. - \mathcal{M}^{n-1} \mathcal{A}^{-n} \sum_{l=k+1}^n \mathcal{M}^{-l} \mathcal{A}^l (\mathcal{N} \mathcal{Y}_{l-k} + \mathbb{F}(l, \mathcal{Y}_{l-k})) \right\|^\beta
 \end{aligned}$$

$$\begin{aligned}
& + \left\| \mathcal{M}^{n-1} \mathcal{A}^{-n} \sum_{l=k+1}^n \mathcal{M}^{-l} \mathcal{A}^l (\mathcal{N} \mathcal{Y}_{l-k} + \mathbf{F}(l, \mathcal{Y}_{l-k})) \right. \\
& \left. - \mathcal{M}^{n-1} \mathcal{A}^{-n} \sum_{l=k+1}^n \mathcal{M}^{-l} \mathcal{A}^l (\mathcal{N} \mathcal{U}_{l-k} + \mathbf{F}(l, \mathcal{U}_{l-k})) \right\|^\beta \\
& \leq \left(\sum_{l=k+1}^n \mathbb{N} e^{kn} \epsilon \phi_n \right)^\beta + \left(\left\| \mathcal{M}^{n-1} \right\| \left\| \mathcal{A}^{-n} \right\| \sum_{l=k+1}^n \left\| \mathcal{M}^{-l} \right\| \left\| \mathcal{A}^l \right\| \left\| \mathcal{N} \mathcal{Y}_{l-k} \right. \right. \\
& \quad \left. \left. + \mathbf{F}(l, \mathcal{Y}_{l-k}) - \mathcal{N} \mathcal{U}_{l-k} - \mathbf{F}(l, \mathcal{U}_{l-k}) \right\| \right)^\beta, \\
\| \mathcal{Y}_n - \mathcal{U}_n \|^\beta & = \left(\sum_{l=k+1}^n \mathbb{N} e^{kn} \epsilon \phi_n \right)^\beta \\
& + \left(\left\| \mathcal{M}^{n-1} \right\| \left\| \mathcal{A}^{-n} \right\| \sum_{l=k+1}^n \left\| \mathcal{M}^{-l} \right\| \left\| \mathcal{A}^l \right\| \left(\left\| \mathcal{N} \mathcal{Y}_{l-k} - \mathcal{N} \mathcal{U}_{l-k} \right\| \right. \right. \\
& \quad \left. \left. + \left\| \mathbf{F}(l, \mathcal{Y}_{l-k}) - \mathbf{F}(l, \mathcal{U}_{l-k}) \right\| \right) \right)^\beta \\
& \leq \left(\sum_{l=k+1}^n \mathbb{N} e^{kn} \epsilon \phi_n \right)^\beta \\
& + \left(\left\| \mathcal{M}^{n-1} \right\| \left\| \mathcal{A}^{-n} \right\| \sum_{l=k+1}^n \left\| \mathcal{M}^{-l} \right\| \left\| \mathcal{A}^l \right\| \left(\left\| \mathcal{N} \mathcal{Y}_{l-k} - \mathcal{N} \mathcal{U}_{l-k} \right\| \right. \right. \\
& \quad \left. \left. + K_n \left\| \mathcal{Y}_{l-k} - \mathcal{U}_{l-k} \right\| \right) \right)^\beta \\
& = \left(\sum_{l=k+1}^n \mathbb{N} e^{kn} \epsilon \phi_n \right)^\beta + \left(\sum_{l=k+1}^n L^4 (N + K_n) \left\| \mathcal{Y}_{l-k} - \mathcal{U}_{l-k} \right\| \right)^\beta \\
& \leq \left(\sum_{l=k+1}^n \mathbb{N} e^{kn} \epsilon \phi_n \right)^\beta + \left(\sum_{l=k+1}^n \mathbb{N} e^{kn} (N + K_n) \left\| \mathcal{Y}_{l-k} - \mathcal{U}_{l-k} \right\| \right)^\beta, \\
\| \mathcal{Y}_n - \mathcal{U}_n \| & = \left[\left(\sum_{l=k+1}^n \mathbb{N} e^{kn} \epsilon \phi_n \right)^\beta + \left(\sum_{l=k+1}^n \mathbb{N} e^{kn} (N + K_n) \left\| \mathcal{Y}_{l-k} - \mathcal{U}_{l-k} \right\| \right)^\beta \right]^{\frac{1}{\beta}}.
\end{aligned}$$

Now, using

$$(\varrho + \xi)^\alpha \leq 3^{\alpha-1} (\varrho^\alpha + \xi^\alpha), \quad \varrho, \xi \geq 0 \text{ and } \alpha > 1,$$

we obtain

$$\| \mathcal{Y}_n - \mathcal{U}_n \| \leq 3^{\frac{1}{\beta}-1} \sum_{l=k+1}^n \mathbb{N} e^{kn} \epsilon \phi_n + 3^{\frac{1}{\beta}-1} (N + K_n) \sum_{l=k+1}^n \mathbb{N} e^{kn} \left\| \mathcal{Y}_{l-k} - \mathcal{U}_{l-k} \right\|,$$

with the help of Lemma(2.2), we get

$$\begin{aligned}\|\mathcal{Y}_n - \mathcal{U}_n\| &\leq 3^{\frac{1}{\beta}-1} \sum_{l=k+1}^n \mathbb{N}e^{kn} \epsilon \phi_n \exp\left(3^{\frac{1}{\beta}-1} \sum_{l=k+1}^n \mathbb{N}e^{kn} (N + K_n)\right), \\ \|\mathcal{Y}_n - \mathcal{U}_n\| &\leq \epsilon \left(3^{\frac{1}{\beta}-1} \sum_{l=k+1}^n \mathbb{N}e^{kn}\right) \exp\left(3^{\frac{1}{\beta}-1} \sum_{l=k+1}^n \mathbb{N}e^{kn} (N + K_n)\right) \\ &= \epsilon \left(3^{\frac{1}{\beta}-1} \sum_{l=k+1}^n \mathbb{N}e^{kn}\right) \exp\left(3^{\frac{1}{\beta}-1} \mathbb{N}e^{kn} \eta_\varphi \varphi_n\right), \\ \|\mathcal{Y}_n - \mathcal{U}_n\|^\beta &= \epsilon^\beta \left(3^{\frac{1}{\beta}-1} \sum_{l=k+1}^n \mathbb{N}e^{kn}\right)^\beta \exp\left(3^{\frac{1}{\beta}-1} \mathbb{N}e^{kn} \eta_\varphi \varphi_n\right)^\beta \\ &= \epsilon^\beta \left(3^{\frac{1}{\beta}-1} \sum_{l=k+1}^n \mathbb{N}e^{kn}\right)^\beta (\exp(\mathbb{N}3^{\frac{1}{\beta}-1}))^\beta (\exp(\eta_\varphi \varphi_n))^\beta \\ &= L_{F, \mathbb{N}, \phi, \beta} \epsilon^\beta \eta_\varphi^\beta \varphi_n^\beta,\end{aligned}$$

where

$$L_{F, \mathbb{N}, \phi, \beta} = \left(3^{\frac{1}{\beta}-1} \sum_{l=k+1}^n \mathbb{N}e^{kn}\right)^\beta (\exp(\mathbb{N}3^{\frac{1}{\beta}-1}))^\beta.$$

So, system (6.1) is β -Hyers–Ulam stable over an unbounded interval. \square

7 An example

Consider we have the following nonsingular delay difference equation:

$$\begin{cases} \mathcal{AG}_{n+1} = \mathcal{MG}_n + \mathcal{NG}_{n-3} + F(n, \mathcal{G}_{n-3}), & \mathcal{G}_0 = 1, n \in \{0, 1, 2, 3\}, \\ \mathcal{G}_n = \Phi_n, & -3 \leq n \leq 0, \end{cases} \quad (7.1)$$

with inequality

$$\begin{cases} \|\mathcal{AG}_{n+1} - \mathcal{MG}_n - \mathcal{NG}_{n-3} - F(n, \mathcal{G}_{n-3})\| \leq 0.7, & n \in \{0, 1, 2, 3\}, \\ \|\mathcal{G}_n - \Phi_n\| \leq 1, & -3 \leq n \leq 0, \end{cases} \quad (7.2)$$

here $k = 3$. If we fix

$$\mathcal{M} = \begin{pmatrix} -3.5 & 1.4 \\ 3.2 & 1.6 \end{pmatrix}, \quad \mathcal{N} = \begin{pmatrix} 3.6 & 1.4 \\ -3.2 & 1.7 \end{pmatrix}, \quad \mathcal{A} = \begin{pmatrix} 1.3 & 0 \\ 0 & 1.3 \end{pmatrix},$$

$F(n, \mathcal{G}_{n-3}) = \mathcal{G}_{n-3}[0.3 \sin(n) \quad 0.15 \sin(n)]$ and $\phi_n = [\cos(n + \frac{\pi}{2}) \quad \cos(n + \frac{\pi}{2})]^t$ (obviously, $\phi_n = [0 \quad 0]^t$, when $n = 0$), hence, we get

$$\mathcal{NM} = \begin{pmatrix} 8.12 & 7.28 \\ -16.64 & -1.76 \end{pmatrix} = \mathcal{MN}, \quad \mathcal{AN} = \begin{pmatrix} 4.68 & 1.28 \\ -4.16 & 2.12 \end{pmatrix} = \mathcal{NA},$$

$$\begin{aligned}\mathcal{A}\mathcal{M} &= \begin{pmatrix} 4.55 & 1.82 \\ -4.16 & 2.08 \end{pmatrix} = \mathcal{M}\mathcal{A}, & \mathcal{A}^{-1} &= \begin{pmatrix} 0.769231 & 0 \\ 0 & 0.769231 \end{pmatrix}, \\ \mathcal{A}^{-1}\mathcal{N} &= \begin{pmatrix} 2.762316 & 1.0769234 \\ -2.4615392 & 1.3076927 \end{pmatrix} = \mathcal{N}\mathcal{A}^{-1}, \\ \mathcal{A}^{-1}\mathcal{M} &= \begin{pmatrix} 2.6923085 & 1.0769234 \\ -2.4615392 & 1.2307696 \end{pmatrix} = \mathcal{M}\mathcal{A}^{-1}.\end{aligned}$$

Moreover, if G satisfies (7.2), then there exists f_n such that $\|f_n\| \leq 0.7$, and

$$\begin{cases} \mathcal{A}\mathcal{G}_{n+1} = \mathcal{M}\mathcal{G}_n + \mathcal{N}\mathcal{G}_{n-3} + F(n, \mathcal{G}_{n-3}) + f_n, & \mathcal{G}_0 = 1, n \in \{0, 1, 2, 3\}, \\ \mathcal{G}_n = \Phi_n, & -3 \leq n \leq 0, \end{cases}$$

also the solution of (7.1) is

$$\begin{aligned}\mathcal{G}_n &= \mathcal{M}^n \mathcal{A}^{-n} \Phi_0 + \mathcal{M}^{n-1} \mathcal{A}^{-n} \sum_{i=0}^k \mathcal{M}^{-i} \mathcal{A}^i (\mathcal{N} \Phi_{i-k} + F(i, \Phi_{i-k})) \\ &\quad + \mathcal{M}^{n-1} \mathcal{A}^{-n} \sum_{i=k+1}^n \mathcal{M}^{-i} \mathcal{A}^i (\mathcal{N} \mathcal{G}_{i-k} + F(i, \mathcal{G}_{i-k})),\end{aligned}\tag{7.3}$$

where $\mathcal{M}\mathcal{A} = \mathcal{A}\mathcal{M}$, $\mathcal{N}\mathcal{A} = \mathcal{A}\mathcal{N}$, and $\mathcal{M}\mathcal{N} = \mathcal{N}\mathcal{M}$.

Let $\epsilon = 0.7$, and $f : \mathcal{Z}_+ \rightarrow \mathcal{R}^2$ be as given below

$$f_n = \left[0.6 \cos\left(n + \frac{\pi}{2}\right) \quad 0.6 \sin\left(n + \frac{\pi}{2}\right) \right]^t,$$

then clearly

$$\begin{aligned}\|f_n\| &= \sqrt{\left(0.6 \cos\left(n + \frac{\pi}{2}\right)\right)^2 + \left(0.6 \sin\left(n + \frac{\pi}{2}\right)\right)^2} \\ &= \left[(0.6)^2 \cos^2\left(n + \frac{\pi}{2}\right) + (0.6)^2 \sin^2\left(n + \frac{\pi}{2}\right) \right]^{\frac{1}{2}} \\ &= \sqrt{0.6^2} = 0.6 \\ &\leq \epsilon = 0.7.\end{aligned}$$

Now the perturbed delay difference system (7.3) has the solution


$$\begin{aligned}H_n &= \mathcal{M}^n \mathbf{A}^{-n} \psi_0 + \mathcal{M}^{n-1} \mathbf{A}^{-n} \sum_{i=0}^k \mathcal{M}^{-i} \mathbf{A}^i (\mathbb{N} \psi_{i-k} + F(i, \psi_{i-k})) \\ &\quad + \mathcal{M}^{n-1} \mathbf{A}^{-n} \sum_{i=k+1}^n \mathcal{M}^{-i} \mathbf{A}^i (\mathbb{N} H_{i-k} + F(i, H_{i-k}) + \mathbf{f}_{i-k}).\end{aligned}$$

Using Mathematica, we get the values given in Table 1.

Table 1 Table for G_n and H_n

n	G_n	H_n	$\ G_n - H_n\ $
1.	$a_1 = 1.11268$	$a_1 = 1.51268$	0.4
	$b_1 = 1.38337$	$b_1 = 1.64668$	0.534
2.	$a_2 = 4.48545$	$a_2 = 4.92105$	0.4356
	$b_2 = 3.03628$	$b_2 = 3.46195$	0.42567
3.	$a_3 = 10.9602$	$a_3 = 11.5274$	0.567234
	$b_3 = 8.3165$	$b_3 = 8.92885$	0.61235

Plotting these values, we have the following graphs.



Hence, we have a solution within a multiple of $\epsilon = 0.7$ and a constant, so system (1.1) has a unique solution in $B(\mathcal{Z}_+, \mathcal{R}^2)$, which is Hyers–Ulam stable on \mathcal{Z}_+ .

Acknowledgements

The authors are grateful to the editor and anonymous referees for their careful review and valuable comments which helped to improve this manuscript. The authors T. Abdeljawad and A. Mukheimer would like to thank Prince Sultan University for paying the publications charges and for the support through the TAS research Lab.

Funding

The authors T. Abdeljawad and A. Mukheimer would like to thank Prince Sultan University for paying the publications charges.

Availability of data and materials

No data were used.

Declarations

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally to the writing of this manuscript. All authors read and approved the final manuscript.

Author details

¹Department of Mathematics, Islamia College University Peshawar, Khyber Pakhtoonkhwa, Pakistan. ²Department of Mathematics, University of Malakand, Chakdara, Dir(L), Pakistan. ³Department of Mathematics and Sciences, Prince Sultan University, P.O. Box 66833, Riyadh 11586, Saudi Arabia. ⁴Department of Medical Research, China Medical University, Taichung, Taiwan. ⁵Department of Computer Science and Information Engineering, Asia University, Taichung, Taiwan.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 12 May 2021 Accepted: 10 August 2021 Published online: 26 October 2021

References

1. Frassu, S., Viglialoro, G.: Boundedness for a fully parabolic Keller–Segel model with sublinear segregation and superlinear aggregation. *Acta Appl. Math.* **171**(1), 19 (2021)
2. Li, T., Pintus, N., Viglialoro, G.: Properties of solutions to porous medium problems with different sources and boundary conditions. *Z. Angew. Math. Phys.* **70**(3), 86 (2019)
3. Li, T., Viglialoro, G.: Boundedness for a nonlocal reaction chemotaxis model even in the attraction-dominated regime. *Differ. Integral Equ.* **34**(5/6), 315–336 (2021)
4. Li, T., Viglialoro, G.: Analysis and explicit solvability of degenerate tensorial problems. *Bound. Value Probl.* **2018**(1), 2 (2018)
5. Brianzoni, S., Mammana, C., Michetti, E., Zirilli, F.: A stochastic cobweb dynamical model. *Discrete Dyn. Nat. Soc.* **2008**, 219653 (2008)
6. Sarkar, B., Mondal, S.P., Hur, S., Ahmadian, A., Salahshour, S., Guchhait, R., Iqbal, M.W.: An optimization technique for national income determination model with stability analysis of differential equation in discrete and continuous process under the uncertain environment. *RAIRO Oper. Res.* **53**(5), 1649–1674 (2019)
7. Kuruklis, S.A.: The asymptotic stability of difference equation. *J. Math. Anal. Appl.* **188**(3), 719–731 (1994)
8. Yu, J.S.: Asymptotic stability for a linear difference equation with variable delay. *Comput. Math. Appl.* **36**(10–12), 203–210 (1998)
9. Kosmala, W., Teixeira, C.: More on the difference equation. *Appl. Anal.* **81**(1), 143–151, 81.1 (2002)
10. Fan, L.Z., Cai, C.X., Zou, Y.: Switching signal design for exponential stability of uncertain discrete-time switched time-delay systems. *J. Appl. Math.* **2013**, 416292 (2013)
11. Marwen, K., Sakly, A.: On stability analysis of discrete-time uncertain switched nonlinear time-delay systems. *Adv. Differ. Equ.* **2014**(1), 1 (2014)
12. Yuan, L., Meng, F.: Stability analysis of a class of higher order difference equations. *Abstr. Appl. Anal.* **2014**, 434621 (2014)
13. Khaliq, A., et al.: On stability analysis of higher-order rational difference equation. *Discrete Dyn. Nat. Soc.* **2020**, 3094185 (2020)
14. Ulam, S.M.: A Collection of Mathematical Problems. Interscience, New York, no. 8 (1960)
15. Hyers, D.H.: On the stability of the linear functional equation. *Proc. Natl. Acad. Sci. USA* **27**(4), 222 (1941)
16. Rassias, T.: On the stability of the linear mapping in Banach spaces. *Proc. Am. Math. Soc.* **72**(2), 297–300 (1978)
17. Obłoz, M.: Connections between Hyers and Lyapunov stability of the ordinary differential equations. *Rocznik Nauk.-Dydakt. Prace Mat.* **14**, 141–146 (1997)
18. Jung, S.M.: Hyers–Ulam stability of the first-order matrix difference equations. *Adv. Differ. Equ.* **2015**(1), 1 (2015)
19. Khan, A., Rahmat, G., Zada, A.: On uniform exponential stability and exact admissibility of discrete semigroups. *Int. J. Differ. Equ.* **2013**, 268309 (2013)
20. Khusainov, D.Ya., Shuklin, G.V.: Linear autonomous time-delay system with permutation matrices solving. *Stud. Univ. Žilina Math. Ser.* **17**(1), 101–108 (2003)
21. Diblík, J., Khusainov, D.Y.: Representation of solutions of discrete delayed system with commutative matrices. *J. Math. Anal. Appl.* **318**(1), 63–76 (2006)
22. You, Z., JinRong, W., O'Regan, D.: Exponential stability and relative controllability of nonsingular delay systems. *Bull. Braz. Math. Soc.* **50**(2), 457–479 (2019)
23. Li, T., Zada, A.: Connections between Hyers–Ulam stability and uniform exponential stability of discrete evolution families of bounded linear operators over Banach spaces. *Adv. Differ. Equ.* **2019**(1), 1 (2016)
24. Shah, S.O., Zada, A., Muzammil, M., Tayyab, M., Rizwan, R.: On the Bielecki–Ulam's type stability results of first order non-linear impulsive delay dynamic systems on time scales. *Qual. Theory Dyn. Syst.* **19**(3), 98 (2020)
25. Shah, S.O., Zada, A., Hamza, A.E.: Stability analysis of the first order non-linear impulsive time varying delay dynamic system on time scales. *Qual. Theory Dyn. Syst.* **18**(3), 825–840 (2019)
26. Zada, A., Pervaiz, B., Alzabut, J., Shah, S.O.: Further results on Ulam stability for a system of first-order nonsingular delay differential equations. *Demonstr. Math.* **53**(1), 225–235 (2020)

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)